



HOW DO REGENERATION STATUS, VEGETATION DIVERSITY, STAND STRUCTURE, AND CARBON STOCK VARY ACROSS ECOLOGICAL REGIONS OF NEPAL?

SANTOSH AYER*, AMAN PRABHAKAR, KISHOR ADHIKARI, SUBRAT POUDEL, SANDHYA BASNET, MANISHA TIMALSINA, MEDHA GIRI, PRASHANT TARAMI MAGAR, SARU LAMICHANE, ANIJA SINGH

*Agriculture and Forestry University, College of Natural Resource Management (CNRM),
Katari, 56310, Nepal*

**Corresponding author: ayer.sant@gmail.com*

Abstract.

*This study compared the regeneration status, vegetation diversity, stand structure, and carbon stock of two community-managed forests located in different ecological regions of Nepal. A total of 61 concentric sample plots were investigated by using systematic random sampling with 1 % sampling intensity. The phytosociological parameters of trees and regeneration density were calculated using standard techniques. The aboveground tree carbon was estimated using a non-destructive method. Correlation analysis was performed to assess the variation of carbon stock with biomass, stand density, tree diameter at breast height (DBH), tree height, basal area, and seedling density. Janata community forest had higher regeneration than the Hazare community forest, with a bell-shaped distribution of DBH in Janata and an interrupted curve in Hazare community forest. Both forests were dominated by *Shorea robusta*. Hazare had higher Shannon diversity index, Simpson's index, and evenness index, while Janata community forest had higher carbon stock. The observed differences in the studied parameters between the two community forests can be attributed to the difference in ecological factors such as temperature, rainfall, soil nutrient availability, and management practices. Future studies focusing on investigating the underlying factors driving the observed patterns and relationships, such as the effects of disturbance, climate, and management practices on forest structure and function are also needed. These findings have important implications for forest management and conservation policies, especially in the context of climate change mitigation and biodiversity conservation.*

Key words: *Sustainable forest management, Vegetation diversity, Carbon sequestration, Biodiversity conservation*

1. INTRODUCTION

Forests are one of the most important terrestrial ecosystems on the planet, providing numerous benefits to both humans and the environment (Brockerhoff et al., 2017). They are home to a diverse range of plant and animal species, many of which are endemic to specific forest types. Forests also provide a wide range of ecosystem services, including climate regulation, water conservation, soil conservation, and carbon sequestration (Bauhus et al., 2010; Thompson et al., 2011; Brockerhoff et al., 2013; Decocq et al., 2016; Liang et al., 2016; Mori et al., 2017). However, forests worldwide are under threat from human activities, such as deforestation, land-use change, and climate change (Liaison, 2013; Thapa, 2021).

In Nepal, community forestry program has been recognized as a model for community-based natural resource management (Ghimire and Lamichhane, 2020). There are 22519 community forests in Nepal, covering 2,312,545 hectares, and 3,088,259 households are in responsible for maintaining them (Pandey and Pokhrel, 2021). This constitutes to 15.67% of the total area of the country, 34.98% of its total forested area, 56.90% of total Households, and 62.68% of the total population of Nepal being engaged in

CF (Pandey and Pokhrel, 2021). Despite the significant number of community forests in Nepal, there is often a lack of detailed information regarding essential technical aspects such as species diversity, stand structure, and carbon stocks (Anup, 2017; Poudyal et al., 2019). These metrics are critical for understanding the role of forests in providing ecosystem services such as carbon sequestration, soil conservation, and biodiversity conservation (Mandal et al., 2013; Bhandari et al., 2021; Baul et al., 2021). Therefore, there is a growing need for assessments studies in community forests in different regions that can provide detailed information about these technical aspects and their implications for ecosystem services (Paudyal et al., 2017).

Regeneration status refers to the ability of a forest to naturally regenerate itself through the growth of new trees and other vegetation (Chazdon and Guariguata, 2016). Assessing the regeneration status of a forest is an important part of forest management and conservation, as it can help identify areas where intervention may be needed to promote natural regeneration and maintain forest health (Rahman et al., 2020). Species diversity is an essential aspect of forest ecosystems, as it reflects the richness and even-

ness of the plant and animal species present in a given area (Magurran, 2005). The diversity of forest ecosystems is a critical indicator of their ecological health and resilience (Brockerhoff et al., 2017). The structure of forest ecosystems, including tree density, size distribution, and canopy cover, also plays an important role in supporting forest biodiversity and ecosystem services (Smith, 2016). The carbon stock of forest ecosystems is a crucial component of the global carbon cycle, as forests act as carbon sinks by sequestering carbon from the atmosphere through the process of photosynthesis (Cannell, 1996; Mukul et al., 2021). The diversity, structure, regeneration status and carbon stock of forests are all critical factors that affect their ability to provide these important ecosystem services (Hooper et al., 2005; Isbell et al., 2011; Gamfeldt et al., 2013). For example, forests with higher levels of biodiversity are typically more resilient to environmental changes and disturbances, and are better able to provide a wide range of ecosystem services (Brockerhoff et al., 2017). Similarly, the structure of forests, including the size, shape, and arrangement of trees, also plays a crucial role in determining the types and amounts of resources that are available to wildlife and humans (Smith, 2016). The composition of forests, including the types and proportions of different tree species, also affects their ability to provide important ecosystem services, such as carbon sequestration and habitat for wildlife (Gamfeldt et al., 2013; Forrester and Bauhus, 2016; Marais et al., 2022). Finally, information on regeneration status can be used to guide management strategies such as tree planting, selective logging, and other interventions aimed at promoting natural regeneration and ensuring the long-term sustainability of the forest ecosystem (Chazdon and Guariguata, 2016; Rahman et al., 2020). Therefore understanding the regeneration status, forest diversity, structure, and carbon stock is essential for effective forest management and conservation.

Biophysical measurements of trees such as number of individuals, diameter at breast height (DBH), height, and crown dimensions can be used to determine diversity and stand structure. Destructive methods such as harvesting trees as well and non-destructive methods such as allometric equations and remote sensing and geographic information system (GIS) can be used to estimate above-ground carbon stock (Shi and Liu, 2017). Allometric equations are based on the relationship between tree biomass and easily measured parameters such as DBH and height (Mugasha et al., 2019). These equations are widely used for estimating forest biomass and carbon stock than harvesting method because they are non-destructive and relatively easy to apply (Shi and Liu, 2017). However, the accuracy of allometric equations can be affected by several factors such as species, forest age, and stand structure (Daba and Soromessa, 2019). Therefore combination of methods such as biophysical measurements and allometric equations, are used in this study.

Although there have been many studies conducted on vegetation analysis and carbon stocks in community forests of Nepal, most of them have focused on a single community forest (Aryal et al., 2018; Sharma et al., 2020; Bhatta and Devkota, 2020; Ghimire and Lamichhane, 2021; Joshi et al., 2021), or compared forests within the same region (Joshi et al., 2021). Additionally, there have been also a few studies that have studied forests under different management regimes (Awasthi et al., 2015; Aryal et al., 2021; Ayer et al., 2022; Khatri et al., 2023). Furthermore, only a limited number of studies have specifically examined differences in vegetation and carbon stocks between different ecological regions (Pandey and Bhusal, 2016). As a result, there is a significant research gap in our understanding in context of Nepal that needs to be filled.

Studying vegetation and carbon stocks in different ecological regions is important because different regions may have different types of forests, different plant species, and different environmental conditions that can affect how much carbon is stored in the vegetation (Pandey and Bhusal, 2016; Karki et al., 2016). Additionally, understanding the differences in vegetation and carbon stocks between ecological regions can help policymakers and land managers make informed decisions about how to manage forests for carbon sequestration and other ecosystem services. Therefore, the purpose of this study is to address this gap in knowledge and assess the diversity, structure, and potential carbon stocks of community forests across two ecological regions.

2. MATERIALS AND METHODS

2.1 Description of the study sites

The study was carried out in the Janata and Hazare community forests (CF) of Katari, Udayapur district of Nepal. It is located in Province no. 1 in the southeastern part of Nepal which lies between 26.8372° N latitude and 86.3213° E longitudes. Both the Mahabharat and Shivalik hills encircle the Udayapur district from the north and the south, and the two hills converge in the west to create the Udayapur valley. The elevation of the district is moderately steep ranging from lower tropical below 300 m and upper tropical ranging from 300-4000 m. Janta CF covers an area of 132 hac and lies in Mahabharat region at 510-770 m whereas Hazare CF covers a range of 177 ha which lies in Chure region at 550-1100 m elevation. Both CF lies in tropical region having 5-50° slope and northeast slope aspect with red, loose, sandy soil. Temperatures in the region exceed 37 °C during the summer period while winter temperatures range from 7 to 23 °C. Leopard (*Panthera pardus*), Rabbit (*Oryctolagus cuniculus*), Barking deer (*Muntiacus muntjak*), Porcupine (*Hystrix brachyura*) etc. are the most spotted animals whereas parrot (*Psittacidae krameri*), sparrow (*Passer montanus*), Common myna (*Acridotheres tristis*), Black Drongo (*Dicrurus macrocerus*) etc. are the most seen bird species in both CF. The




Figure 1: Map of study area

endangered species like Leopard (*Panthera pardus*) and chinese pangolin (*Manis pentadactyla*) are strictly protected in this area.

2.2 Data Collection

GPS and Arc Map 10.8 software were used to demarcate the targeted community forest for the research purpose. The study employed a systematic random sampling approach to gather data on plant species diversity and regeneration status. The plants were classified into three categories based on their size - tree (dbh > 10 cm), sapling (dbh < 10 cm and height > 1.0 m), and seedling (height < 1.0 m). A total of 61 circular sample plots, each covering an area of 500 m² (with a 1% intensity), were established according to the Forest Inventory Guideline recommended by the Government of Nepal (DoF, 2004). Within each sample plot, three concentric plots with radii of 12.62 m, 5.64 m, and 1 m were used to sample trees, saplings, and seedlings, respectively. Diameter at breast height (dbh) and tree height were measured using Diameter Tape and Clinometer respectively.

2.3 Data Analysis

The phytosociological parameters of trees including density (D), frequency (F), basal area (m²), relative density (RD), relative frequency (RF), relative basal area (RBA), and important value index (IVI), were calculated using the Zobel et al. (1987) technique. Similarly, regeneration density (individuals per hac) was also calculated using Zobel et al. (1987). The number of woody species in each plot

was recorded to determine species richness 'S', following Whittaker (1972). The Shannon-diversity Weiner's index (H) (Shannon and Weaver, 1949), Simpson's index 'D' (Simpson, 1949), and Species Evenness (e) (Wilsey et al., 2005) were also evaluated. To analyze the community structure, all plant populations of all species found in all plots were categorized into various size classes based on diameter at breast height (DBH).

In this study, a non-destructive method was used to estimate biomass. The aboveground tree biomass (AGTB) was calculated using the allometric equation provided by Chave et al. (2005), which takes into account the tree diameter at breast height (D), wood specific density (WD) in kg m⁻³, and tree height (H) in meters for trees (dbh > 10 cm).

$$AGTB = 0.0509 \times D^2 \times WD \times H$$

To calculate the carbon stock of individual tree species in community forests, the density values of each species across the entire forest were added together. The percentage contribution of carbon stock of each tree species in a forest was then determined by taking the proportional sum of carbon stock per hectare of all species present in the forest and dividing it by the sum of carbon stock of the particular species in the same forest. This method, described by Joshi et al. (2020), allows for the quantification of the contribution of each tree species to the total carbon stock of a forest and provides insight into the relative importance of different species in carbon sequestration efforts.

Carbon stock of tree species (%) = (Sum of carbon stock of tree species (ha) *100)/ Sum of carbon stock of all tree species (ha)

Correlation analysis was performed to access the variation of C stock with biomass, stand density, tree DBH, tree height, basal area and seedling density. Prior to the correlation analysis, the data were tested for the normality (Shapiro-Wilk test, $p > 0.05$). In Janata CF, tree DBH, tree height and seedling density meet the assumption of normality ($p > 0.05$). So, they were correlated by parametric test i.e. Pearson correlation. Carbon, biomass, tree density and basal area were log transformed to meet the assumption of normality ($p > 0.05$). So, they were correlated by parametric test i.e. Pearson correlation. Similarly, in Hazare CF, carbon, biomass, stand density, tree height, and basal area meet the assumption of normality ($p > 0.05$). So, they were correlated by parametric test i.e. Pearson correlation. Tree DBH and seedling density didn't meet the assumption of normality ($p < 0.05$) even after transformation, so they were correlated by Spearman correlation, a non-parametric test.

3. RESULT AND DISCUSSION

3.1 Regeneration status

A total of 14 species of seedlings and saplings were recorded, with varying degrees of occurrence in the two forest areas. The total number of seedlings (7385 individuals ha⁻¹) and saplings (138 individuals ha⁻¹) in Janata CF was higher than the total number of seedlings (3629 individuals ha⁻¹) and saplings (86 individuals ha⁻¹) in Hazare CF (Table 1). *Shorea robusta*, the dominant species in both forests, was found in higher numbers in Janata CF with 4769 seedlings and 69 saplings per ha, compared to Hazare CF where it had

2200 seedlings and 63 saplings per ha. Other species that were more abundant in Janata CF than Hazare CF included *Symplocos sps*, *Myrsine capitellata*, and *Cassia fistula*. In contrast, Hazare CF had higher abundance of *Schima wallichii* and *Terminalia tomentosa* with 500 ha⁻¹ and 86 seedlings ha⁻¹, respectively, compared to Janata CF where they were not recorded. Additionally, *Alangium salvifolium* and *Buchanania latifolia* were found only in Janata CF, while *Dillenia pentagyna*, *Croton roxburghii*, and *lagerstroemia parviflora* were found only in Hazare CF (Table 1).

Regeneration is a crucial element that shows the community's and species' evolutionary trends and preserves the integrity or resilience of the forest in the long run (Napit 2015; Bose et al. 2016). *Shorea robusta* and *Schima wallichii* have a higher proportion of seedlings and saplings in both community forests, which indicates that they have dominated the regeneration (Table 1). This may be due to the fact that different species have different ecological preferences, and their growth and survival may be affected by factors such as climate, soil type, and topography (Sapkota et al., 2009; Joshi et al., 2021).

Janata CF has good regeneration condition where as Hazare CF has medium regeneration condition according to criteria suggested by Community Forest Resource Inventory guideline (DoF, 2004): a forest is considered to have poor, medium and good regeneration if it has regeneration (Seedlings and saplings) <2000, 2000-5000 and > 5000 in number per hectare (Pandey et al. 2012). However seedlings and sapling density was found lower than Basyal et al. (2011), Awasthi et al. (2015), Napit (2015) and Joshi et al. (2021). This might be due topography, climate, stand intensity, type of disturbances, and soil nutrients of the study site (Khaine et al., 2018; Joshi et al., 2021).

Table 1: Overview of particular species seedlings and saplings regeneration status in Janata and Hazare CFs, Udaypur District, Nepal

Species Name	Hazare CF		Janata CF	
	Individual ha ⁻¹		Individual ha ⁻¹	
	Seedlings	Saplings	Seedlings	Saplings
<i>Alangium salvifolium</i>	0	0	231	0
<i>Buchanania latifolia</i>	0	0	154	12
<i>Cassia fistula</i>	400	0	0	0
<i>Dillenia pentagyna</i>	0	0	0	8
<i>Croton roxburghii</i>	57	0	0	0
<i>lagerstroemia parviflora</i>	143	0	0	0
<i>Lyonia ovalifolia</i>	0	0	154	0
<i>Myrsine capitellata</i>	57	3	0	0
<i>Other hill species</i>	343	3	346	0
<i>Schima wallichii</i>	343	6	500	31
<i>Shorea robusta</i>	2200	63	4769	69
<i>Symplocos sps</i>	0	11	154	19
<i>Syzygium cumini</i>	0	0	1077	0.00
<i>Terminalia tomentosa</i>	86	0	0	0.00
Total	3629	86	7385	138




Figure 2: Distribution of DBH in Janata and Hazare CF.

3.2 DBH distributions

The diameter class-wise distribution of trees in a forest is an important indicator of its structural heterogeneity and can provide insights into forest dynamics and functioning (Lutz et al. 2013). The bell-shaped curve observed in Janata CF with the highest peak in the 30-39.9 cm DBH size class indicates mature nature of forest (Clatterbuck et al., 2011; Sharma et al., 2020) with well-structured forest with a wide range of tree diameters and age classes (Figure 2). The current state of the forest may be a result of past disruptions that have decreased its ability to regenerate, as well as the removal of large trees (Nizami, 2012; Dar et al., 2017). On the other hand, the interrupted curve in Hazare CF with the highest peak in the 20-29.9 cm DBH size class indicates that the forest has gaps in its size structure, which could be problematic to regeneration because not all diameter class are represented (Figure 2) (Temgoua et al., 2020). The interrupted pattern in diameter class distribution observed in the Hazare CF suggests that the forest may have experienced some type of disturbance or disruption in its natural regeneration process. Our result is in contrast with the findings of Chauhan et al. (2008),

Acharya et al. (2007), Basyal et al. (2011), Awasthi et al. (2015), Manna and Mishra (2017) and Joshi et al. (2021) that indicated an inverse J-shaped curve showing higher number of young trees.

3.3 Community attributes

Both the CF are dominated by *Shorea robusta*, followed by *Schima wallichii* and *Lagerstroemia parviflora* (Table 2 and 3). The importance value index (IVI) of *Shorea robusta* is higher in both forests, with 185.89 and 200.69 in Hazare CF and Janata CF, respectively. *Schima wallichii* also has high IVI in both forests, with 38.94 in Hazare CF and 77.44 in Janata CF. Other species have lower IVI values indicating their lesser ecological significance (Table 2 and 3). The high IVI value of these species may be due to their ability to adapt to the environmental conditions of the forest and to their competitive ability to utilize the available resources (Shameem & Kangroo, 2011). It also suggests that they are preferred by the local communities for their ecological and economic value (Sarkar and Devi 2014; Das et al. 2017).

Table 2: Frequency, relative frequency (F), density (D), relative density (RD), basal area (BA), relative basal area (RBA) and importance value index (IVI) of plant species in Hazare CF.

Species	F %	RF%	D (tree ha ⁻¹)	RD %	BA (m ² ha ⁻¹)	RBA%	IVI
<i>Adina cardifolia</i>	2.86	1.16	0.57	0.53	0.10	0.74	2.44
<i>Buchanania latifolia</i>	14.29	5.81	2.86	2.67	0.20	1.46	9.95
<i>Cassia fistula</i>	8.57	3.49	1.71	1.60	0.09	0.68	5.77
<i>Dillenia pentagyna</i>	2.86	1.16	0.57	0.53	0.07	0.53	2.23
<i>Lagerstroemia parviflora</i>	31.43	12.79	6.29	5.88	0.46	3.46	22.13
<i>Myrsine capitellata</i>	2.86	1.16	0.57	0.53	0.01	0.04	1.74
Other hill species	5.71	2.33	1.14	1.07	0.05	0.39	3.79
<i>Schima wallichii</i>	42.86	17.44	8.57	8.02	1.81	13.47	38.94
<i>Semicarpus anacardum</i>	5.71	2.33	1.14	1.07	0.09	0.64	4.04
<i>Shorea robusta</i>	100.00	40.70	76.00	71.12	9.96	74.07	185.89
<i>Symplocos sps</i>	11.43	4.65	2.29	2.14	0.11	0.80	7.59
<i>Terminalia bellerica</i>	8.57	3.49	1.71	1.60	0.16	1.19	6.28
<i>Terminalia tomentosa</i>	8.57	3.49	3.43	3.21	0.34	2.51	9.21
Total	245.71	100.00	106.86	100.00	13.44	100.00	299.99

Table 3: Frequency, relative frequency (F), density (D), relative density (RD), basal area (BA), relative basal area (RBA) and importance value index (IVI) of plant species in Janata CF.

Species	F %	RF%	D (tree ha ⁻¹)	RD %	BA (m ² ha ⁻¹)	RBA%	IVI
<i>Anogeissus latifolia</i>	7.69	3.85	1.54	2	0.352	3.344	9.19
<i>Lagerstroemia parviflora</i>	15.38	7.69	3.08	4	0.261	2.480	14.17
<i>Schima wallichii</i>	65.38	32.69	19.23	25	2.079	19.748	77.44
<i>Semicarpus anacardum</i>	11.54	5.77	2.31	3	0.078	0.741	9.51
<i>Shorea robusta</i>	100	50.00	59.23	77	7.760	73.691	200.69
Total	200	100.00	85.38	111	10.53	100.00	311.01

3.4 Biodiversity parameters

The Shannon-Wiener diversity index (H') in the studied area was ranging from 0.9 to 1.21 (Table 4), which is similar to range (0.9- 1.0) reported by (Sharma et al, 2018) in Resunga Sacred Grove, Gulmi. However, our result is significantly lower than the range (2.69-4.53) reported by (Sahu et al. 2012), (1.80-1.91) reported by (Shrestha et al. 2020) in sacred groves in Kathmandu Valley, and Ganesh Community Forest and Ramnagar Community Forest in Nepal tropical (Joshi et al. 2019). This could be due to differences in forest type and management practices, as well as variations in the degree of anthropogenic disturbances such as grazing, logging, and fire that can significantly affect the diversity of the forest (Chapagain et al., 2021). The Simpson's index (D) reflected the difference in species richness, with the Janata forest having a value of 0.87 and Hazare having a value of 2.29. The Evenness Index, on the other hand, indicated that the Janata forest had a slightly higher evenness ($e=0.56$) compared to the Hazare forest ($e=0.47$), suggesting a more balanced distribution of species in the Janata forest.

Table 4: Tree Biodiversity parameters in Janata and Hazare Community Forest, Udaypur District, Nepal

Parameters	Janata CF	Hazare CF
Shannon-Wiener Index (H')	0.90	1.21
Simpson's Index (D)	0.87	2.29
Evenness Index (e)	0.56	0.47

3.5 Above Ground Tree Biomass and Carbon stock

Mean AGTB and carbon stock in Janata CF were 98.01 ± 7.90 t ha⁻¹ and 46.06 ± 3.71 t ha⁻¹, respectively. Similarly, the AGTB and carbon stock in Hazare CF were 64.92 ± 9.70 t ha⁻¹ and 30.51 ± 4.56 t ha⁻¹, respectively. As a result, total AGTB and carbon stock were comparatively greater in Janata CF than Hazare CF (Figure 3). This may be due to higher tree density in Janata CF, which could result in a greater AGTB and carbon stock (Joshi et al., 2023). Our Study have reported higher AGTB and carbon stock than Joshi et al. (2023) and lower than Paudel and Bhusal (2016), Karki et al. (2016) and Joshi et al. (2021). This might be due the history of land use, management practices, soil quality, and climate conditions in each area (Rossi et al., 2007).




Figure 3: Total AGTB accumulation and carbon stock in both CFs.

The t-test showed that there was significant difference ($P < 0.05$) in the carbon stock densities of these pools with respective to the ecological regions; the amount of carbon stock density was found to be significantly higher in the Janata CF than in the Hazare CF (Table 5).

The results indicated that *Shorea robusta* (Sal) was the highest contributor to the carbon stock in both forests, with a contribution of 79.51% (36.62 t ha⁻¹) and 77.91% (23.77 t ha⁻¹) in Janata and Hazare CF, respectively. This could be because *Shorea robusta* are dominant species in both forests, and their abundance could contribute to high-

er carbon stock. The other tree species also contributed a significantly higher percentage of carbon stock in both CFs (Table 6). This finding is consistent with previous studies, which showed that Sal is the highest contributor to carbon stock in the Terai region of Nepal including the FRA report (2014), Neupane and Sharma (2014) and Joshi et al., (2021).

3.6 Correlation Analysis

The forest biometric variables are statistically significant at the 0.05 level (2 tailed). The correlation analysis

Table 5: Aboveground Tree carbon stock densities in the study areas

Carbon Pool	df	Janata CF	Hazare CF	Mean	p-value	remarks
Aboveground Tree Carbon stock (t ha ⁻¹)	40.81	46.06	30.51	38.29	0.0015	*

Remarks: * = significant at $P < 0.05$

Table 6: The percentage share of species contribution on carbon stock of Janata and Hazare CFs, Udaypur District, Nepal

Janata CF			Hazare CF		
Species	C ± SE (t ha ⁻¹)	C (%)	Species	C ± SE (t ha ⁻¹)	C (%)
<i>Anogeissus latifolia</i>	1.08	2.34	<i>Adina cardifolia</i>	0.17	0.56
<i>Lagerstroemia parviflora</i>	1.13	2.45	<i>Buchanania latifolia</i>	0.32	1.04
<i>Schima wallichii</i>	7.08	15.36	<i>Cassia fistula</i>	0.13	0.41
<i>Semicarpus anacardum</i>	0.16	0.35	<i>Dillenia pentagyna</i>	0.12	0.41
<i>Shorea robusta</i>	36.62	79.51	<i>Lagerstroemia parviflora</i>	0.99	3.24
			<i>Myrsine capitellata</i>	0.01	0.02
			<i>Other hill species</i>	0.08	0.27
			<i>Schima wallichii</i>	3.58	11.72
			<i>Semicarpus anacardum</i>	0.13	0.43
			<i>Shorea robusta</i>	23.77	77.91
			<i>Symplocos sps</i>	0.15	0.50
			<i>Terminalia bellerica</i>	0.24	0.80
			<i>Terminalia tomentosa</i>	0.83	2.71
Total	46.06 ± 3.71	100.00	Total	30.51 ± 4.56	100.00

Table 7: Correlation analysis between the carbon stock (t ha⁻¹), biomass (t ha⁻¹), tree density (trees ha⁻¹), tree DBH (cm), tree height (m), basal area (m² ha⁻¹) and seedling density (seedlings ha⁻¹) of Janata CF.

Variables	Carbon	Biomass	Density	DBH	Height	Basal Area	Seedling Density
Carbon	1	1*	0.66*	0.44*	0.3	0.97*	0.12
Biomass	1*	1	0.66*	0.44*	0.3	0.97*	0.12
Density	0.66*	0.66*	1	-0.34	-0.25	0.76*	-0.06
DBH	0.44*	0.44*	-0.34	1	0.60*	0.34	0.2
Height	0.3	0.3	-0.25	0.60*	1	0.11	0.145
Basal Area	0.97*	0.97*	0.76*	0.34	0.11	1	0.06
Seedling Density	0.12	0.12	-0.06	0.2	0.15	0.06	1

No. of plots. N=26, *Represents the significant difference, p<0.05

Table 8: Correlation analysis between the carbon stock (t ha⁻¹), biomass (t ha⁻¹), tree density (trees ha⁻¹), tree DBH (cm), tree height (m), basal area (m² ha⁻¹) and seedling density (seedlings ha⁻¹) of Hazare CF.

Variables	Carbon	Biomass	Density	DBH	Height	Basal Area	Seedling Density
Carbon	1	1*	0.41*	0.63*	0.48*	0.98*	0.08
Biomass	1*	1	0.41*	0.63*	0.48*	0.98*	0.08
Density	0.41*	0.41	1	-0.31	-0.29	0.48*	0.18
DBH	0.63*	0.63*	-0.31	1	0.74*	0.63*	-0.05
Height	0.48*	0.48*	-0.29	0.74*	1	0.40*	-0.17
Basal Area	0.98*	0.98*	0.48*	0.63*	0.40*	1	0.08
Seedling Density	0.08	0.08	0.18	-0.05	-0.17	0.08	1

No. of plots. N=35, *Represents the significant difference, p<0.05

(Table 7 and 8) showed that the forest C stock in both forest was positively correlated with tree DBH, height, basal area (BA) and biomass. A similar finding was also observed by Thapa-Magar and Shrestha (2015), Shaheen et al. (2016) and Sharma et al. (2020). Thus, tree diameter, height, BA and biomass are the determinant variable for forest C. The negative correlation of stand density with tree DBH and tree height in both CF is in accordance with Nizami et al. (2009), Nizami (2012), Amir et al. (2018) and Sharma et al. (2020). Natural thinning process due to resource competition resulting in lower stand density but allowing the remaining trees to grow bigger and selective removal of big trees by humans for timber extraction or other purposes could be the possible reasons for such negative correlation (Nizami 2012). Similarly, the positive correlation between DBH and height observed in this study is consistent with previous research by Nizami et al. (2009) and Sharma et al. (2020). This could be attributed to the similar environmental conditions, which influenced their growth rate in a similar manner (Ekoungoulou et al., 2018). The negative correlation between stand density and seedling density in Janata CF could be due to competition among trees for resources such as light, nutrients, and water, leading to reduced availability for seedlings to establish and grow (Jackson, 1994). However, stand density and seedling density were positively correlated in Hazare

CF which could be due to the presence of suitable environmental conditions such as light availability, nutrient availability, and soil moisture, as well as management practices such as selective removal of big trees and thinning operations which may have promoted both the growth and survival of both established trees and seedlings.

4. CONCLUSION

In this comparative study, we explored the richness of community-managed forests in terms of regeneration status, vegetation diversity, stand structure, and carbon stock across two ecological regions. Our findings indicate that Janata CF and Hazare CF exhibit significant differences in their regeneration, DBH distribution, species diversity, and carbon stock. Janata CF has higher regeneration compared to Hazare CF, which may be attributed to its better forest management practices. The distribution of DBH in Janata CF shows a bell-shaped curve, whereas Hazare CF exhibits an interrupted curve, which may indicate differences in the growth patterns of the dominant species. Both forests are dominated by *Shorea robusta*, which suggests the importance of this species for the overall forest structure and function. However, Hazare CF shows higher species diversity indices, indicating a more diverse forest ecosystem compared to Janata CF. Interestingly, despite lower species diversity, Janata CF has higher carbon stock, which may

be attributed to the higher density of *Shorea robusta* in this forest. *Shorea robusta* was found to be the highest contributor to the carbon stock in both forests, highlighting the significance of this species in the carbon sequestration potential of the forests. The difference in ecological factors such as temperature, rainfall, soil nutrient availability, and management practices could be attributed to the observed differences in the studied parameters between the two ecological regions. This study emphasizes the importance of community-managed forests in maintaining biodiversity and ecosystem services and underscores the need for tailored forest management practices based on the specific ecological and social contexts of the forests. It is also recommended to conduct similar studies in other community-managed forests across different ecological regions to better understand the patterns of regeneration, vegetation diversity, stand structure, and carbon stock. Future studies focusing on investigating the underlying factors driving the observed patterns and relationships, such as the effects of disturbance, climate, and management practices on forest structure and function are also needed.

Author's contribution statement

S. Ayer: Conceptualization (lead); data curation (lead); formal analysis (lead); writing original draft (lead); writing—review and editing (lead). **A. Prabhakar, K. Adhikari and S. Poudel:** Formal analysis (equal); writing original draft (equal); writing—review and editing (equal). **S. Basnet, M. Timilsina, M. Giri, P.T. Magar, S. Lamichane, A. Singh:** Writing—review and editing (equal).

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

Data will be made available on request.

Acknowledgments

We are indebted to Divisional Forest Office (DFO), Udaypur (Triveni) for providing the platform to conduct this research. Also, we are most sincerely grateful to Assistant Professor Mr. Jeetendra Gautam of the Faculty of Forestry, Hetauda, Agriculture and Forestry University for his assistance in analysis.

REFERENCES

- Acharya, K.P., Khadka, S., Lekhak, H.D., Chaudhary, R.P. & Vetaas, O.R. (2007). Species composition and regeneration of coniferous forest in Manang. In: R.P. Chaudhary, T.H. Aase, O.R. Vetaas and B.P. Subedi (eds.) Local Effects of Global Changes in the Himalayas: Manang, Nepal (pp. 131–138). Tribhuvan: Tribhuvan University and Bergen: University of Bergen.
- Amir, M., Liu, X., Ahmad, A., Saeed, S., Mannan, A., & Muneer, M. A. (2018). Patterns of biomass and carbon allocation across chronosequence of chir pine (*Pinus roxburghii*) forest in Pakistan: inventory-based estimate. *Advances in Meteorology*, 2018.
- Anup, K. C. (2017, March). Community forestry management and its role in biodiversity conservation in Nepal. In *Global Exposition of Wildlife Management* (Vol. 51). InTech. <https://doi.org/10.5772/65926>
- Aryal, B., Regmi, S., & Timilsina, S. (2021). Regeneration status and species diversity of major tree species under scientific forest management in Kapilbastu district, Nepal. *Banko Janakari*, 31(2), 26-39. <https://doi.org/10.3126/banko.v31i2.41898>
- Aryal, S., Shrestha, S., Maraseni, T., Wagle, P. C., & Gaire, N. P. (2018). Carbon stock and its relationships with tree diversity and density in community forests in Nepal. *International Forestry Review*, 20(3), 263-273. <https://doi.org/10.1505/146554818824063069>
- Awasthi, N., Bhandari, S. K., & Khanal, Y. (2015). Does scientific forest management promote plant species diversity and regeneration in Sal (*Shorea robusta*) forest? A case study from Lumbini collaborative forest, Rupandehi, Nepal. *Banko Janakari*, 25(1), 20-29. Does scientific forest management promote plant species diversity and regeneration in Sal (*Shorea robusta*) forest? <https://doi.org/10.3126/banko.v25i1.13468>
- Ayer, K., Kandel, P., Gautam, D., Khadka, P., & Miya, M. S. (2022). Comparative Study of Carbon Stock and Tree Diversity between Scientifically and Conventionally Managed Community Forests of Kanchanpur District, Nepal: 10.32526/ennrj/20/202200010. *Environment and Natural Resources Journal*, 20(5), 494-504. <https://doi.org/10.32526/ennrj/20/202200010>
- Basyal, S., Lekhak, H.D. & Devkota, A. (2011) Regeneration of *Shorea robusta* Gaertn in tropical forest of Palpa district, central Nepal. *Scientific World*, 9(9), 53–56. <https://doi.org/10.3126/sw.v9i9.5519>
- Bauhus, J., Pokorny, B., van der Meer, P. J., Kanowski, J., & Kanninen, M. (2010). Ecosystem goods and services - the key for sustainable plantations. In J. Bauhus, P. J. van der Meer, & M. Kanninen (Eds.), *Ecosystem Goods and Services from Plantation Forests* (pp. 205-227). Earthscan.
- Baul, T. K., Chakraborty, A., Nandi, R., Mohiuddin, M., Kilpeläinen, A., & Sultana, T. (2021). Effects of tree species diversity and stand structure on carbon stocks of homestead forests in Maheshkhali Island, Southern Bangladesh. *Carbon Balance and Management*, 16(1), 1-15. <https://doi.org/10.1186/s13021-021-00175-6>
- Bhandari, S. K., Maraseni, T., Timilsina, Y. P., & Parajuli, R. (2021). Species composition, diversity, and carbon stock in trees outside forests in middle hills of Nepal. *Forest Policy and Economics*, 125, 102402.

- <https://doi.org/10.1016/j.forpol.2021.102402>
- Bhatta, S. P., & Devkota, A. (2020). Community structure and regeneration status of Sal (*Shorea robusta* Gaertn.) forests of Dadeldhura district, Western Nepal. *Community Ecology*, 21(2), 191-201. <https://doi.org/10.1007/s42974-020-00021-8>
- Bose, A. K., Weiskittel, A., Wagner, R. G., & Kuehne, C. (2016). Assessing the factors influencing natural regeneration patterns in the diverse, multi-cohort, and managed forests of Maine, USA. *Journal of Vegetation Science*, 27, 1140–1150. <https://doi.org/10.1111/jvs.12433>
- Brockerhoff, E. G., Barbaro, L., Castagnyrol, B., Forrester, D. I., Gardiner, B., González-Olabarria, J. R., ... & Jactel, H. (2017). Forest biodiversity, ecosystem functioning and the provision of ecosystem services. *Biodiversity and Conservation*, 26, 3005-3035 <https://doi.org/10.1007/s10531-017-1453-2>
- Brockerhoff, E. G., Jactel, H., Parrotta, J. A., & Ferraz, S. F. (2013). Role of eucalypt and other planted forests in biodiversity conservation and the provision of biodiversity-related ecosystem services. *Forest Ecology and Management*, 301, 43-50. <https://doi.org/10.1016/j.foreco.2012.09.018>
- Cannell, M. G. (1996). Forests as carbon sinks mitigating the greenhouse effect. *The Commonwealth Forestry Review*, 92-99. <https://doi.org/10.1007/s40725-016-0031-2>
- Chapagain, U., Chapagain, B. P., Nepal, S., & Manthey, M. (2021). Impact of disturbances on species diversity and regeneration of Nepalese Sal (*Shorea robusta*) forests managed under different management regimes. *Earth*, 2(4), 826-844. <https://doi.org/10.3390/earth2040049>
- Chauhan, P.S., Negi, J.D.S., Singh, L., & Manhas, R.K. (2008). Regeneration status of Sal forests of Doon Valley. *Annals of Forestry*, 16(2), 178–182.
- Chave, J., Andalo, C., Brown, S., Cairns, M.A., Chambers, J.Q., Eamus, D. et al. (2005). Tree allometry and improved estimation of carbon stocks and balance in tropical forests. *Oecologia*, 145: 87–99. <https://doi.org/10.1007/s00442-005-0100-x>
- Chazdon, R. L., & Guariguata, M. R. (2016). Natural regeneration as a tool for large-scale forest restoration in the tropics: prospects and challenges. *Biotropica*, 48(6), 716-730.
- Clatterbuck, W. K., Stringer, J. W., & Tankersley, L. (2011). PB1798 Uneven-age Management in Mixed Species, Southern Hardwoods: Is It Feasible and Sustainable?.
- Daba, D. E., & Soromessa, T. (2019). The accuracy of species-specific allometric equations for estimating aboveground biomass in tropical moist montane forests: case study of *Albizia grandibracteata* and *Trichilia dregeana*. *Carbon balance and management*, 14, 1-13. <https://doi.org/10.1186/s13021-019-0134-8>
- Dar JA, Rather MY, Subashree K, Sundarapandian S, Khan ML.(2017). Distribution patterns of tree, understorey, and detritus biomass in coniferous and broad-leaved forests of Western Himalaya, India. *J Sustain Forest* 36: 787-805. <https://doi.org/10.1080/10549811.2017.1363055>
- Decocq, G., Andrieu, E., Brunet, J., Chabrierie, O., De Frenne, P., De Smedt, P., ... & Wulf, M. (2016). Ecosystem services from small forest patches in agricultural landscapes. *Current Forestry Reports*, 2, 30-44. <https://doi.org/10.1007/s40725-016-0028-x>
- DoF. 2004. Community Forestry Resource Inventory Guideline. Department of Forest (DoF), Ministry of Forest and Soil Conservation, Kathmandu, Nepal.
- Ekoungoulou, R., Nzala, D., Liu, X., & Niu, S. (2018). Tree biomass estimation in central African forests using allometric models. *Open Journal of Ecology*, 8(3), 209-237.
- Forrester DI, Bausch J.(2016). A review of processes behind diversity—productivity relationships in forests. *Curr For Rep* 2:45–61 <https://doi.org/10.1007/s40725-016-0031-2>
- Gamfeldt L, Snäll T, Bagchi R, Bengtsson J. et al. (2013). Higher levels of multiple ecosystem services are found in forests with more tree species. *Nat Commun* 4:1340 <https://doi.org/10.1038/ncomms2328>
- Ghimire, P. and Lamichhane, U. (2020). Community Based Forest Management in Nepal: Current Status, Successes and Challenges. *Grassroots Journal of Natural Resources*, 3(2): 16-29. Doi: <https://doi.org/10.33002/nr2581.6853.03022>
- Ghimire, P., & Lamichhane, U. (2021). Plant species diversity and crown cover response to vegetation regeneration in community-managed forest in Makawanpur District, Nepal. *Asian Journal of Forestry*, 5(1). <https://doi.org/10.13057/asianjfor/r050105>
- Hooper DU, Chapin FS, Ewel JJ, Hector A, Inchausti P, Lavorel S, Lawton JH, Lodge DM, Loreau M, Naeem S, Schmid B. (2005). Effects of biodiversity on ecosystem functioning: a consensus of current knowledge. *Ecol Monogr* 75:3–35 <https://doi.org/10.1890/04-0922>
- Isbell F, Calcagno V, Hector A et al. (2011). High plant diversity is needed to maintain ecosystem services. *Nature* 477:199–202. <https://doi.org/10.1038/nature10282>
- Jackson, J. K. (1994). Manual of afforestation in Nepal, vol. 1. *Forest Research and Survey Center, Ministry of Forests and Soil Conservation, Kathmandu, Nepal.*
- JOSHI, P., JOSHI, R., MAHARJAN, A., PANTA, M., & CHAND, P. (2023). Vegetation diversity, structure, composition and carbon stock of community managed forests of Mid-hills, Nepal. *Asian Journal of Forest-*

- ry, 7(1). <https://doi.org/10.13057/asianjfor/r070104>
- JOSHI, P., JOSHI, R., MAHARJAN, A., PANTA, M., & CHAND, P. (2023). Vegetation diversity, structure, composition and carbon stock of community managed forests of Mid-hills, Nepal. *Asian Journal of Forestry*, 7(1). <https://doi.org/10.13057/asianjfor/r070104>
- Joshi, R., Pangen, M., Neupane, S. S., & Yadav, N. P. (2021). Regeneration status and carbon accumulation potential in community managed sal (*Shorea robusta*) forests of Far-Western Terai Region, Nepal. *European Journal of Ecology*, 7(1). <https://doi.org/10.17161/eurojecol.v7i1.15005>
- Karki S, Joshi NR, Udas E, Adhikari MD, Sherpa S, Kotru R, Karky BS, Chettri N, Ning W. (2016). Assessment of Forest Carbon Stock and Carbon Sequestration Rates at the ICIMOD Knowledge Park in Godavari. ICIMOD Working Paper, Kathmandu, Nepal.
- Karki, S., Joshi, N. R., Udas, E., Adhikari, M. D., Sherpa, S., Kotru, R., ... & Ning, W. (2016). Assessment of forest carbon stock and carbon sequestration rates at the ICIMOD Knowledge Park at Godavari. *ICIMOD Working Paper*, (2016/6). <https://doi.org/10.53055/ICIMOD.622>
- Khaine, I., Woo, S. Y., Kwak, M., Lee, S. H., Je, S. M., You, H., ... & Kim, J. (2018). Factors affecting natural regeneration of tropical forests across a precipitation gradient in Myanmar. *Forests*, 9(3), 143. <https://doi.org/10.3390/f9030143>
- KHATRI, B., GAUTAM, D., GAIRE, D., KHANAL, Y., BHATTARAI, S., & KHATRI, S. K. (2021). Species Diversity and Regeneration under the Scientific Forest Management Practice in Tropical Region of Nepal. *Agriculture and Forestry Journal*, 5(2), 89-95. <https://doi.org/10.5281/zenodo.5796310>
- Liaison, L. (2013). *Conservation on Biological Diversity. UNCCD, 3. Climate change, Biodiversity and Land degradation*
- Liang, J., Crowther, T. W., Picard, N., Wiser, S., Zhou, M., Alberti, G., ... & Reich, P. B. (2016). Positive biodiversity-productivity relationship predominant in global forests. *Science*, 354(6309), aaf8957. <https://doi.org/10.1126/science.aaf8957>
- Magurran, A. E. (2005). Biological diversity. *Current Biology*, 15(4), R116-R118.
- Mandal, R. A., Dutta, I. C., Jha, P. K., & Karmacharya, S. (2013). Relationship between carbon stock and plant biodiversity in collaborative forests in Terai, Nepal. *International Scholarly Research Notices*, 2013. <https://doi.org/10.1155/2013/625767>
- Manna, S. S., & Mishra, S. P. (2017). Diversity, population structure and regeneration of tree species in Lalgah forest range of West Bengal, India. *International Journal of Botany Studies*, 2, 191–195.
- Marais, Z. E., Baker, T. P., Hunt, M. A., & Mendham, D. (2022). Shelterbelt species composition and age determine structure: Consequences for Ecosystem Services. *Agriculture, Ecosystems & Environment*, 329, 107884. <https://doi.org/10.1016/j.agee.2022.107884>.
- Mori, A. S., Lertzman, K. P., & Gustafsson, L. (2017). Biodiversity and ecosystem services in forest ecosystems: a research agenda for applied forest ecology. *Journal of Applied Ecology*, 54(1), 12-27. <https://doi.org/10.1111/1365-2664.12669>
- Mugasha, W. A., Mauya, E. W., Njana, A. M., Karlsson, K., Malimbwi, R. E., & Ernest, S. (2019). Height-diameter allometry for tree species in tanzania mainland. *International Journal of Forestry Research*, 2019. <https://doi.org/10.1155/2019/4832849>
- Mukul, S. A., Halim, M. A., Herbohn, J. (2021). Forest carbon stock and fluxes: Distribution, biogeochemical cycles, and measurement techniques. *Life on land, encyclopedia of the UN sustainable development goals*, 365-380. https://doi.org/10.1007/978-3-319-71065-5_23-1
- Napit, R., & Paudel, P. K. (2015). Response of natural forest regeneration to human disturbance in Banke national park. *Banko Janakari*, 25, 39–49. DOI: <https://doi.org/10.3126/banko.v25i1.13470>.
- Nizami SM, Mirza SN, Livesley S, Arndt S, Fox JC, Khan IA, Mahmood T. (2009). Estimating carbon stocks in sub-tropical pine (*Pinus roxburghii*) forests of Pakistan. *Pak J Agr Sci* 46: 266-270
- Nizami, S. M. (2012). The inventory of the carbon stocks in sub tropical forests of Pakistan for reporting under Kyoto Protocol. *Journal of Forestry Research*, 23(3), 377-384.
- Nizami, S. M., Mirza, S. N., Livesley, S., Arndt, S., Fox, J. C., Khan, I. A., & Mahmood, T. (2009). Estimating carbon stocks in sub-tropical pine (*Pinus roxburghii*) forests of Pakistan. *Pakistan Journal of Agricultural Sciences*, 46(4), 266-270.
- Pandey H.P. & Bhusal M.(2016). A comparative study on carbon stock in Sal. *Banko Janakari* 26 (1): 24-31. <https://doi.org/10.3126/banko.v26i1.15498>
- Pandey, H. P., & Bhusal, M. (2016). A comparative study on carbon stock in Sal (*Shorea robusta*) forest in two different ecological regions of Nepal. *Banko Janakari*, 26(1), 24-31. <https://doi.org/10.3126/banko.v26i1.15498>
- Pandey, H. P., & Pokhrel, N. P. (2021). Formation trend analysis and gender inclusion in community forests of Nepal. *Trees, Forests and People*, 5, 100106. <https://doi.org/10.1016/j.tfp.2021.100106>
- Pandey, H.P., Maren, I.E. & Dutta, I.C. (2012). REDD+ in community forests, western Nepal: A case from Gorkha district, central Himalaya. Chisinau: LAMBERT Academic Publishing
- Paudyal, K., Baral, H., Lowell, K., & Keenan, R. J. (2017). Ecosystem services from community-based forestry

- in Nepal: Realising local and global benefits. *Land use policy*, 63, 342-355. <https://doi.org/10.1016/j.landusepol.2017.01.046>
- Poudyal, B. H., Maraseni, T., & Cockfield, G. (2019). Impacts of forest management on tree species richness and composition: Assessment of forest management regimes in Tarai landscape Nepal. *Applied Geography*, 111, 102078. <https://doi.org/10.1016/j.apgeog.2019.102078>
- Rahman, M. R., Rahman, M. M., & Chowdhury, M. A. (2020). Assessment of natural regeneration status: the case of Durgapur hill forest, Netrokona, Bangladesh. *Geology, Ecology, and Landscapes*, 4(2), 121-130. <https://doi.org/10.1080/24749508.2019.1600911>
- Rossi S, Deslauriers A, Anfodillo & Carraro V.(2007). Evidence of threshold temperatures for xylogenesis in conifers at high altitudes. *Oecologia* 152: 1-12. <https://doi.org/10.1007/s00442-006-0625-7>
- Sahu SC, Dhal NK, Lal B, Mohanty RC. (2012). Differences in tree species diversity and soil nutrient status in a tropical sacred forest ecosystem on Niyamgiri hill range, Eastern Ghats, India. *J Mount Sci* 9 (4): 492-500. <https://doi.org/10.1007/s11629-012-2302-0>
- Sapkota, I.P., Tigabu, M. & Odén, P.C. (2009). Spatial distribution, advanced regeneration and stand structure of Nepalese Sal (*Shorea robusta*) forests subject to disturbances of different intensities. *Forest Ecology and Management*, 257(9), 1966–1975. <https://doi.org/10.1016/j.foreco.2009.02.008>
- Sarkar, M., & Devi, A. (2014). Assessment of diversity, population structure and regeneration status of tree species in Hollongapar Gibbon wildlife sanctuary, Assam, Northeast India. *Tropical Plant Research*, 1, 26–36.
- Shaheen, H., Khan, R. W. A., Hussain, K., Ullah, T. S., Nasir, M., & Mehmood, A. (2016). Carbon stocks assessment in subtropical forest types of Kashmir Himalayas. *Pak. J. Bot*, 48(6), 2351-2357.
- Shameem, S.A. & Kangroo, I.N. (2011). Comparative assessment of edaphic features and phytodiversity in lower Dachigam National Park, Kashmir Himalaya, India. *African Journal of Environmental Science and Technology*, 5(11): 972–984. <https://doi.org/10.5897/AJEST11.099>
- Shannon, C. E., & Weaver, W. (1949). *The Mathematical Theory of Communication*. Urbana, IL: The University of Illinois Press, 1-117.
- Sharma BK, Pokharel CP, Shrestha L. (2018). Forest diversity and carbon sequestration in Resunga Sacred Grove, Gulmi, Nepal. *J Nat His Mus* 29: 60-69. <https://doi.org/10.3126/jnhm.v29i0.19038>.
- Sharma, K. P., Bhatta, S. P., & Lamsal, S. K. (2020). Species diversity and regeneration status of community-managed hill sal (*Shorea robusta*) forest in Central Nepal. *Current Science*, 119(1), 83-92. <https://doi.org/10.18520/cs/v119/i1/83-92>
- Sharma, K. P., Bhatta, S. P., Khatri, G. B., Pajiyar, A., & Joshi, D. K. (2020). Estimation of carbon stock in the chir pine (*Pinus roxburghii* Sarg.) plantation forest of Kathmandu valley, Central Nepal. *Journal of Forest and Environmental science*, 36(1), 37-46. <https://doi.org/10.7747/JFES.2020.36.1.37>
- Shi, L., & Liu, S. (2017). Methods of Estimating Forest Biomass: A Review. *Biomass Volume Estimation and Valorization for Energy*. <https://doi.org/10.5772/65733>
- Shrestha, LJ, Devkota M, Sharma BK. (2020). Tree diversity conservation initiatives in sacred groves of Kathmandu Valley, Nepal. *Nepal J Sci Technol* 19 (1): 60-68. <https://doi.org/10.3126/njst.v19i1.29768>
- Simpson EH. (1949). Measurement of diversity. *Nature* 163 (4148): 688-688. <https://doi.org/10.1038/163688a0>
- Smith, C. T. (2016). *Sustainable Forestry: Ecology and Silviculture for Resilient Forests*.
- Temgoua, L. F., Momo Solefack, M. C., Nyong, P. A., & Tadjou, P. (2020). Floristic diversity and exploitable potential of commercial timber species in the Cobaba community forest in Eastern Cameroon: implications for forest management. *Forest Science and Technology*, 16(2), 56-67. <https://doi.org/10.1080/21580103.2020.1750493>
- Thapa, P. (2021). The relationship between land use and climate change: a case study of Nepal. *The Nature, Causes, Effects and Mitigation of Climate Change on the Environment*. <https://doi.org/10.5772/intechopen.98282>
- Thapa-Magar, K. B., & Shrestha, B. B. (2015). Carbon stock in community managed hill sal (*Shorea robusta*) forests of central Nepal. *Journal of Sustainable forestry*, 34(5), 483-501. <https://doi.org/10.2989/20702620.2019.1686690>
- Thompson, I. D., Okabe, K., Tylianakis, J. M., Kumar, P., Brockerhoff, E. G., Schellhorn, N. A., ... & Nasi, R. (2011). Forest biodiversity and the delivery of ecosystem goods and services: translating science into policy. *BioScience*, 61(12), 972-981. <https://doi.org/10.1525/bio.2011.61.12.7>
- Whittaker, R. H. (1972). Evolution and measurement of species diversity. *Taxon*, 21(2-3), 213-251. <https://doi.org/10.2307/1218190>
- Wilsey, B. J., Chalcraft, D. R., Bowles, C. M., & Willig, M. R. (2005). Relationships among indices suggest that richness is an incomplete surrogate for grassland biodiversity. *Ecology*, 86(5), 1178-1184. <https://doi.org/10.1890/04-0394>
- Zobel, D. B., Yadav, U. K. R., Jha, P. K., & Behan, M. J. (1987). *A practical manual for ecology*. Kathmandu: Ratna Pustak Bhandar.