# Treatise on Invertebrate Paleontology

## Part L MOLLUSCA 4

## Revised

## Volume 2: Carboniferous and Permian Ammonoidea (Goniatitida and Prolecanitida)

WILLIAM M. FURNISH, BRIAN F. GLENISTER, JÜRGEN KULLMANN, and ZHOU ZUREN

Prepared under Sponsorship of

The Paleontological Society The Palaeontographical Society SEPM (Society for Sedimentary Geology) The Palaeontological Association

RAYMOND C. MOORE Founder PAUL A. SELDEN Editor

JANE KERNS, JILL HARDESTY, MICHAEL CORMACK, DENISE MAYSE Assistant Editors and Editorial Staff

> The University of Kansas Paleontological Institute Lawrence, Kansas 2009

> > © 2009 University of Kansas Paleontological Institute

© 2009 by

The University of Kansas Paleontological Institute

All Rights Reserved

Library of Congress Catalogue Card Number 53-12913 ISBN 9781-891276-61-1

## This volume is dedicated to the memory of WILLIAM MADISON FURNISH 1912–2007

Distributed by the Paleontological Institute, The University of Kansas, 1475 Jayhawk Blvd., Room 119, Lawrence, Kansas 66045-7613, USA, www.paleo.ku.edu, from which current price lists of parts in print may be obtained and to which all orders and related correspondence should be directed. Editorial office of the *Treatise*: Paleontological Institute, The University of Kansas, 1475 Jayhawk Blvd., Room 119, Lawrence, Kansas 66045-7613, USA, www.paleo.ku.edu.

Citation information: Selden, P. A., ed. 2009. Treatise on Invertebrate Paleontology. Part L, Mollusca 4, Revised, vol. 2. The University of Kansas Paleontological Institute. Lawrence, Kansas. xxix + 258 p., 139 fig., 1 table. The *Treatise on Invertebrate Paleontology* has been made possible by (1) funding principally from the National Science Foundation of the United States in its early stages, from The Geological Society of America through the bequest of Richard Alexander Fullerton Penrose, Jr., and from The Kansas University Endowment Association through the bequest of Raymond C. and Lillian B. Moore; (2) contribution of the knowledge and labor of specialists throughout the world, working in cooperation under sponsorship of the Paleontological Society, the SEPM (Society for Sedimentary Geology), the Palaeontographical Society, and the Palaeontological Association; (3) acceptance by The University of Kansas of publication without any financial gain to the University; and (4) generous contributions by our individual and corporate sponsors.

## PART L, Revised MOLLUSCA 4

## Volume 2: Carboniferous and Permian Ammonoidea (Goniatitida and Prolecanitida)

WILLIAM M. FURNISH, BRIAN F. GLENISTER, JÜRGEN KULLMANN, and ZHOU ZUREN

#### CONTENTS

INFORMATION ON <i>TREATISE</i> VOLUMES
Editorial Preface (Paul A. Selden)x
STRATIGRAPHIC DIVISIONS
COORDINATING AUTHORS' PREFACE (Jürgen Kullman & Zhou Zuren) xxiv
Repositories and Their Abbreviations
CARBONIFEROUS AND PERMIAN AMMONOIDEA (Goniatitida and Prolecanitida)
(William M. Furnish, Brian F. Glenister, Jürgen Kullmann, & Zhou Zuren)1
Order GONIATITIDA1
Suborder TORNOCERATINA1
Superfamily TORNOCERATOIDEA1
Superfamily DIMEROCERATOIDEA2
Superfamily PRAEGLYPHIOCERATOIDEA
PRIONOCERATOIDEA (Jürgen Kullmann)2
Superfamily PRIONOCERATOIDEA
Family PRIONOCERATIDAE
Family GATTENDORFIIDAE
Family PSEUDARIETITIDAE8
Family VOEHRINGERITIDAE9
KARAGANDOCERATOIDEA (Jürgen Kullmann)11
Superfamily KARAGANDOCERATOIDEA11
Family KARAGANDOCERATIDAE11
Family PRODROMITIDAE11
PSEUDOHALORITOIDEA (Brian F. Glenister, William M. Furnish, & Zhou Zuren)16
Superfamily PSEUDOHALORITOIDEA16
Family MAXIMITIDAE16
Family PSEUDOHALORITIDAE16
Suborder GONIATITINA
PERICYCLOIDEA (Jürgen Kullmann)
Superfamily PERICYCLOIDEA
Family PERICYCLIDAE

Eamily INTOCED ATIDAE	20
Family INTOCERATIDAE Family MUENSTEROCERATIDAE	
Family MOENSTEROCERATIDAE	
NOMISMOCERATOIDEA (Jürgen Kullmann)	
Superfamily NOMISMOCERATOIDEA	
Family NOMISMOCERATIDAE	
Family ENTOGONITIDAE	
DIMORPHOCERATOIDEA (Jürgen Kullmann)	
Superfamily DIMORPHOCERATOIDEA	
Family DIMORPHOCERATIDAE	
Family GIRTYOCERATIDAE	
Family BERKHOCERATIDAE	
Family EOGONIOLOBOCERATIDAE	
Family ANTHRACOCERATIDAE	50
GONIATITOIDEA (Jürgen Kullmann)	
Superfamily GONIATITOIDEA	
Family GONIATITIDAE	
Family DELEPINOCERATIDAE	
AGATHICERATOIDEA (Jürgen Kullmann)	
Superfamily AGATHICERATOIDEA	
Family AGATHICERATIDAE	
NEOGLYPHIOCERATOIDEA (Jürgen Kullmann)	
Superfamily NEOGLYPHIOCERATOIDEA	
Family NEOGLYPHIOCERATIDAE	63
Family CRAVENOCERATIDAE	
Family FERGANOCERATIDAE	71
Family RHYMMOCERATIDAE	71
NEODIMORPHOCERATOIDEA (Jürgen Kullmann)	73
Superfamily NEODIMORPHOCERATOIDEA	73
Family RAMOSITIDAE	73
Family NEODIMORPHOCERATIDAE	
GASTRIOCERATOIDEA (Jürgen Kullmann)	
Superfamily GASTRIOCERATOIDEA	
Family GLAPHYRITIDAE	
Family HOMOCERATIDAE	
Family RETICULOCERATIDAE	
Family GASTRIOCERATIDAE	
THALASSOCERATOIDEA (Jürgen Kullmann)	
Superfamily THALASSOCERATOIDEA	102
Family BISATOCERATIDAE	
Family THALASSOCERATIDAE	
SCHISTOCERATOIDEA (Jürgen Kullmann)	
Superfamily SCHISTOCERATOIDEA	
Family SCHISTOCERATIDAE	
Family PSEUDOPARALEGOCERATIDAE	
Family ORULGANITIDAE	
Family WELLERITIDAE	116

Family CHRISTIOCERATIDAE	118
GONIOLOBOCERATOIDEA (Jürgen Kullmann)	120
Superfamily GONIOLOBOCERATOIDEA	120
Family WIEDEYOCERATIDAE	120
Family GONIOLOBOCERATIDAE	
ADRIANITOIDEA (Brian F. Glenister, William M. Furnish, & Zhou Zuren)	125
Superfamily ADRIANITOIDEA	125
Family ADRIANITIDAE	
SHUMARDITOIDEA (Brian F. Glenister, William M. Furnish, & Zhou Zuren)	136
Superfamily SHUMARDITOIDEA	
Family SÓMOHOLITIDAE	
Family PARASHUMARDITIDAE	136
Family SHUMARDITIDAE	139
Family PERRINITIDAE	
CYCLOLOBOIDEA (Brian F. Glenister, William M. Furnish, & Zhou Zuren)	145
Superfamily CYCLOLOBOIDEA	145
Family VIDRIOCERATIDAE	145
Family CYCLOLOBIDAE	
MARATHONITOIDEA (Brian F. Glenister, William M. Furnish, & Zhou Zuren)	158
Superfamily MARATHONITOIDEA	158
Family MARATHONITIDAE	
Family HYATTOCERATIDAE	163
NEOICOCERATOIDEA	
(William M. Furnish, Brian F. Glenister, Jürgen Kullmann, & Zhou Zuren)	
Superfamily NEOICOCERATOIDEA	166
Family NEOICOCERATIDAE	
Family EUPLEUROCERATIDAE	
Family PARAGASTRIOCERATIDAE	
Family METALEGOCERATIDAE	
POPANOCERATOIDEA (Brian F. Glenister, William M. Furnish, & Zhou Zuren)	
Superfamily POPANOCERATOIDEA	180
Family POPANOCERATIDAE	
Family MONGOLOCERATIDAE	181
PROLECANITIDA	
(William M. Furnish, Brian F. Glenister, Jürgen Kullmann, & Zhou Zuren)	
Order PROLECANITIDA	185
Superfamily PROLECANITOIDEA	
Family PROLECANITIDAE	
Family DARAELITIDAE	
MEDLICOTTIOIDEA (Brian F. Glenister, William M. Furnish, & Zhou Zuren)	
Superfamily MEDLICOTTIOIDEA	
Family PRONORITIDAE	193
Family MEDLICOTTIIDAE	198

215
216
217
220
253

## INFORMATION ON TREATISE VOLUMES

Parts of the *Treatise* are distinguished by assigned letters with a view to indicating their systematic sequence while allowing publication of units in whatever order each is made ready for the press. Copies can be obtained from the Paleontological Institute, The University of Kansas, 1475 Jayhawk Blvd., Room 119, Lawrence, Kansas 66045-7613, USA, www.paleo. ku.edu.

#### PUBLISHED VOLUMES

- Part A. INTRODUCTION: Fossilization (Taphonomy), Biogeography, and Biostratigraphy, xxiii + 569 p., 169 fig., 1979.
- Part B. PROTOCTISTA 1 (Charophyta), xvi + 170 p., 79 fig., 9 tables, 2005.
- Part C. PROTISTA 2 (Sarcodina, Chiefly "Thecamoebians" and Foraminiferida), Volumes 1 and 2, xxxi + 900 p., 653 fig., 1964.
- Part D. PROTISTA 3 (Protozoa: Chiefly Radiolaria, Tintinnina), xii + 195 p., 92 fig., 1954.
- Part E. Archaeocyatha and Porifera, xviii + 122 p., 89 fig., 1955.
- Part E, Revised. Archaeocyatha, Volume 1, xxx + 158 p., 107 fig., 1972.
- Part E, Revised. PORIFERA, Volume 2 (Introduction to the Porifera), xxvii + 349 p., 135 fig., 10 tables, 2003.
- Part E, Revised. PORIFERA, Volume 3 (Demospongea, Hexactinellida, Heteractinida, Calcarea), xxxi + 872 p., 506 fig., 1 table, 2004.
- Part F. COELENTERATA, xx + 498 p., 358 fig., 1956.
- Part F. COELENTERATA, Supplement 1 (Rugosa and Tabulata), Volumes 1 and 2, xl + 762 p., 462 fig., 1981.
- Part G. BRYOZOA, xiii + 253 p., 175 fig., 1953.
- Part G, Revised. BRYOZOA, Volume 1 (Introduction, Order Cystoporata, Order Cryptostomata), xxvi + 625 p., 295 fig., 1983.
- Part H. BRACHIOPODA, Volumes 1 and 2, xxxii + 927 p., 746 fig., 1965.
- Part H, Revised. BRACHIOPODA, Volume 1 (Introduction), xx + 539 p., 417 fig., 40 tables, 1997.
- Part H, Revised. BRACHIOPODA, Volumes 2 and 3 (Linguliformea, Craniiformea, Rhynchonelliformea [part]), xxx + 919 p., 616 fig., 17 tables, 2000.
- Part H, Revised. BRACHIOPODA, Volume 4 (Rhynchonelliformea [part]), xxxix + 768 p., 484 fig., 3 tables, 2002.
- Part H, Revised. BRACHIOPODA, Volume 5 (Rhynchonelliformea [part]), xlvi + 631 p., 398 fig., 2006.
- Part H, Revised. BRACHIOPODA, Volume 6 (Supplement), l + 906 p., 461 fig., 38 tables, CD of compiled references from volumes 1–6, 2007.
- Part I. MOLLUSCA 1 (Mollusca General Features, Scaphopoda, Amphineura, Monoplacophora, Gastropoda General Features, Archaeogastropoda, Mainly Paleozoic Caenogastropoda and Opisthobranchia), xxiii + 351 p., 216 fig., 1960.
- Part K. MOLLUSCA 3 (Cephalopoda General Features, Endoceratoidea, Actinoceratoidea, Nautiloidea, Bactritoidea), xxviii + 519 p., 361 fig., 1964.

Part L. MOLLUSCA 4 (Cephalopoda: Ammonoidea), xxii + 490 p., 558 fig., 1957.

Part L, Revised. MOLLUSCA 4, Volume 4 (Cretaceous Ammonoidea), xx + 362 p., 216 fig., 1996.

- Part N. MOLLUSCA 6 (Bivalvia), Volumes 1 and 2 (of 3), xxxvii + 952 p., 613 fig., 1969; Volume 3, iv + 272 p., 153 fig., 1971.
- Part O. ARTHROPODA 1 (Arthropoda General Features, Protarthropoda, Euarthropoda General Features, Trilobitomorpha), xix + 560 p., 415 fig., 1959.
- Part O, Revised. ARTHROPODA 1 (Trilobita: Introduction, Order Agnostida, Order Redlichiida), xxiv + 530 p., 309 fig., 1997.
- Part P. ARTHROPODA 2 (Chelicerata, Pycnogonida, Palaeoisopus), xvii + 181 p., 123 fig., 1955 [1956].
- Part Q. ARTHROPODA 3 (Crustacea, Ostracoda), xxiii + 442 p., 334 fig., 1961.
- Part R. ARTHROPODA 4, Volumes 1 and 2 (Crustacea Exclusive of Ostracoda, Myriapoda, Hexapoda), xxxvi + 651 p., 397 fig., 1969.
- Part R. ARTHROPODA 4, Volumes 3 and 4 (Hexapoda), xxii + 655 p., 265 fig., 1992.
- Part S. ECHINODERMATA 1 (Echinodermata General Features, Homalozoa, Crinozoa, exclusive of Crinoidea), Volumes 1 and 2, xxx + 650 p., 400 fig., 1967 [1968].
- Part T. ECHINODERMATA 2 (Crinoidea), Volumes 1-3, xxxviii + 1,027 p., 619 fig., 1978.
- Part U. ECHINODERMATA 3 (Asterozoans, Echinozoans), xxx + 695 p., 534 fig., 1966.
- Part V. GRAPTOLITHINA, xvii + 101 p., 72 fig., 1955.
- Part V, Revised. GRAPTOLITHINA, xxxii + 163 p., 109 fig., 1970.
- Part W. MISCELLANEA (Conodonts, Conoidal Shells of Uncertain Affinities, Worms, Trace Fossils, Problematica), xxv + 259 p., 153 fig., 1962.
- Part W, Revised. MISCELLANEA, Supplement 1 (Trace Fossils and Problematica), xxi + 269 p., 110 fig., 1975.
- Part W, Revised. MISCELLANEA, Supplement 2 (Conodonta), xxviii + 202 p., frontis., 122 fig., 1981.

#### THIS VOLUME

Part L, Revised. MOLLUSCA 4, Volume 2 (Carboniferous and Permian Ammonoidea), xxix + 258 p., 139 fig., 1 table, 2009.

#### VOLUMES IN PREPARATION

- Part B. PROTISTA 1 (Chrysomonadida, Coccolithophorida, Diatomacea, etc.).
- Part E, Revised. PORIFERA (additional volumes).
- Part F, Revised. CNIDARIA (Scleractinia).
- Part G, Revised. BRYOZOA (additional volumes).
- Part K, Revised. MOLLUSCA 3 (Nautiloidea).
- Part L, Revised. MOLLUSCA 4 (Ammonoidea) (additional volumes).
- Part M. MOLLUSCA 5 (Coleoidea).
- Part O, Revised. ARTHROPODA 1 (Trilobita) (additional volumes).
- Part P, Revised. ARTHROPODA 2 (Chelicerata).
- Part Q, Revised. ARTHROPODA 3 (Ostracoda).
- Part R, Revised. ARTHROPODA 4 (Crustacea Exclusive of Ostracoda).
- Part T, Revised. ECHINODERMATA 2 (Crinoidea).
- Part V, Revised. GRAPTOLITHINA.
- Part W, Revised. TRACE FOSSILS.

## EDITORIAL PREFACE

PAUL A. SELDEN [The University of Kansas]

From the outset the aim of the Treatise on Invertebrate Paleontology has been to present a comprehensive and authoritative yet compact statement of knowledge concerning groups of invertebrate fossils. Typically, preparation of early Treatise volumes was undertaken by a small group with a synoptic view of the taxa being monographed. Two or perhaps three specialists worked together, sometimes co-opting others for coverage of highly specialized taxa. Recently, however, both new Treatise volumes and revisions of existing ones have been undertaken increasingly by teams of specialists led by a coordinating author. This volume, Part L, Revised, Ammonoidea, Volume 2, has been prepared by a team of specialists, each taking equal part in the work involved in this producing this revision. Editorial matters specific to this volume are discussed near the end of this editorial preface.

#### ZOOLOGICAL NAMES

Questions about the proper use of zoological names arise continually, especially questions regarding both the acceptability of names and alterations of names that are allowed or even required. Regulations prepared by the International Commission on Zoological Nomenclature (ICZN) and published in 1999 in the International Code of Zoological Nomenclature, hereinafter referred to as the Code, provide procedures for answering such questions. The prime objective of the Code is to promote stability and universality in the use of the scientific names of animals, ensuring also that each generic name is distinct and unique, while avoiding unwarranted restrictions on freedom of thought and action of systematists. Priority of names is a basic principle of the *Code;* but, under specified conditions and by following prescribed procedures, priority may be set aside by the Commission. These procedures apply especially where slavish adherence to the principle of priority would hamper or even disrupt zoological nomenclature and the information it conveys.

The Commission, ever aware of the changing needs of systematists, revised the Code in 1999 to enhance further nomenclatorial stability, specifying that the revised Code should take effect at the start of 2000. Among other requirements, the revised Code is clear in Chapter 14 that the type genus of family-level taxa must be specified. In this volume we have continued the practice that has characterized most previous volumes of the Treatise, namely that the type genus of all family-level taxa is the first listed and diagnosed. In spite of the revisions, the nomenclatorial tasks that confront zoological taxonomists are formidable and have often justified the complaint that the study of zoology and paleontology is too often merely the study of names rather than the study of animals. It is incumbent upon all systematists, therefore, at the outset of their work to pay careful attention to the Code to enhance stability by minimizing the number of subsequent changes of names, too many of which are necessitated by insufficient attention to detail. To that end, several pages here are devoted to aspects of zoological nomenclature that are judged to have chief importance in relation to procedures adopted in the Treatise, especially in this volume. Terminology is explained, and examples are given of the style employed in the nomenclatorial parts of the systematic descriptions.

## GROUPS OF TAXONOMIC CATEGORIES

Each taxon belongs to a category in the Linnaean hierarchical classification. The Code recognizes three groups of categories, a species-group, a genus-group, and a family-group. Taxa of lower rank than subspecies are excluded from the rules of zoological nomenclature, and those of higher rank than superfamily are not regulated by the Code. It is both natural and convenient to discuss nomenclatorial matters in general terms first and then to consider each of these three, recognized groups separately. Especially important is the provision that within each group the categories are coordinate, that is, equal in rank, whereas categories of different groups are not coordinate.

### FORMS OF NAMES

All zoological names can be considered on the basis of their spelling. The first form of a name to be published is defined as the original spelling (*Code*, Article 32), and any form of the same name that is published later and is different from the original spelling is designated a subsequent spelling (*Code*, Article 33). Not every original or subsequent spelling is correct.

#### **ORIGINAL SPELLINGS**

If the first form of a name to be published is consistent and unambiguous, the original is defined as correct unless it contravenes some stipulation of the *Code* (Articles 11, 27 to 31, and 34) or unless the original publication contains clear evidence of an inadvertent error in the sense of the *Code*, or, among names belonging to the familygroup, unless correction of the termination or the stem of the type genus is required. An original spelling that fails to meet these requirements is defined as incorrect.

If a name is spelled in more than one way in the original publication, the form adopted by the first reviser is accepted as the correct original spelling, provided that it complies with mandatory stipulations of the *Code* (Articles 11 and 24 to 34).

Incorrect original spellings are any that fail to satisfy requirements of the Code, represent an inadvertent error, or are one of multiple original spellings not adopted by a first reviser. These have no separate status in zoological nomenclature and, therefore, cannot enter into homonymy or be used as replacement names. They call for correction. For example, a name originally published with a diacritical mark, apostrophe, dieresis, or hyphen requires correction by deleting such features and uniting parts of the name originally separated by them, except that deletion of an umlaut from a vowel in a name derived from a German word or personal name unfortunately requires the insertion of e after the vowel. Where original spelling is judged to be incorrect solely because of inadequacies of the Greek or Latin scholarship of the author, nomenclatorial changes conflict with the primary purpose of zoological nomenclature as an information retrieval system. One looks forward with hope to further revisions of the Code wherein rules are emplaced that enhance stability rather than classical scholarship, thereby facilitating access to information.

#### SUBSEQUENT SPELLINGS

If a subsequent spelling differs from an original spelling in any way, even by the omission, addition, or alteration of a single letter, the subsequent spelling must be defined as a different name. Exceptions include such changes as an altered termination of adjectival specific names to agree in gender with associated generic names (an unfortunate impediment to stability and retrieval of information); changes of family-group names to denote assigned taxonomic rank; and corrections that eliminate originally used diacritical marks, hyphens, and the like. Such changes are not regarded as spelling changes conceived to produce a different name. In some instances, however, speciesgroup names having variable spellings are regarded as homonyms as specified in the *Code* (Article 58).

Altered subsequent spellings other than the exceptions noted may be either intentional or unintentional. If "demonstrably intentional" (*Code*, Article 33), the change is designated as an emendation. Emendations may be either justifiable or unjustifiable. Justifiable emendations are corrections of incorrect original spellings, and these take the authorship and date of the original spellings. Unjustifiable emendations are names having their own status in nomenclature, with author and date of their publication. They are junior, objective synonyms of the name in its original form.

Subsequent spellings, if unintentional, are defined as incorrect subsequent spellings. They have no status in nomenclature, do not enter into homonymy, and cannot be used as replacement names.

## AVAILABLE AND UNAVAILABLE NAMES

Editorial prefaces of some previous volumes of the Treatise have discussed in appreciable detail the availability of the many kinds of zoological names that have been proposed under a variety of circumstances. Much of that information, while important, does not pertain to the present volume, in which authors have used fewer terms for such names. The reader is referred to the Code (Articles 10 to 20) for further details on availability of names. Here, suffice it to say that an available zoological name is any that conforms to all mandatory provisions of the Code. All zoological names that fail to comply with mandatory provisions of the Code are unavailable and have no status in zoological nomenclature. Both available and unavailable names are classifiable into groups that have been recognized in previous volumes of the *Treatise*, although not explicitly differentiated in the *Code*. Among names that are available, these groups include inviolate names, perfect names, imperfect names, vain names, transferred names, improved or corrected names, substitute names, and conserved names. Kinds of unavailable names include naked names (see *nomina nuda* below), denied names, impermissible names, null names, and forgotten names.

Nomina nuda include all names that fail to satisfy provisions stipulated in Article 11 of the Code, which states general requirements of availability. In addition, they include names published before 1931 that were unaccompanied by a description, definition, or indication (Code, Article 12) and names published after 1930 that (1) lacked an accompanying statement of characters that differentiate the taxon, (2) were without a definite bibliographic reference to such a statement, (3) were not proposed expressly as a replacement (nomen novum) of a preexisting available name (Code, Article 13.1), or (4) for genus-group names, were unaccompanied by definite fixation of a type species by original designation or indication (Code, Article 13.2). Nomina nuda have no status in nomenclature, and they are not correctable to establish original authorship and date.

### VALID AND INVALID NAMES

Important considerations distinguish valid from available names on the one hand and invalid from unavailable names on the other. Whereas determination of availability is based entirely on objective considerations guided by articles of the *Code*, conclusions as to validity of zoological names may be partly subjective. A valid name is the correct one for a given taxon, which may have two or more available names but only a single correct, hence valid, name, which is also generally the oldest name that it has been given. Obviously, no valid name can also be an unavailable name, but invalid names may be either available or unavailable. It follows that any name for a given taxon other than the valid name, whether available or unavailable, is an invalid name.

One encounters a sort of nomenclatorial no-man's land in considering the status of such zoological names as *nomina dubia* (doubtful names), which may include both available and unavailable names. The unavailable ones can well be ignored, but names considered to be available contribute to uncertainty and instability in the systematic literature. These can ordinarily be removed only by appeal to the ICZN for special action. Because few systematists care to seek such remedy, such invalid but available names persist in the literature.

## NAME CHANGES IN RELATION TO GROUPS OF TAXONOMIC CATEGORIES SPECIES-GROUP NAMES

Detailed consideration of valid emendation of specific and subspecific names is unnecessary here, both because the topic is well understood and relatively inconsequential and because the Treatise deals with genus-group names and higher categories. When the form of adjectival specific names is changed to agree with the gender of a generic name in transferring a species from one genus to another, one need never label the changed name as nomen correctum. Similarly, transliteration of a letter accompanied by a diacritical mark in the manner now called for by the Code, as in changing originally bröggeri to broeggeri, or eliminating a hyphen, as in changing originally published cornu-oryx to cornuoryx, does not require the designation nomen correctum. Of course, in this age of computers and electronic databases, such changes of name, which are perfectly valid for the purposes of scholarship, run counter to the requirements of nomenclatorial stability upon which the preparation of massive, electronic databases is predicated.

#### **GENUS-GROUP NAMES**

Conditions warranting change of the originally published, valid form of generic and subgeneric names are sufficiently rare that lengthy discussion is unnecessary. Only elimination of diacritical marks and hyphens in some names in this category and replacement of homonyms seem to furnish basis for valid emendation. Many names that formerly were regarded as homonyms are no longer so regarded, because two names that differ only by a single letter or in original publication by the presence of a diacritical mark in one are now construed to be entirely distinct (but see *Code*, Article 58).

As has been pointed out above, difficulty typically arises when one tries to decide whether a change of spelling of a name by a subsequent author was intentional or unintentional, and the decision has to be made often arbitrarily.

#### FAMILY-GROUP NAMES

Family-Group Names: Authorship and Date

All family-group taxa having names based on the same type genus are attributed to the author who first published the name of any of these groups, whether tribe, subfamily, or family (superfamily being almost inevitably a later-conceived taxon). Accordingly, if a family is divided into subfamilies or a subfamily into tribes, the name of no such subfamily or tribe can antedate the family name. Moreover, every family containing differentiated subfamilies must have a nominate subfamily (sensu stricto), which is based on the same type genus as the family. Finally, the author and date set down for the nominate subfamily invariably are identical with those of the family, irrespective of whether the author of the family or some subsequent author introduced subdivisions.

Corrections in the form of family-group names do not affect authorship and date of the taxon concerned, but in the *Treatise* recording the authorship and date of the correction is desirable because it provides a pathway to follow the thinking of the systematists involved.

#### Family-Group Names: Use of *nomen translatum*

The Code (Article 29.2) specifies the suffixes for tribe (-ini), subfamily (-inae), family (-idae) and superfamily (-oidea), the formerly widely used ending (-acea) for superfamily having been disallowed. All these family-group categories are defined as coordinate (Code, Article 36.1): "A name established for a taxon at any rank in the family group is deemed to have been simultaneously established for nominal taxa at other ranks in the family group; all these taxa have the same type genus, and their names are formed from the stemof the name of the type genus (Art. 29.3] with appropriate change of suffix [Art. 34.1]. The name has the same authorship and date at every rank." Such changes of rank and concomitant changes of endings as elevation of a subfamily to family rank or of a family to superfamily rank, if introduced subsequent to designation of the original taxon or based on the same nominotypical genus, are nomina translata. In the Treatise it is desirable to distinguish the valid alteration in the changed ending of each transferred family-group name by the term nomen translatum, abbreviated to nom. transl. Similarly for clarity, authors should record the author, date, and page of the alteration, as in the following example.

#### Family HEXAGENITIDAE Lameere, 1917

[nom. transl. DEMOULIN, 1954, p. 566, ex Hexagenitinae LAMEERE, 1917, p. 74]

This is especially important for superfamilies, for the information of interest is the author who initially introduced a taxon rather than the author of the superfamily as defined by the *Code*. For example:

## Superfamily AGNOSTOIDEA M'Coy, 1849

[nom. transl. SHERGOLD, LAURIE, & SUN, 1990, p. 32, ex Agnostinae M'Coy, 1849, p. 402]

The latter is merely the individual who first defined some lower-ranked, family-group taxon that contains the nominotypical genus of the superfamily. On the other hand, the publication that introduces the superfamily by *nomen translatum* is likely to furnish the information on taxonomic considerations that support definition of the taxon.

#### Family-Group Names: Use of *nomen correctum*

Valid name changes classed as *nomina correcta* do not depend on transfer from one category of the family group to another but most commonly involve correction of the stem of the nominotypical genus. In addition, they include somewhat arbitrarily chosen modifications of endings for names of tribes or superfamilies. Examples of the use of *nomen correctum* are the following.

#### Family STREPTELASMATIDAE Nicholson, 1889

[nom. correct. WEDEKIND, 1927, p. 7, pro Streptelasmidae NICHOLSON in NICHOLSON & LYDEKKER, 1889, p. 297]

#### Family PALAEOSCORPIDAE Lehmann, 1944

[nom. correct. PETRUNKEVITCH, 1955, p. 73, pro Palaeoscorpionidae LEHMANN, 1944, p. 177]

#### Family-Group Names: Replacements

Family-group names are formed by adding combinations of letters, which are prescribed for all family-group categories, to the stem of the name belonging to the nominotypical genus first chosen as type of the assemblage. The type genus need not be the first genus in the family to have been named and defined, but among all those included it must be the first published as name giver to a familygroup taxon. Once fixed, the family-group name remains tied to the nominotypical genus even if the generic name is changed by reason of status as a junior homonym or junior synonym, either objective or subjective. Seemingly, the Code requires replacement of a family-group name only if the nominotypical genus is found to have been a junior homonym when it was proposed (Code, Article 39), in which case "... it must be replaced either by the next oldest available name from among its synonyms [Art. 23.3.5], including the names of its subordinate family-group taxa, or, if there is no such synonym, by a new name based on the valid name . . . of the former type genus." Authorship and date attributed to the replacement family-group name are determined by first publication of the changed family-group name. Recommendation 40A of the Code, however, specifies that for subsequent application of the rule of priority, the family-group name ". . . should be cited with its original author and date (see Recommendation 22A.2.2), followed by the date of its priority as determined by this Article; the date of priority should be enclosed in parentheses." Many family-group names that have been in use for a long time are nomina nuda, since they fail to satisfy criteria of availability (Code, Article 11.7). These demand replacement by valid names.

The aim of family-group nomenclature is to yield the greatest possible stability and uniformity, just as in other zoological names. Both taxonomic experience and the Code (Article 40) indicate the wisdom of sustaining family-group names based on junior subjective synonyms if they have priority of publication, for opinions of the same worker may change from time to time. The retention of first-published, family-group names that are found to be based on junior objective synonyms, however, is less clearly desirable, especially if a replacement name derived from the senior objective synonym has been recognized very long and widely. Moreover, to displace a widely used, familygroup name based on the senior objective synonym by disinterring a forgotten and

virtually unused family-group name based on a junior objective synonym because the latter happens to have priority of publication is unsettling.

A family-group name may need to be replaced if the nominotypical genus is transferred to another family group. If so, the first-published of the generic names remaining in the family-group taxon is to be recognized in forming a replacement name.

#### SUPRAFAMILIAL TAXA: TAXA ABOVE FAMILY-GROUP

International rules of zoological nomenclature as given in the Code affect only lower-rank categories: subspecies to superfamily. Suprafamilial categories (suborder to kingdom) are either not mentioned or explicitly placed outside of the application of zoological rules. The Copenhagen Decisions on Zoological Nomenclature (1953, Articles 59 to 69) proposed adopting rules for naming suborders and higher taxa up to and including phylum, with provision for designating a type genus for each, in such manner as not to interfere with the taxonomic freedom of workers. Procedures were outlined for applying the rule of priority and rule of homonymy to suprafamilial taxa and for dealing with the names of such taxa and their authorship, with assigned dates, if they should be transferred on taxonomic grounds from one rank to another. The adoption of terminations of names, different for each category but uniform within each, was recommended.

The Colloquium on Zoological Nomenclature, which met in London during the week just before the 15th International Congress of Zoology convened in 1958, thoroughly discussed the proposals for regulating suprafamilial nomenclature, as well as many others advocated for inclusion in the new *Code* or recommended for exclusion from it. A decision that was supported by a wide majority of the participants in the colloquium was against the establishment of rules for naming taxa above family-group rank, mainly because it was judged that such regulation would unwisely tie the hands of taxonomists. For example, a class or order defined by an author at a given date, using chosen morphologic characters (e.g., gills of bivalves), should not be allowed to freeze nomenclature, taking precedence over another class or order that is proposed later and distinguished by different characters (e.g., hinge teeth of bivalves). Even the fixing of type genera for suprafamilial taxa would have little, if any, value, hindering taxonomic work rather than aiding it. Beyond mere tidying up, no basis for establishing such types and for naming these taxa has yet been provided.

The considerations just stated do not prevent the editors of the Treatise from making rules for dealing with suprafamilial groups of animals described and illustrated in this publication. Some uniformity is needed, especially for the guidance of *Treatise* authors. This policy should accord with recognized general practice among zoologists; but where general practice is indeterminate or nonexistent, our own procedure in suprafamilial nomenclature needs to be specified as clearly as possible. This pertains especially to decisions about names themselves, about citation of authors and dates, and about treatment of suprafamilial taxa that, on taxonomic grounds, are changed from their originally assigned rank. Accordingly, a few rules expressing Treatise policy are given here, some with examples of their application.

1. The name of any suprafamilial taxon must be a Latin or Latinized, uninominal noun of plural form or treated as such, with a capital initial letter and without diacritical mark, apostrophe, diaeresis, or hyphen. If a component consists of a numeral, numerical adjective, or adverb, this must be written in full.

2. Names of suprafamilial taxa may be constructed in almost any manner. A name may indicate morphological attributes (*e.g.*, Lamellibranchiata, Cyclostomata, Toxoglossa) or be based on the stem of an included genus (e.g., Bellerophontina, Nautilida, Fungiina) or on arbitrary combinations of letters (e.g., Yuania); none of these, however, can end in -idae or -inae, which terminations are reserved for family-group taxa. No suprafamilial name identical in form to that of a genus or to another published suprafamilial name should be employed (e.g., order Decapoda LATREILLE, 1803, crustaceans, and order Decapoda LEACH, 1818, cephalopods; suborder Chonetoidea MUIR-WOOD, 1955, and genus Chonetoidea JONES, 1928). Worthy of notice is the classificatory and nomenclatorial distinction between suprafamilial and family-group taxa that, respectively, are named from the same type genus, since one is not considered to be transferable to the other (e.g., suborder Bellerophontina ULRICH & SCOFIELD, 1897 is not coordinate with superfamily Bellerophontacea McCoy, 1851 or family Bellerophontidae McCoy, 1851).

3. The rules of priority and homonymy lack any force of international agreement as applied to suprafamilial names, yet in the interest of nomenclatorial stability and to avoid confusion these rules are widely applied by zoologists to taxa above the family-group level wherever they do not infringe on taxonomic freedom and long-established usage.

4. Authors who accept priority as a determinant in nomenclature of a suprafamilial taxon may change its assigned rank at will, with or without modifying the terminal letters of the name, but such changes cannot rationally be judged to alter the authorship and date of the taxon as published originally. A name revised from its previously published rank is a transferred name (*nomen translatum*), as illustrated in the following.

## Order CORYNEXOCHIDA Kobayashi, 1935

[nom. transl. MOORE, 1959, p. 217, ex suborder Corynexochida KOBAYASHI, 1935, p. 81]

A name revised from its previously published form merely by adoption of a different termination without changing taxonomic rank is a *nomen correctum*.

## Order DISPARIDA Moore & Laudon, 1943

[nom. correct. MOORE in MOORE, LALICKER, & FISCHER, 1952, p. 613, pro order Disparata MOORE & LAUDON, 1943, p. 24]

A suprafamilial name revised from its previously published rank with accompanying change of termination, which signals the change of rank, is recorded as a *nomen translatum et correctum*.

## Order HYBOCRINIDA Jaekel, 1918

[nom. transl. et correct. MOORE in MOORE, LALICKER, & FISCHER, 1952, p. 613, ex suborder Hybocrinites JAEKEL, 1918, p. 90]

5. The authorship and date of nominate subordinate and supraordinate taxa among suprafamilial taxa are considered in the *Treatise* to be identical since each actually or potentially has the same type. Examples are given below.

## Subclass ENDOCERATOIDEA Teichert, 1933

[nom. transl. TEICHERT in TEICHERT & others, 1964, p. 128, ex order Endoceroidea TEICHERT, 1933, p. 214]

## Order ENDOCERIDA Teichert, 1933

[nom. correct. TEICHERT in TEICHERT & others, 1964, p. 165, pro order Endoceroidea TEICHERT, 1933, p. 214]

### TAXONOMIC EMENDATION

Emendation has two distinct meanings as regards zoological nomenclature. These are alteration of a name itself in various ways for various reasons, as has been reviewed, and alteration of the taxonomic scope or concept for which a name is used. The *Code* (Article 33.1 and Glossary) concerns itself only with the first type of emendation, applying the term to intentional, either justified or unjustified changes of the original spelling of a name. The second type of emendation primarily concerns classification and inherently is not associated with change of name. Little attention generally has been paid to this distinction in spite of its significance.

Most zoologists, including paleontologists, who have emended zoological names refer to what they consider a material change in application of the name such as may be expressed by an importantly altered diagnosis of the assemblage covered by the name. The abbreviation emend. then must accompany the name with statement of the author and date of the emendation. On the other hand, many systematists think that publication of emend. with a zoological name is valueless because alteration of a taxonomic concept is introduced whenever a subspecies, species, genus, or other taxon is incorporated into or removed from a higher zoological taxon. Inevitably associated with such classificatory expansions and restrictions is some degree of emendation affecting diagnosis. Granting this, still it is true that now and then somewhat more extensive revisions are put forward, generally with a published statement of the reasons for changing the application of a name. To erect a signpost at such points of most significant change is worthwhile, both as an aid to subsequent workers in taking account of the altered nomenclatorial usage and to indicate where in the literature cogent discussion may be found. Authors of contributions to the Treatise are encouraged to include records of all especially noteworthy emendations of this nature, using the abbreviation emend. with the name to which it refers and citing the author, date, and page of the emendation. Examples from Treatise volumes follow.

## Order ORTHIDA Schuchert & Cooper, 1932

[nom. transl. et correct. MOORE in MOORE, LALICKER, & FISCHER, 1952, p. 220, ex suborder Orthoidea SCHUCHERT & COOPER, 1932, p. 43; emend., WILLIAMS & WRIGHT, 1965, p. 299]

#### Subfamily ROVEACRININAE Peck, 1943

[Roveacrininae Реск, 1943, р. 465; *emend.*, Реск in Мооке & Теіснегт, 1978, р. 921]

## STYLE IN GENERIC DESCRIPTIONS CITATION OF TYPE SPECIES

In the *Treatise* the name of the type species of each genus and subgenus is given immediately following the generic name with its accompanying author, date, and page reference or after entries needed for definition of the name if it is involved in homonymy. The originally published combination of generic and trivial names of this species is cited, accompanied by an asterisk (\*), with notation of the author, date, and page of original publication, except if the species was first published in the same paper and by the same author as that containing definition of the genus of which it is the type. In this instance, the initial letter of the generic name followed by the trivial name is given without repeating the name of the author and date. Examples of these two sorts of citations follow.

- Orionastraea SMITH, 1917, p. 294 [\*Sarcinula phillipsi McCoy, 1849, p. 125; OD].
- Schoenophyllum SIMPSON, 1900, p. 214 [\*S. aggregatum; OD].

If the cited type species is a junior synonym of some other species, the name of this latter is given also, as follows.

Actinocyathus D'ORBIGNY, 1849, p. 12 [\**Cyathophyllum crenulate* PHILLIPS, 1836, p. 202; M; =*Lons-daleia floriformis* (MARTIN), 1809, pl. 43; validated by ICZN Opinion 419].

In some instances the type species is a junior homonym. If so, it is cited as shown in the following example.

Prionocyclus MEEK, 1871b, p. 298 [\*Ammonites serratocarinatus MEEK, 1871a, p. 429, non STOLICZKA, 1864, p. 57; =Prionocyclus wyomingensis MEEK, 1876, p. 452].

In the *Treatise* the name of the type species is always given in the exact form it had in the original publication except that diacritical marks have been removed. Where other mandatory changes are required, these are introduced later in the text, typically in the description of a figure.

#### Fixation of Type Species Originally

It is desirable to record the manner of establishing the type species, whether by original designation (OD) or by subsequent designation (SD). The type species of a genus or subgenus, according to provisions of the Code, may be fixed in various ways in the original publication; or it may be fixed subsequently in ways specified by the Code (Article 68) and described in the next section. Type species fixed in the original publication include (1) original designation (in the Treatise indicated by OD) when the type species is explicitly stated or (before 1931) indicated by n. gen., n. sp. (or its equivalent) applied to a single species included in a new genus, (2) defined by use of typus or typicus for one of the species included in a new genus (adequately indicated in the Treatise by the specific name), (3) established by monotypy if a new genus or subgenus has only one originally included species (in the Treatise indicated as M), and (4) fixed by tautonymy if the genus-group name is identical to an included species name not indicated as the type.

#### Fixation of Type Species Subsequently

The type species of many genera are not determinable from the publication in which the generic name was introduced. Therefore, such genera can acquire a type species only by some manner of subsequent designation. Most commonly this is established by publishing a statement naming as type species one of the species originally included in the genus. In the *Treatise* such fixation of the type species by subsequent designation in this manner is indicated by the letters SD accompanied by the name of the subsequent author (who may be the same person as the original author) and the publication date and page number of the subsequent designation. Some genera, as first described and named, included no mentioned species (for such genera established after 1930, see below); these necessarily lack a type species until a date subsequent to that of the original publication when one or more species is assigned to such a genus. If only a single species is thus assigned, it becomes automatically the type species. Of course, the first publication containing assignment of species to the genus that originally lacked any included species is the one concerned in fixation of the type species, and if this publication names two or more species as belonging to the genus but did not designate a type species, then a later SD designation is necessary. Examples of the use of SD as employed in the Treatise follow.

- Hexagonaria GURICH, 1896, p. 171 [\**Cyathophyllum hexagonum* GOLDFUSS, 1826, p. 61; SD LANG, SMITH, & THOMAS, 1940, p. 69].
- Mesephemera HANDLIRSCH, 1906, p. 600 [\**Tineites lithophilus* GERMAR, 1842, p. 88; SD CARPENTER, herein].

Another mode of fixing the type species of a genus is through action of the International Commission of Zoological Nomenclature using its plenary powers. Definition in this way may set aside application of the *Code* so as to arrive at a decision considered to be in the best interest of continuity and stability of zoological nomenclature. When made, it is binding and commonly is cited in the *Treatise* by the letters ICZN, accompanied by the date of announced decision and reference to the appropriate numbered opinion.

Subsequent designation of a type species is admissible only for genera established prior to 1931. A new genus-group name established after 1930 and not accompanied by fixation of a type species through original designation or original indication is invalid (*Code*, Article 13.3). Effort of a subsequent author to validate such a name by subsequent designation of a type species constitutes an original publication making the name available under authorship and date of the subsequent author.

#### HOMONYMS

Most generic names are distinct from all others and are indicated without ambiguity by citing their originally published spelling accompanied by name of the author and date of first publication. If the same generic name has been applied to two or more distinct taxonomic units, however, it is necessary to differentiate such homonyms. This calls for distinction between junior homonyms and senior homonyms. Because a junior homonym is invalid, it must be replaced by some other name. For example, Callophora HALL, 1852, introduced for Paleozoic trepostomate bryozoans, is invalid because Gray in 1848 published the same name for Cretaceous-Holocene cheilostomate bryozoans. Bassler in 1911 introduced the new name Hallophora to replace Hall's homonym. The Treatise style of entry is given below.

Hallophora Bassler, 1911, p. 325, nom. nov. pro Callophora Hall, 1852, p. 144, non Gray, 1848.

In like manner, a replacement generic name that is needed may be introduced in the *Treatise* (even though first publication of generic names otherwise in this work is generally avoided). An exact bibliographic reference must be given for the replaced name as in the following example.

Mysterium DE LAUBENFELS, herein, nom. nov. pro Mystrium SCHRAMMEN, 1936, p. 183, non ROGER, 1862 [\*Mystrium porosum SCHRAMMEN, 1936, p. 183; OD].

Otherwise, no mention is made generally of the existence of a junior homonym.

#### Synonymous Homonyms

An author sometimes publishes a generic name in two or more papers of different date, each of which indicates that the name is new. This is a bothersome source of errors for later workers who are unaware that a supposed first publication that they have in hand is not actually the original one. Although the names were published separately, they are identical and therefore definable as homonyms; at the same time they are absolute synonyms. For the guidance of all concerned, it seems desirable to record such names as synonymous homonyms. In the *Treatise* the junior of one of these is indicated by the abbreviation *jr. syn. hom.* 

Not infrequently, identical family-group names are published as new names by different authors, the author of the name that was introduced last being ignorant of previous publication(s) by one or more other workers. In spite of differences in taxonomic concepts as indicated by diagnoses and grouping of genera and possibly in assigned rank, these family-group taxa, being based on the same type genus, are nomenclatorial homonyms. They are also synonyms. Wherever encountered, such synonymous homonyms are distinguished in the *Treatise* as in dealing with generic names.

A rare but special case of homonymy exists when identical family names are formed from generic names having the same stem but differing in their endings. An example is the family name Scutellidae RICHTER & RICHTER, 1925, based on *Scutellum* PUSCH, 1833, a trilobite. This name is a junior homonym of Scutellidae GRAY, 1825, based on the echinoid genus *Scutella* LAMARCK, 1816. The name of the trilobite family was later changed to Scutelluidae (ICZN, Opinion 1004, 1974).

#### **SYNONYMS**

In the *Treatise*, citation of synonyms is given immediately after the record of the type species. If two or more synonyms of differing date are recognized, these are arranged in chronological order. Objective synonyms are indicated by accompanying designation *obj.*, others being understood to constitute subjective synonyms, of which the types are also indicated. Examples showing *Treatise* style in listing synonyms follow.

Mackenziephyllum Pedder, 1971, p. 48 [\*M. insolitum; OD] [=Zonastraea TsyGANKO in SPASSKIY, KRAVTSOV, & TSYGANKO, 1971, p. 85, nom. nud.; =Zonastraea TsyGANKO, 1972, p. 21 (type, Z. graciosa, OD)]. Kodonophyllum WEDEKIND, 1927, p. 34 [\*Streptelasma Milne-Edwardsi DyBowski, 1873, p. 409; OD; =Madrepora truncata LINNE, 1758, p. 795, see SMITH & TREMBERTH, 1929, p. 368] [=Patrophontes LANG & SMITH, 1927, p. 456 (type, Madrepora truncata LINNE, 1758, p. 795, OD); =Codonophyllum LANG, SMITH, & THOMAS, 1940, p. 39, obj.].

Some junior synonyms of either the objective or the subjective sort may be preferred over senior synonyms whenever uniformity and continuity of nomenclature are served by retaining a widely used but technically rejectable name for a genus. This requires action of the ICZN, which may use its plenary powers to set aside the unwanted name, validate the wanted one, and place the concerned names on appropriate official lists.

## OTHER EDITORIAL MATTERS BIOGEOGRAPHY

Purists, Treatise editors among them, would like nothing better than a stable world with a stable geography that makes possible a stable biogeographical classification. Global events of the past few years have shown how rapidly geography can change, and in all likelihood we have not seen the last of such change as new, so-called republics continue to spring up all over the globe. One expects confusion among readers in the future as they try to decipher such geographical terms as USSR, Yugoslavia, or Ceylon. Such confusion is unavoidable, as books must be completed and published at some real time. Libraries would be limited indeed if publication were always to be delayed until the political world had settled down. In addition, such terms as central Europe and western Europe are likely to mean different things to different people. Some imprecision is introduced by the use of all such terms, of course, but it is probably no greater than the imprecision that stems from the fact that the work of paleontology is not yet finished, and the geographical ranges of many genera are imperfectly known.

Other geographic terms can also have varying degrees of formality. In general, *Treatise* policy is to use adjectives rather than nouns to refer to directions. Thus we have used *southern* and *western* in place of *South* and *West* unless a term has been formally defined as a geographic entity (e.g., South America or West Virginia). Note that we have referred to western Texas rather than West Texas, which is said to be not a state but a state of mind.

#### NAMES OF AUTHORS: TRANSLATION AND TRANSLITERATION

Chinese scientists have become increasingly active in systematic paleontology in the past two decades. Chinese names cause anguish among English-language bibliographers for two reasons. First, no scheme exists for one-to-one transliteration of Chinese characters into roman letters. Thus, a Chinese author may change the roman-letter spelling of his name from one publication to another. For example, the name Chang, the most common family name in the world reportedly held by some one billion people, has been spelled more recently Zhang. The principal purpose of a bibliography is to provide the reader with entry into the literature. Quite arbitrarily, therefore, in the interest of information retrieval, the Treatise editorial staff has decided to retain the roman spelling that a Chinese author has used in each of his publications rather than attempting to adopt a common spelling of an author's name to be used in all citations of his work. It is entirely possible, therefore, that the publications of a Chinese author may be listed in more than one place under more than one name in the bibliography.

Second, most but by no means all Chinese list their family name first followed by given names. People with Chinese names who study in the West, however, often reverse the order, putting the family name last as is the Western custom. Thus, for example, Dr. Yi-Maw Chang, formerly of the staff of the Paleontological Institute, was Chang Yi-Maw when he lived in Taiwan. When he came to America, he became Yi-Maw Chang. In the *Treatise*, authors' names are used in the text and listed in the references as they appear in the source being cited.

Several systems exist for transliterating the Cyrillic alphabet into the roman alphabet. On the recommendation of skilled bibliographic librarians, we have adopted the American Library Association/Library of Congress romanization table for Russian and other languages using the Cyrillic alphabet.

## MATTERS SPECIFIC TO THIS VOLUME

Authorship entails both credit and responsibility. As the knowledge of paleontology grows and paleontologists become more specialized, preparation of *Treatise* volumes must necessarily involve larger and larger teams of researchers, each focusing on increasingly narrow aspects of the higher taxon under revision. In this volume, we have taken special pains to acknowledge authorship of small subsections. Readers citing the volume are encouraged to pay close attention to the actual authorship of a section or subsection.

Stratigraphic ranges of taxa listed in the systematic descriptions herein have been compiled from the ranges of lower taxa. In all instances, we have used the *range-through* method of describing ranges. In instances, therefore, where the work of paleontology is not yet finished, some ranges of higher taxa will not show gaps between the ranges of their subtaxa and may seem to be more complete than the data warrant. In this volume, stratigraphic range charts typical of previous *Treatise* volumes present a much more precise picture of the biostratigraphy of the ammonoids (see Table 1, p. 220 herein).

#### ACKNOWLEDGMENTS

The Paleontological Institute's Assistant Editor for Text, Jill Hardesty, and the Assistant Editor for Illustrations, Jane Kerns, have faced admirably the formidable task of moving this volume through the various stages of editing and into production. In this they have been ably assisted by other members of the editorial team, including Mike Cormack with his outstanding computer skills, and Denise Mayse with her work on reference checking and general support.

This editorial preface and other, recent ones are extensive revisions of the prefaces prepared for previous *Treatise* volumes by former editors, including the late Raymond C. Moore, the late Curt Teichert, Richard A. Robison, and the late Roger L. Kaesler. I am indebted to them for preparing earlier prefaces and for the leadership they have provided in bringing the *Treatise* project to its present status.

#### REFERENCES

- International Commission on Zoological Nomenclature. 1999. International Code of Zoological Nomenclature, 4th edition. International Trust for Zoological Nomenclature. London. 306 p.
- Moore, R. C., and C. Teichert. 1978. Treatise on Invertebrate Paleontology. Part T, Echinodermata 2(1). The Geological Society of America and The University of Kansas. Boulder & Lawrence. 401 p.
- Robison, R. A., and C. Teichert. 1979. Treatise on Invertebrate Paleontology. Part A, Introduction. The Geological Society of America and The University of Kansas. Boulder & Lawrence. 569 p.

Paul A. Selden Lawrence, Kansas February 17, 2009

## STRATIGRAPHIC DIVISIONS

The major divisions of the geological time scale are reasonably well established throughout the world, but minor divisions (e.g., subseries, stages, and substages) are more likely to be provincial in application. The stratigraphic units listed here represent an authoritative version of the stratigraphic column for all taxonomic work relating to the revision of Part L (any provincial terms are presented in brackets in taxonomic descriptions). They are adapted from the International Stratigraphic Chart, compiled by the International Commission on Stratigraphy (ICS; ©2008).

**Cenozoic Erathem** Quaternary System Holocene Series **Pleistocene Series** Neogene System **Pliocene Series** Miocene Series Paleogene System **Oligocene Series Eocene Series Paleocene** Series Mesozoic Erathem Cretaceous System Upper Cretaceous Series Lower Cretaceous Series Jurassic System Upper Jurassic Series Middle Jurassic Series Lower Jurassic Series **Triassic System** Upper Triassic Series Middle Triassic Series Lower Triassic Series Paleozoic Erathem Permian System Lopingian Series Guadalupian Series Cisuralian Series

Carboniferous System Pennsylvanian Series Gzhelian Stage Kasimovian Stage Moscovian Stage Bashkirian Stage Mississippian Series Serpukhovian Stage Visean Stage Tournaisian Stage **Devonian System** Upper Devonian Series Middle Devonian Series Lower Devonian Series Silurian System Pridoli Series Ludlow Series Wenlock Series Llandovery Series Ordovician System Upper Ordovician Series Middle Ordovician Series Lower Ordovician Series **Cambrian System** Furongian Series Series 3 Series 2 Terreneuvian Series

## COORDINATING AUTHORS' PREFACE

JÜRGEN KULLMANN and ZHOU ZUREN

[University of Tübingen, Germany; and Nanjing Institute of Geology and Palaeontology]

In 1938, the first volume of a comprehensive treatment of every valid taxon (especially genus and species) in invertebrate paleozoology was published in the Handbuch der Paläozoologie by Otto H. Schindewolf (1938-1944). This very ambitious work resulted only in a few parts being issued: Graptolithina, Gastropoda, and Hydrozoa; it was stopped by World War II. Less than ten years later, Raymond C. Moore in Lawrence, Kansas, founded the Treatise on Invertebrate Paleontology, with the first volume, part G, Bryozoa, being issued in December 1953. In his Editorial Preface, R. C. Moore (1953, p. vii) defined the intent of the Treatise: "The aim of the Treatise on Invertebrate Paleontology, as originally conceived and consistently pursued, is to present the most comprehensive and authoritative, yet compact statement of knowledge concerning invertebrate fossil groups that can be formulated by collaboration of competent specialists."

The first volume dedicated solely to the Ammonoidea was published in 1957. It covered not only the different groups of ammonoid cephalopods, but included also chapters on morphology, evolution, paleoecology, and geographic distribution. The chapter on Paleozoic Ammonoidea was authored by A. K. Miller and W. M. Furnish, and embraces the treatment by O. H. Schindewolf (Clymeniida) and B. Kummel (Permian constituents of Ceratitida).

Between 1954 and 1965, more than 100 new genera were established worldwide. Therefore, in February 1967, plans to revise *Treatise* Part L were discussed in Lawrence, Kansas. At this meeting, W. M. Furnish, B. F. Glenister, J. Kullmann, and B. Kummel agreed upon exact assignments, based on taxonomy or stratigraphy. W. M. Furnish and B. F. Glenister would be responsible for all families occuring in the Permian except Xenodiscidae. J. Kullmann would work mostly with simpler Goniatitina, M. R. House would handle the bulk of Devonian forms, and O. H. Schindewolf the clymeniids. Despite a considerable expansion of the introductory portion, a single chapter would be processed by editor C. Teichert, who was appointed as coordinator of the volume. Originally, spring 1970 was considered as the target date for completion of manuscripts. The first to present a manuscript was Schindewolf in 1971, and the others were expected at latest by 1975.

Curt Teichert worked as coordinator of the ammonoid *Treatise* revision for more than a decade. Though not a specialist, Teichert was recognized for his wide and deep knowledge of the Cephalopoda and regarded as a superb generalist. A number of revised editions of parts of the *Treatise* were directed and edited by him, beginning with Part V, Graptolithina (1970). Teichert resigned from the *Treatise* editorship at end of 1976, but he continued working for the *Treatise* for several more years. A broad correspondence with the involved specialists ensured that things went smoothly, although slowly.

Several so-called first drafts were assembled. The Devonian chapter of 138 pages and more than 100 plates, by M. House; the Carboniferous part, 277 pages and 75 plates by Kullmann; the Permian part, 100 pages and more than 100 plates by Furnish and Glenister; and orthocone ammonoids, by R. T. Becker.

A great event for the *Treatise* was the international gathering in 1979 in London at the Systematics Association Symposium on the Ammonoidea, the first international symposium ever held to discuss scientific problems of the group. On this occasion, *Treatise* authors summarized in "Keynote Lectures" their views on the classification and evolution of the group. The final session of the symposium addressed the biostratigraphic usefulness of the ammonoids and the problem of correlation and distribution.

The advent of the computer with its enormous storage and retrieval capacity allowed another step to be taken to provide access to the treasures of information about fossils, including information about taxa insufficiently described and figured from earlier times. In 1990 the Database System GONIAT (and later GONIAT Online 2007) was founded, based on Treatise classification. After 15 years of work, about 4000 valid species and more than 700 genera and subgenera of about 120 families have been included, and approximately 7500 localities are described. The information provided in GONIAT is intensively used for the present volume. GONIAT can be found at www.goniat.org. Updates of the entire database system provided by Svetlana V. Nikolaeva, Scientific Editor, Bulletin of Zoological Nomenclature, International Commission on Zoological Nomenclature, The National Museum London, SW7 58D, U.K. (s.nikolaeva@nhm.ac.uk); technical assistance provided by Peter S. Kullmann, Humboldtstr. 32, D-70771 Leinfelden-Echterdingen, Germany (info@elementec. de).

In 2004, W. M. Furnish, B. F. Glenister, J. Kullmann, and Zhou Zuren decided to put together the Carboniferous and Permian parts and asked the Editorial Office at the Paleontological Institute of the University of Kansas in Lawrence to accept both unified parts for the *Treatise* series of revised parts. The uninterrupted evolutionary lines between the Carboniferous and Permian made possible a similar treatment of taxonomic descriptions and literature.

The present volume reflects a considerable amount of paleontological research of the last century. In 1988 Zhou Zuren joined the Iowa group of scientists specializing in Permian ammonoids. The new interpretation of the sutural pattern of the early stages of coiling, EALUI [German] and VLU:ID [Russian] instead of  $ELU_{2:}U_1$ :I or VUU<sup>1</sup>:ID in Prolecanitida may be of great consequence for the evaluation of phylogeny (p. 185 herein). It has led to the idea of a much closer affinity of the orders Prolecanitida and Goniatitida. Many representatives of aberrant endemics, mainly the pseudohaloritaceans, kufengoceratins, and paragastrioceratids reported from South China, suggest a widespread ecological differentiation among Permian ammonoids. The combination of these special faunas with special environments has led to a new understanding of ammonoid provincialism. Dimorphism is known to be a common phenomenon in the Permian period. A regressive evolution pattern existed in some late Paleozoic ammonoid families: a kind of terminal paedomorphosis has been verified in many genera and their family groups. As is explained in one of the old Treatise prefaces: the making of a reasonably complete inventory of present knowledge may be expected to yield needed foundation for future research, and it is hoped that the Treatise will serve this end.

#### ACKNOWLEDGMENTS

Many colleagues have supported our investigations with significant contributions to systematic descriptions and by providing essential illustrations. A large number of individuals and institutions have generously helped in bringing together extensive information on the Carboniferous and Permian ammonoids. The authors wish to express special thanks to R. T. Becker, Münster; V. E. Ruzhentsev,<sup>†</sup> B. I. Bogoslovskii,<sup>†</sup> M. F. Bogoslovskaia, and T. B. Leonova, Moscow; J. Dzik, Warzawa; M. R. House,<sup>†</sup> Hull; D. Weyer, and D. Korn, Berlin; L. F. Kuzina, Moscow; Liang Xi-luo and Ruan Yiping,<sup>†</sup> Nanjing; Sheng Huaibin, Beijing; Claude Spinosa, Boise; W. W. Nassichuk, Calgary; W. B. Saunders, Boston; W. L. Manger, Fayetteville; L. K. Meeks, Warrensburg; S. V. Nikolaeva, London and Moscow, the Palaeontographical Society, London; J. D. Price, formerly University of Hull; A. L. Titus, Kanab; H. W. J. van Amerom, Heerlen; W. H. C. Ramsbottom, Sheffield; and D. M. Work, Augusta.

## **REPOSITORIES AND THEIR ABBREVIATIONS**

Abbreviations and locations of museums and institutions holding type material, which are used throughout the systematic sections of this volume, are listed below.

- AM: Australian Museum, Sydney, Australia
- AMF: Australian Museum, Sydney, Australia
- AMNH: American Museum of Natural History, New York, USA
- ANU: Australian National University, Canberra, Australia
- AU: Geology Department, Auckland University, Auckland, Australia
- BAU: Buenos Aires University, Buenos Aires, Argentina
- BGS, GSM, IGS: British Geological Survey (formerly Geological Survey Museum; Institute of Geological Sciences, London) Keyworth, Nottinghamshire, United Kingdom
- BMNH: The Natural History Museum, London, United Kingdom [formerly British Museum (Natural History)]
- BMR: see CPC
- Br: see TAGI Br
- BSM: Bavarian State Museum, Munich, Germany
- BU: Department of Geology, Birmingham University, Birmingham, United Kingdom
- BUM: Bristol University Museum, Bristol, United Kingdom
- CAGS: Institute of Geology, Chinese Academy of Geological Sciences, Beijing, China
- CAS: California Academy of Sciences, Types Collections, San Francisco, California, USA
- CB: Muséum d'Histoire Naturelle, Geneva, Switzerland
- CEGH: see CORD-PZ
- CFP UA: Compagnie Française Petroles, Paris, France
- CGS: Czech Geological Survey, Prague, Czech Republic
- CIGMR: Chengdu Institute of Geology and Mineral Resources, Chengdu, China
- CMB: City Museum and Art Gallery, Bristol, UK
- CMNH: Carnegie Museum, Pittsburgh, USA
- **CNIGR:** Central Scientific Geological Exploration Museum (Tschernyshev Museum), St. Petersburg, Russia
- CORD-PZ: Universidad Nacional de Córdoba, Argentina
- CPC: Commonwealth Palaeontological Collections, Australian Geological Survey Organisation, Canberra, Australia
- CRMGE: Central Research Museum of Geological Explorations, St. Petersburg, Russia
- D, EM, ENSM, FSI, FSL, SSL, TA: Université Claude Bernard, Lyon I, Villeurbanne, France
- DNGM: Servicio Nacional Minero Geológico, Buenos Aires, Argentina
- DP, DPO: Departamento de Geología, Oviedo University, Oviedo, Spain

DPO: see DP

- DPUCM: Departamento de Paleontologia, Universidad Complutense, Madrid, Spain
- EM: see D
- ENSM: see D
- FD: Geological College of Eastern China, Fuzhou, China
- FSI: see D
- FSL: see D
- GB: Xian Institute of Geology and Mineral Resources, Xian, China
- GBA: Geologisches Bundesanstalt Museum, Vienna, Austria
- GIB: Geological Institute, Bonn, Germany
- GIBAS: Geological Institute, Bulgarian Academy of Sciences, Sofia, Bulgaria
- GIN KAZ: Institute of Geology, Kazakh Academy of Sciences, Alma-Ata, Kazakhstan
- GIN TAD: Institute of Geology, Dushanbe, Tadzhikistan
- GIN UZ: Institute of Geology, Uzbek Academy of Sciences, Tashkent, Uzbekistan
- GIUS: Department of Earth Sciences, Silesian University, Sosnowiec, Poland
- GLAHM: Hunterian Museum, Glasgow University, Scotland, United Kingdom
- GMC, IV: Geological Museum of China, Beijing, China
- GMG: State Museum of Georgia (named after S. N. Djanashia), Academy of Sciences of the Georgian SSR, Tbilisi
- GMUT: see TUG
- GM YaRGTS: Geological Museum of the Regional Geological Centre, Yakutsk, Yakutia
- GPIBo: Palaontological Institute, Bonn, Germany
- GPIT: Institut für Geowissenschaften, Tübingen, Germany
- GPZ: Department of Geology and Palaeontology, Zagreb, Croatia
- GSC: Geological Survey of Canada, Ottawa, Ontario, Canada
- GSE: see IGS GSE
- GSI: Geological Survey of India, Calcutta, India GSM: see BGS
- GSQ: Geological Survey, Queensland, Australia
- GSV: Geological Survey of Victoria, Australia
- GSWA: Geological Survey of Western Australia, Perth, Australia
- GS YA: see CGS
- HB: Bureau of Geology and Mineral Resources of Hunan Province, Hunan, China
- HGI: Hungarian Geological Institut, Budapest, Hun-
- HIGS: Hangzhou Institute for Geological Science, Hangzhou, China
- HM: see GLAHM

**HNHMB:** Hungarian Natural History Museum, Budapest, Hungary

- I: New York State Geological Survey, Albany, New York, USA
- ICPSB: Institute of Geology, University of Padua, Italy
- IG: Palaeontological Collections of L'Institut Royal des Sciences Naturelles de Belgique, Brussels
- IGAS: Institute of Geology, Chinese Academy of Sciences, Beijing, China
- IGiG: Institute of Geology and Geophysics, Siberian Branch, Academy of Sciences, Akademgorodok, Russia
- **IGM:** Instituto de Geología, Universidad Autónoma de México, Ciudad Univesitaria, México City, Mexico
- IGN: Institute of Geological Sciences, Kiev, Ukraine
- **IGN SO RAN:** Geological Museum of the Institute of Geological Sciences of Yakutia Sakha Scientific Centre, Siberian Division, Russian Academy of Sciences, Yakutsk, Yakutia
- IGNA: Geological Museum of the Institute of Geological Sciences, Almaty, Kazakhstan
- IGR: Institute of Geology, University of Rennes, Rennes, France
- IGS GSE: Institute of Geological Sciences, Edinburgh, United Kingdom
- IGS GSM: see BGS
- IMGPT: Geological-Paleontological Institute and Museum of Tübingen University, Germany
- Inst. Geol.: Geological Institute, Bishkek, Kyrgyzstan
- IO: P. P. Shirshov Institute of Oceanology, Moscow, Russia
- IV: see GMC
- IPW: Institut für Paläontologie der Universität (Geozentrum), Vienna, Austria
- **IRScNB:** Palaeontological Collections of L'Institut Royal des Sciences Naturelles de Belgique, Brussels
- JCF: James Cook University, Townsville, Queensland, Australia
- KAS, MANK: Geological Museum of Institute of Geological Sciences, Almaty, Kazakhstan
- KHGU: Kharkov State University, Ukraine
- **KIGLGU:** Geology Faculty of Leningrad State University, Paleontology-Stratigraphy Museum, St. Petersburg, Russia
- L: National Museum, Prague, Czech Republic, Barrande specimens
- LGE: St. Petersburg State University, St. Petersburg, Russia
- LGI: Leningrad Geological Institute, Leningrad, Russia
- LM: see LO
- LMT: Loodus Museum, Tallinn, Estonia
- LO (formerly LM): Lund University Museum, Sweden
- **LPB:** Laboratoire de Paléontologie, Université de Bretagne Occidentale, Brest, France
- LS: Linnean Society of London, United Kingdom
- MANK: see KAS

- **MB (formerly HUB):** Humboldt University, Berlin, Germany
- M.Ch: Museum Chabarovsk, Verkhoyan, eastern Siberia, Russia
- **MBHR:** Museum of Dr. B. Horák, Rokycany, Czech Republic
- MCMB: Department of Geology, University of Beijing, Beijing, China
- MCZ: Museum of Comparative Zoology, Harvard University, Cambridge, Massachusetts, USA
- MDSGF: Museo del Dipartimento di Scienze Geolgiche dell'Università di Ferrara, Ferrara, Italy
- MDSGF: Museo del Dipartimento di Scienze Geolgiche dell'Università di Ferrara, Ferrara, Italy
- MDTF: see MDSGF
- MFLV: Museo dei Fossili della Lessinia, Verona, Italy
- MFMGB: Museum of the Faculty of Mining and Geology, Belgrade University, Belgrade, Yugoslavia
- MG: Institute of Geology, Ashkhabad, Turkmenistan
- MGBW: Museum of the Geologische Bundesanstalt of Wien, Austria
- MGRI: Moscow Geological Prospecting Institute, Moscow, Russia
- **MGSB:** Museo Geológico del Seminario de Barcelona, Barcelona, Spain
- MGU: Moscow State University, Russia
- MGUH: Geological Museum, Copenhagen, Denmark
- MGUP: Museum of Geology, University of Palermo, Sicily, Italy
- MIP: see MLP
- **MLP:** Invertebrate Paleontology Department, La Plata Natural Sciences Museum, La Plata, Argentina
- MM: Geological Survey, Prague, Czech Republic
- MM: Moravian Museum, Brno, Czech Republic
- **MMF:** Geological and Mining Museum, Department of Mines, Sydney, Australia
- MNB: see MB
- **MNHN:** Muséum National d'Histoire Naturelle, Paris, France
- MONZ: see NMNZ
- MPL: see MLP
- **MPM:** Milwaukee Public Museum, Milwaukee, Wisconsin, USA
- **MPUM:** Museo di Paleontologia del Dipartimento di Scienze della Terra dell'Università degli Studi di Milano, Italy
- MUGT: see GIN TAD
- **Muz IG:** Geological Museum of the Geological Institute, Warsaw, Poland
- MV: see NMVP
- NHM: Natural History Museum, London, UK
- NHMB: Natural History Museum, Basel, Switzerland (Naturhistorisches Museum Basel)
- NHMW: Natural History Museum in Vienna, Naturhistorisches Museum, Wien, Austria
- NIGP: Nanjing Institute of Geology and Palaeontology, Academia Sinica, Nanjing, China
- NM: National Museum, Prague, Czech Republic
- NMING: National Museum of Ireland, Dublin, Ireland
- NMNZ: Te Papa, Museum of New Zealand, Wellington, New Zealand

xxvii

HUB: see MB

- NMVP: Victoria Museum, Melbourne, Victoria, Australia
- **NMW:** National Museum of Wales, Cardiff, United Kingdom
- **NMV P:** Department of Invertebrate Palaeontology, Museum of Victoria, Australia
- NS: Northeastern Institute of Geology, Inner Mongolia
- NUF: Department of Geology, University of Newcastle, New South Wales, Australia
- NYSM: New York State Museum, Albany, USA
- NZGS: New Zealand Geological Survey, Lower Hutt, New Zealand (presently called Institute of Geological and Nuclear Sciences)
- NZOI: New Zealand Oceanographic Institute, National Institute of Water and Atmospheric Research, Wellington, New Zealand
- OKGS: Oklahoma Geological Survey, Norman, Oklahoma, USA
- OMR: District Museum, Rokycany, Czech Republic
- OMR VH: see OMR
- **OSU:** Orton Geological Museum, Ohio State University, Columbus, Ohio, USA
- OU: University of Oklahoma, Norman, USA
- OUM: Oxford University Museum, United Kingdom
- **OU NZ:** Geology Department, Otago University, Dunedin, New Zealand

PAN: see PIN

- PCZCU: Department of Biology, Západočeská univerzita, Plzeň, Czech Republic
- **PIN:** Palaeontological Institute, Russian Academy of Sciences, Moscow, Russia
- PIN RAS: see PIN
- **PIW:** Paleontological Institute, Würzburg University, Würzburg, Germany
- **PKUM:** Geological Museum of Beijing University, China
- **PM (formerly PMU):** Palaeontological Museum, Uppsala University, Uppsala, Sweden
- **PMNUF:** Paleontological Museum, University of Naples 'Federico II', Naples, Italy
- **PMO:** Paleontologisk Museum, University of Oslo, Norway
- PMU: see PM
- **PRI:** Paleontological Research Institute, Ithaca, New York, USA
- PUC: Princeton University Collections, Princeton, New Jersey, USA
- PUM: Geology, Peking University, Beijing, China
- **QMF:** Queensland Museum, South Brisbane, Australia
- RCCBYU: Research Center for the Chengjiang Biota, Yunnan University, Yunnan, China
- RGF VR: Institute of Regional Geology and Paleontology, Faculty of Mining and Geology, University of Belgrade, Belgrade, Serbia RM, RMS: Swedish Museum of Natural History, Stockholm, Sweden
- ROM: Royal Ontario Museum, Toronto, Ontario, Canada
- **RX:** Rowley Collection, University of Illinois, Urbana, Illinois, USA

- SAM.P: South Australian Museum, Adelaide, South Australia
- SBNML: National Museum, Prague, Czech Republic
- SGU: Geological Survey of Sweden, Uppsala, Sweden
- SIGM: Shenyang Institute of Geology and Mineral Resources, Shenyang, Liaoning, China
- SM (formerly SMA): Sedgwick Museum, University of Cambridge, United Kingdom
- SMF: Senckenbergische Museum, Frankfurt, Germany
- SNM: Slovakian National Museum, Bratislava, Slovakia (Slovenské Narodné Múzeum, Bratislava)
- SSL: see D
- **SUI:** University of Iowa, Department of Geology, Iowa City, USA
- SUP: Palaeontological collections, University of Sydney, New South Wales, Australia
- T: Paleontological Museum, University of Naples, Naples, Italy
- TA: see D
- **TAGI BR:** Geological Museum, Institute of Geology, Tallinn Technical University, Tallinn, Estonia
- TBR: see TF
- TF: Geological Survey Division, Department of Mineral Resources, Bangkok, Thailand
- TsGM: see CNIGR
- TsNIGRA: see CNIGR
- TUBr: Paläontologische Sammlung, Institut für Geowissenschaften, Universität Tübingen, Germany
- **TUG:** Museum of Geology, University of Tartu, Tartu, Estonia
- **UA:** Geology Department, University of Alberta, Edmonton, Canada
- UC: Field Museum of Natural History, Chicago, Illinois, USA
- UCF: The University, Calgary, Canada
- UCLA: University of California at Los Angeles, Los Angeles, California,
- USA
- UCM: University of Canterbury, Christchurch, New Zealand
- UCMP: University of California, Museum of Paleontology, USA
- UD: University of Dijon, Dijon, France
- UHR: Hokkaido University, Sapporo, Japan
- UI: University of Illinois, Urbana, Illinois, USA
- UL: Department of Geology and Palaeontology, University of Ljubljana, Slovenia
- UM: Museum of Paleontology, University of Michigan, Ann Arbor, Michigan, USA
- **UMC (formerly UMO):** University of Missouri, Columbia, Missouri, USA
- **UMMF:** Department of Geology, University of Montpellier, Montpellier, France
- **UMUT:** University Museum of the University of Tokyo, Tokyo, Japan
- UND: University of Notre Dame, Indiana, USA
- U.N.E: University of New England, Armidale, Australia
- UPS: Université de Paris-Sud, France

© 2009 University of Kansas Paleontological Institute

- **UQF:** University of Queensland, Department of Geology, Brisbane, Australia
- USNM: United States National Museum, Washington, D.C., USA
- UT: Department of Geology, University of Texas, Austin, Texas, USA
- UTC: Department of Geology, University of Toronto, Toronto, Canada
- **UTGD:** University of Tasmania Geology Department, Hobart, Tasmania, Australia
- U.W.A.: University of Western Australia, Nedlands, Western Australia

VH: see OMR

- VSEGEI: Russian Geology Institute, St. Petersburg, Russia
- XAGM: Xi'an Institute of Geology and Mineral Resources, Shaanxi, China

- **XB:** Palaeontological Collections of the Xi'an Institute of Geology and Mineral Resources, Chinese Academy of Geological Sciences, Xi'an, Shaanxi Province, China
- **XIGMR:** Xi'an Institute of Geology and Mineral Resources, Shaanxi, China
- YaTGU: Geological Museum, Yakutsk, Yakutia
- YIGM: Yichang Institute of Geology and Mineral Resources, Yichang, China
- **YPM:** Yale University, Peabody Museum of Natural History, New Haven, Connecticut, USA
- **ZI:** Zhejiang Institute of Geology and Mineralogy, Zhejiang, China
- ZPAL Br: Institute of Palaeobiology, Polish Academy of Sciences, Warsaw, Poland