

FIG. 129. Permobiellidae, Amboneuridae, and Pleisiogrammatidae (p. 202–204).

SC extending well beyond midwing; 3A vestigial or absent. *Perm.*

Pleisiogramma CARPENTER, 1943a, p. 73 [**P. mediale*; OD]. M unbranched. *Perm.*, USA (Kansas). —FIG. 129,2. **P. mediale*; wing, $\times 3.2$ (Carpenter, 1943a).

Family SYNOMALOPTILIDAE
Martynov, 1938

[Synomaloptilidae MARTYNOV, 1938a, p. 76]

Related to Euthygrammatidae, but veins CUA and CUP anastomosed for a considerable distance. *Perm.*

Synomaloptila MARTYNOV, 1938a, p. 76 [**S. longipennis*; OD]. SC terminating well before wing

apex; separation of CUA and CUP at about level of midwing. *Perm.*, USSR (Asian RSFSR). —FIG. 128,1. **S. longipennis*; wing, $\times 2$ (Martynov, 1938a).

Family UNCERTAIN

The following genera, apparently belonging to the order Caloneuroidea, are too poorly known to permit assignment to families.

Caloneurella CARPENTER, 1934, p. 324 [**C. carbonaria*; OD]. Apical wing fragment, probably related to *Caloneura*. *U. Carb.*, USA (Pennsylvania). —FIG. 128,2. **C. carbonaria*; $\times 1.8$ (Carpenter, 1934).

Pruvostiella HANDLIRSCH, 1922, p. 82 [**Euthyneura lecomtei* PRUVOST, 1919, p. 115; OD]. Small wing fragment. *U. Carb.*, Europe (France).

Order MIOMOPTERA
Martynov, 1927

[Miomoptera MARTYNOV, 1927d, p. 101, *emend.* MARTYNOV, 1938b, p. 138]

Small to very small insects, with wings nearly homonomous. Fore wing with vein SC ending before or at midwing; R commonly with a distal twig; RS arising before midwing, with at least 3 terminal branches; M commonly coalesced with CUA basally to varying amounts, but diverging in basal third of wing; M deeply forked; CUA with 2 or 3 terminal branches; CUP unbranched; 2 anal veins typically present. Hind wing similar in form to fore wing, without an anal lobe or fan; M usually arising from CU very near wing base. Body structure little known; head of moderate size; mouthparts apparently mandibulate; antennae conspicuous, relatively thick, with 15 to 20 segments; tarsi with 4 segments (*Palaeomantis*); cerci short. Immature stages unknown. *U. Carb.*—*Perm.*

The status of this order is uncertain. As originally proposed by MARTYNOV (1927d) it included five Permian families, previously placed in the order Protorthoptera, but it was based mainly on one of them, the Palaeomanteidae (=Delopteridae). The following year TILLYARD (1928b), obviously unaware of MARTYNOV's article, proposed the new order Protoperlaria for the same series of families except the Palaeomanteidae. Subsequent studies of extensive collections of

Palaeomanteidae and Lemmatophoridae (CARPENTER, 1933a, 1935a; MARTYNOV, 1938b; MARTYNOVA, 1958, 1961b, 1962a) supported the view that the Palaeomanteidae were sufficiently distinctive to justify ordinal separation and that the Lemmatophoridae and related families were in reality part of the order Protorthoptera.

The Miomoptera, as exemplified by the Palaeomanteidae, stand apart from the protorthopterous families, with which they have been associated, by two distinct features. One of these is the absence of the anal lobe or fan on the hind wing, probably a secondary condition, as in the Isoptera and Embioptera. The other is the very small number of crossveins and their virtual absence from the costal area of both wings.

Several families have been added to the Miomoptera since MARTYNOV's original publication on the order. Two of these, Archaeomipteridae (GUTHÖRL, 1939) and Metroptoridae HANDLIRSCH (1906a) almost certainly belong in the order (MARTYNOVA, 1958, 1961b; CARPENTER, 1965). The evidence for the others, however, is very weak and in my opinion insufficient to justify their inclusion in the Miomoptera. The family Permembiiidae TILLYARD (1937b), originally described in the Psocoptera, has been transferred to the Miomoptera by KUKALOVÁ (1963a) and RIEK (1973, 1976a); *Permembia* itself is known from a very few, poorly preserved specimens (CARPENTER, 1976), with a venation that has little in common with that of the Palaeomanteidae. Some details of body structure are preserved in two specimens, but since almost nothing is known of the body of the Palaeomanteidae we have no basis for comparing those details. The family Permosialidae MARTYNOV (1928b), originally in the Neuroptera, has been placed in the Miomoptera by RIEK (1976a) and RASNITSYN (1977c), as have the families Permonkidae RASNITSYN (1977c) and Palaeomantiscidae RASNITSYN (1977c). These families, however, are characterized by broadened anal areas or anal lobes on the hind wings and numerous crossveins on both wings including the costal areas. Since no revisions

of the definition or diagnosis of the Miomoptera have been proposed for the accommodation of these families, the order is treated here essentially as it was defined by MARTYNOV (1938b) and MARTYNOVA (1961b, 1962a). The families Permembiiidae, Permosialidae, Permonkidae, and Palaeomantiscidae, along with a nymphal form, *Permonympha* SHAROV (1957b), are included under Neoptera, Order Uncertain.

The homologies of the wings of the Miomoptera have not been definitely determined. In the fore wing, veins R and CUA are clearly convex, but RS and M show no definite topography. Whether the branches of M represent MA and MP (KUKALOVÁ, 1963a) or only one of these veins is uncertain; they are designated here as M1+2 and M3+4. The amount of anastomosis of M with CUA in the fore wing varies from genus to genus; in some species (e.g., *Permodelopterum obscurum* KUKALOVÁ; see Fig. 131,4a), M appears to arise independently of CUA, which joins it later; in others (e.g., *Palaeomantis minutum* (SELLARDS); see Fig. 131,1b) the stem of M seems to be coalesced with that of CUA from the very base of the wing. In all species, however, M diverges from CUA before midwing. In some individual wings, M1+2 seems to arise from RS or RS3+4; this may be a specific or even a generic characteristic, and there is some evidence that it occurs as an individual variation (CARPENTER, 1939).

The Miomoptera are among the smallest insects known from the Upper Carboniferous and Permian. However, their affinities are not clear. They are generally considered to have been related to the Psocoptera, although RASNITSYN (1980b) believes them to have been endopterygote insects, close to the ancestral stock of the Hymenoptera. MARTYNOV concluded (1938b) that they were an early, aberrant branch of protorthopterous or perlarian stock.

Family METROPATORIDAE Handlirsch, 1906

[Metroptoridae HANDLIRSCH, 1906a, p. 681]

Hind wing nearly oval; vein SC short, weakly developed, and close to R; RS forked




FIG. 130. Metropatoridae and Archaemiopteri-
dae (p. 206).

before midwing; RS1+2 with 4 terminal branches; RS3+4 forked once; M apparently arising from CUA at base of wing and forked almost to level of origin of RS; CUA with short terminal fork. Fore wing and body unknown. *U. Carb.*

Metropator HANDLIRSCH, 1906a, p. 681 [**M. pusillus*; OD]. RS3+4 more deeply forked than RS1+2; M1+2 forked distally; M3+4 forked twice. TILLYARD, 1926c; CARPENTER, 1965. *U. Carb.*, USA (Pennsylvania).—FIG. 130,2. **M. pusillus*; hind wing, $\times 6.5$ (Carpenter, 1965).

Family ARCHAEMIOPTERIDAE Guthörl, 1939

[Archaemiopteridae GUTHÖRL, 1939, p. 320]

Fore and hind wings broadly oval; vein RS1+2 forked, at least distally. Body unknown. *U. Carb.*—*Perm.*

Archaemioptera GUTHÖRL, 1939, p. 320 [**A. carbonaria*; OD]. RS3+4 unbranched; M forked to more than half its length; CUA with distal fork only. *U. Carb.*, Europe (Germany).—FIG. 130,1. **A. carbonaria*; fore(?) wing, $\times 10$ (Carpenter, new, based on holotype).

Eodelopterum SCHMIDT, 1962, p. 838 [**E. priscum*; OD]. Similar to *Tychtodelopterum*, but CUA with 3 terminal branches; M1+2 terminating at wing apex. GUTHÖRL, 1963. *U. Carb.*, Europe (Germany).—FIG. 130,4. **E. priscum*; hind wing, $\times 12$ (Guthörl, 1963).

Saaromioptera GUTHÖRL, 1963, p. 22 [**S. jordani*; OD]. Similar to *Tychtodelopterum*, but RS3+4 unbranched. *U. Carb.*, Europe (Germany).—FIG. 130,5. **S. jordani*; fore(?) wing, $\times 11$ (Guthörl, 1963).

Tychtodelopterum MARTYNOVA, 1958, p. 70 [**T. relictum*; OD]. Similar to *Archaemioptera*, but RS3+4 deeply forked; CUA forked to at least half its length. *Perm.*, USSR (Asian RSFSR).—FIG. 130,3. **T. relictum*; hind wing, $\times 16$ (Martynova, 1962a).

Family PALAEOMANTEIDAE Handlirsch, 1906

[*Nom. correct.* BRUES, MELANDER, & CARPENTER, 1954, p. 811, *pro* Palaeomantidae HANDLIRSCH, 1906b, p. 348] [=Delopteridae SEL-
LARDS, 1909, p. 168]

Fore wing membranous or coriaceous; vein SC usually ending before midwing, less commonly near midwing; RS usually with 3 terminal branches; basal stem of M apparently independent of both R and CU in some genera, but anastomosed with one of these in other genera; M forked deeply, usually to midwing; rarely, M1+2 apparently anastomosed with stem of RS (or connected by crossvein), resembling additional branch of RS; CUA with 2 or 3 terminal branches; distinct marginal indentation at end of CUP in some genera. Hind wing similar to fore

wing except for differences characteristic of the order. Antennae with about 15 short segments; tarsi with 4 segments; abdomen short, with wings projecting far beyond abdomen at rest; cerci very short. *U. Carb.*—*Perm.*

Palaeomantis HANDLIRSCH, 1904b, p. 4 [**P. schmidii*; OD] [= *Delopterum* SELLARDS, 1909, p. 168 (type, *D. minutum*); *Pseudodelopterum* MARTYNOV, 1928b, p. 66 (type, *Delopterum latum* SELLARDS, 1909); *Pseudomantis* MARTYNOV, 1928b, p. 73 (type, *P. minuta*); *Leptoneurula* MARTYNOV, 1928b, p. 77 (type, *L. insignis*); *Delopsocus* TILLYARD, 1928f, p. 474 (type, *D. elongatus*); *Miomantisca* ZALESSKY, 1956a, p. 275 (type, *M. clara*); *Miomatoneurites* ZALESSKY, 1956a, p. 278 (type, *M. sylvanensis*); *Delopteriella* ZALESSKY, 1956a, p. 284 (type, *D. graciosa*); *Stefanomioptera* GUTHÖRL, 1962a, p. 67 (type, *S. hangardi*)]. Fore wing slender, membranous; SC terminating at about midwing; RS diverging from R in basal third of wing and usually with branches RS1, RS2, and RS3+4; M separating from CUA near level of origin of RS; M1+2 and M3+4 diverging near midwing; CUA forked; crossveins very few and weakly developed; posterior margin of wing either smoothly curved or with an indentation at end of CUP. Hind wing slightly shorter than fore wing; SC short, usually terminating about one-third wing length from base. Hind wing similar in form to fore wing but apparently lacking a separate and distinct CUP; CUA not preserved as a definite, convex vein as in fore wing, but forming with CUP a compound vein (CU) and preserved as a strong ridge within a furrow. Head with large compound eyes; first tarsal segment longer than others; cerci short. CARPENTER, 1933a, 1967c; GUTHÖRL, 1962a; MARTYNOVA, 1962a; KUKALOVÁ, 1963a. *U. Carb.*, Europe (Germany); *Perm.*, USA (Kansas, Oklahoma), Europe (Czechoslovakia), USSR (European RSFSR).—FIG. 131,1a. *P. hangardi* (GUTHÖRL), *U. Carb.*, Germany; fore wing, $\times 9$ (Carpenter, 1967c).—FIG. 131,1b,c. **P. minutum* (SELLARDS), *Perm.*, Kansas; b, fore and c, hind wings, $\times 12$ (Carpenter, 1933a).

Miomatoneura MARTYNOV, 1927d, p. 106 [**M. frigida*; OD]. Fore wing as in *Palaeomantis*, but M1+2 arising from stem of RS or connected to it by crossvein; CUA with 2 or 3 terminal branches. Hind wing unknown. MARTYNOVA, 1961b, 1962a; KUKALOVÁ, 1963a. *Perm.*, USSR (European RSFSR), Europe (Czechoslovakia).—FIG. 131,2a. **M. frigida*, *Perm.*, USSR; fore wing, $\times 8$ (Martynova, 1962a).—FIG. 131,2b. *M. candida* KUKALOVÁ, *Perm.*, Czechoslovakia; fore wing, $\times 14$ (Kukalová, 1963a).

Miomatoneurella MARTYNOVA, 1958, p. 71 [**M. reducta*; OD]. Fore wing similar to that of *Miomatoneura*, but RS1+2 unbranched. MARTYNOVA, 1961b. *Perm.*, USSR (Asian RSFSR).—FIG.

131,3. **M. reducta*; fore wing, $\times 15$ (Martynova, 1962a).

Permodelopterum KUKALOVÁ, 1963a, p. 25 [**P. obscurum*; OD]. Fore wing similar to that of *Perunopterum* but broader and base of M apparently coalesced with R basally. Hind wing and body unknown. *Perm.*, Europe (Czechoslovakia).—FIG. 131,4a. **P. obscurum*; fore wing, $\times 8$ (Kukalová, 1963a).—FIG. 131,4b. *P. lumbiforme* KUKALOVÁ; fore wing, $\times 12$ (Kukalová, 1963a).

Perunopterum KUKALOVÁ, 1963a, p. 16 [**P. peruni*; OD]. Fore wing membranous or distinctly coriaceous, densely covered with minute hairs, and more slender than in *Palaeomantis*; indentation of hind margin at end of CUP pronounced; SC terminating near midwing; R with distal twig; RS arising before midwing, typically with branches RS1, RS2, and RS3+4; stem of M free from R basally; M arising from CUA at about level of origin of RS, with 2 long branches; CUA forked. Hind wing similar to fore but with SC shorter and weaker and costal area narrower; M separating from CUA nearer wing base. Both wings tend to show more crossveins than in *Palaeomantis*, even having reticulation in anal area. Body structure apparently much as in *Palaeomantis*; short cerci present. *Perm.*, Europe (Czechoslovakia).—FIG. 131,5a. **P. peruni*; fore wing, $\times 12$ (Kukalová, 1963a).—FIG. 131,5b. *P.(?) corium* KUKALOVÁ; fore wing, $\times 8$ (Kukalová, 1963a).

Order THYSANOPTERA Haliday, 1836

[Thysanoptera HALIDAY, 1836, p. 439]

Small or minute insects, with slender body (Fig. 132); head usually quadrangular; compound eyes small but prominent, with relatively large, rounded facets; ocelli commonly present; antennae with 6 to 10 segments; labrum and labium forming a short cone, containing as stylets the left mandible (right one absent or vestigial) and extensions of the 2 maxillae; maxillary and labial palpi present; prothorax free from mesothorax and well developed; wings usually present, nearly homonomous, membranous but very narrow, often strap-shaped, with not more than 2 longitudinal veins; wings fringed with long setae, at least along posterior margins; both brachypterous and apterous individuals may occur in some species; legs short, with 1 to 2 tarsal segments; abdomen elongate, seg-




FIG. 131. Palaeomanteidae (p. 206–207).

mentation distinct; ovipositor present (Terebrantia) or absent (Tubulifera); cerci absent. *Perm.—Holo.*

The Thysanoptera, or thrips, feed by lacerating the surface of plants with their stylets and drawing the plant juices through the mouth cone. Some species are predaceous on small arthropods. The nymphs hatch from eggs laid on or in plant tissue; they resemble the adults in general form and feeding habits. Nymphs of most species pass through two or three quiescent stages (prepupa and pupa) in which wing pads occur.

The thrips, like the Psocoptera, are a dis-

trictive and homogeneous order of insects. Two suborders, Terebrantia and Tubulifera, are generally recognized; the most obvious differences between them are in the presence or absence of the ovipositor in the female, and in the shape of the terminal abdominal segment. However, other differences, of more phylogenetic significance, are found in the detailed structure of the mouthparts and wings. Such morphological evidence indicates that the Terebrantia are more primitive than the Tubulifera and that the family Aeolothripidae of the Terebrantia is the most primitive of the recent families.

Thysanoptera are well represented in the Tertiary deposits, mainly the Baltic amber; these Oligocene species have been studied in detail by two authorities on recent thrips, BAGNALL and PRIESNER. Two pre-Tertiary thrips have been described, one (*Liassothrips*) from the Jurassic (MARTYNOV, 1927b) and the other (*Permothrips*) from the Permian (MARTYNOV, 1935a). That these are thrips seems almost certain, although their subordinal positions are obscure. The fossil record of the Thysanoptera shows little to date about the evolution of the order. The Permian species appear to have had somewhat larger wings than any existing species, but no veins are preserved and the structure of the mouthparts is unknown. The classification used here is that of PRIESNER (1949).

Suborder TEREBRANTIA

Haliday, 1836

[Terebrantia HALIDAY, 1836, p. 439]

Terminal abdominal segment conical (rarely tubular) in females, bluntly rounded in male; female with sawlike ovipositor; fore wings nearly always with at least 1 longitudinal vein (in addition to costal vein) extending to apex. *Perm.—Holo.*

Family PERMOTHRIPIDAE

Martynov, 1935

[Permothripidae MARTYNOV, 1935a, p. 334]

Head somewhat extended; pronotum transverse; legs short, tibiae more slender than femora; wings broad and long, extending beyond abdomen; ovipositor apparently present. [Subordinal position of the family uncertain; it probably represents an extinct suborder.] *Perm.*

Permothrips MARTYNOV, 1935a, p. 334 [**P. longipennis*; OD]. Abdomen narrowed distally, not tubular. *Perm.*, USSR (Asian RSFSR).—FIG. 133, 5. **P. longipennis*; general form of body and wings, $\times 16$ (Martynov, 1935a).

Family AEOLOTHRIPIDAE Uzel, 1895

[Aeolothripidae UZEL, 1895, p. 42]

Wings broad and rounded at apex; ovipositor curved upward; antennae with 9 segments. *Eoc.—Holo.*




FIG. 132. Thysanoptera; dorsal view of whole insect, *Desmothrips propinquus*, Holocene, $\times 20$ (CSIRO, 1970).

Aeolothrips HALIDAY, 1836, p. 451. COCKERELL, 1917b; BAGNALL, 1924a; PRIESNER & QUIÉVREUX, 1935. *Oligo.*, England, Europe (France, Germany)—*Holo.*

Archanothrips PRIESNER, 1924, p. 132 [**A. pugionifer*; OD]. Similar to *Ankothrips* (recent), but hind angles of prothorax with short bristles; ninth antennal segment with 4 pale, transverse sutures. PRIESNER, 1949. *Oligo.*, Europe (Baltic).

Eocranothrips BAGNALL, 1926, p. 17 [**Melanothrips annulicornis* BAGNALL, 1923, p. 36; OD]. Similar to *Cranothrips* (recent), but all antennal segments simple, without projections. PRIESNER, 1949. *Oligo.*, Europe (Baltic).

Lithadothrips SCUDDER, 1875b, p. 221 [**L. vetusta*; OD]. Similar to *Orothrips* (recent), but fore wings widened toward apex. BAGNALL, 1924a; PRIESNER, 1949. *Eoc.*, USA (Utah); *Oligo.*, Europe (Germany).

Melanthrips HALIDAY, 1836, p. 450. SCUDDER, 1890. ?*Eoc.*, USA (Colorado)—*Holo.*

Palaeothrips SCUDDER, 1875b, p. 222 [**P. fossilis*; OD]. Apparently related to *Rhipidothrips* (recent); antennae with 7 segments, apical segments not conical. PRIESNER, 1949. *Eoc.*, USA (Utah).

Promelanthrips PRIESNER, 1930, p. 113 [**P. spiniger*; OD]. Similar to *Ankothrips* (recent), but hind angles of prothorax with one long bristle only. USINGER, 1942. *Oligo.*, Europe (Baltic).

Rhipidothripoides BAGNALL, 1923, p. 36 [**R. abdominalis*; OD]. Similar to *Rhipidothrips* (recent), but ninth segment of abdomen unusually elongate and third, fourth, and fifth antennal segments of about equal length. *Oligo.*, Europe (Baltic).

Stenurothrips BAGNALL, 1914, p. 483 [**S. succineus*; OD]. Ovipositor straight or nearly so; terminal abdominal segment tubular; setae of hind margin of fore wing wavy. [Family position




FIG. 133. Permothripidae, Aeolothripidae, Thripidae, Uncertain, and Liassothripidae (p. 209–212).

uncertain; possibly belonging to Heterothripidae.] BAGNALL, 1923; PRIESNER, 1949; STANNARD, 1956. *Oligo.*, Europe (Baltic).—FIG. 133,2. *S. bagnalli* STANNARD; *a*, terminal abdominal segment, $\times 66$; *b*, fore wing, $\times 55$ (Stannard, 1956).

Family THIRIPIDAE Stephens, 1829

[Thripidae STEPHENS, 1829b, p. 363]

Similar to Heterothripidae, but antennae with 6 to 9 segments and with slender sense cones; cones simple or forked; tarsal claws, if present, at apex of first or second tarsal segment. *Oligo.*—*Holo.*

Thrips LINNÉ, 1761, p. 266. *Holo.*

Amorphothrips BAGNALL, 1924c, p. 252 [**A. klebsi*; OD]. Similar to *Procerothrips*, but eye occupying whole side of head; pronotum transverse, without setae; hind legs very long and stout. BAGNALL, 1923; PRIESNER, 1949. *Oligo.*, Europe (Baltic).

Anaphothrips UZEL, 1895, p. 142. PRIESNER, 1930; PALMER, 1957. *Oligo.*, Europe (Baltic); *Mio.*, USA (California)—*Holo.*

Frankliniella KARNY, 1910, p. 46. PRIESNER & QUIÉVREUX, 1935. *Oligo.*, Europe (France)—*Holo.*
Geronthrips PRIESNER, 1949, p. 41, *nom. subst. pro Archaeothrips* PRIESNER, 1924, p. 138, *non* FIELD, 1910 [**Archaeothrips latipennis* PRIESNER, 1924, p. 138; OD]. Wings very broad, entire surface reticulate. PRIESNER, 1930. *Oligo.*, Europe (Baltic).—FIG. 133,4. **G. latipennis* (PRIESNER); fore wing, $\times 66$ (Priesner, 1924).

Heliothrips HALIDAY, 1836, p. 443. BAGNALL, 1924b. *Oligo.*, Europe (Baltic)—*Holo.*

Hercinothrips BAGNALL, 1932, p. 506. STANNARD, 1956. *Oligo.*, Europe (Baltic)—*Holo.*

Homothrips BAGNALL, 1915, p. 588. BAGNALL, 1924b. *Oligo.*, Europe (Baltic)—*Holo.*

Lipsanothrips PRIESNER, 1930, p. 119 [**L. skewarvae*; OD]. Antennae with 8 segments, sixth segment much shorter than fifth; wing with 1 or 2 longitudinal veins. *Oligo.*, Europe (Baltic).

Oxythrips UZEL, 1895, p. 141. PRIESNER, 1924, 1930. *Oligo.*, Europe (Baltic)—*Holo.*

Praedendrothrips PRIESNER, 1924, p. 139 [**P. avus*; OD]. Antennae with 9 segments, last 4 clearly separate from one another; posterior angles of

pronotum with at least 1 conspicuous bristle. PRIESNER, 1930. *Oligo.*, Europe (Baltic).

Procerothrips BAGNALL, 1924c, p. 252 [**P. cylindricornis*; OD]. Antennae with 8 segments and style with 2 segments; third through sixth antennal segments with parallel sides and of same width. BAGNALL, 1923. *Oligo.*, Europe (Baltic).

Selenothrips KARNY, 1911, p. 180. BAGNALL, 1923. *Oligo.*, Europe (Baltic)—*Holo.*

Taeniothrips SERVILLE, 1843, p. 644. BAGNALL, 1924b; PRIESNER, 1930, 1949. *Oligo.*, Europe (Baltic)—*Holo.*

Telothrips PRIESNER, 1930, p. 116 [**T. klebsi*; OD]. Similar to *Praedendrotbrips* but with sixth antennal segment large and stout and seventh, eighth, and ninth segments minute. PRIESNER, 1949. *Oligo.*, Europe (Baltic).

Family HETEROTHRIPIDAE

Bagnall, 1912

[Heterothripidae BAGNALL, 1912, p. 222] [=Hemithripidae BAGNALL, 1923, p. 37; Stenurothripidae BAGNALL, 1923, p. 37; Opadothripidae BAGNALL, 1927, p. 562]

Wings narrow, usually pointed distally; ovipositor curved downward; antennae with 9 or 10 segments; fore tarsi usually with clawlike appendage at base of second segment. *Oligo.*—*Holo.*

Heterothrips HOOD, 1908, p. 361. *Holo.*

Electrothrips BAGNALL, 1924c, p. 251 [**E. bystrix*; OD]. Cephalic, pronotal, and wing bristles abnormally long and stout; wings and legs long and slender. *Oligo.*, Europe (Baltic).

Hemithrips BAGNALL, 1923, p. 37 [**H. femoralis*; OD]. Similar to *Heterothrips* (recent), but third and fourth antennal segments cylindrical. BAGNALL, 1924a; PRIESNER, 1949. *Oligo.*, Europe (Baltic, Germany).

Opadothrips PRIESNER, 1924, p. 133 [**O. fritschianus*; OD]. Similar to *Oligothrips* (recent), but antennal segments more elongate; terminal segment slender. BAGNALL, 1924a, 1927; PRIESNER, 1949. *Oligo.*, Europe (Baltic).

Family MEROTHRIPIDAE

Hood, 1914

[Merothripidae HOOD, 1914, p. 17]

Wings narrow, pointed distally, surface smooth (not pubescent); ovipositor curved downward; pronotum with dorsal longitudinal sutures; anterior and posterior femora greatly enlarged. *Oligo.*—*Holo.*

Merothrips ZIMMERMANN, 1900, p. 12. Antennae with 8 segments. PRIESNER, 1924. *Oligo.*, Europe (Baltic)—*Holo.*

Praemerothrips PRIESNER, 1930, p. 130 [**P. boodi*; OD]. Antennae with 9 segments. PRIESNER, 1949. *Oligo.*, Europe (Baltic).

Family UNCERTAIN

The following genera, apparently belonging to the order Thysanoptera, suborder Terebrantia, are too poorly known to permit assignment to families.

Calothrips OUSTALET, 1873, p. 24 [**C. scudderii*; OD]. Little-known thysanopteron, probably belonging to the Terebrantia. *Oligo.*, Europe (France).

Eucephalothrips BAGNALL, 1924a, p. 161 [**Thrips capito* SCHLECHTENDAL, 1887, p. 579; OD] [=Protothrips PRIESNER, 1924, p. 136 (type, *P. speratus*)]. Head quadrate; wings moderately broad, apex pointed. PRIESNER, 1949. *Oligo.*, Europe (Baltic).—FIG. 133, 1. *E. speratus* (PRIESNER); *a*, head and prothorax; $\times 66$; *b*, fore wing, $\times 66$ (Priesner, 1924).

Suborder TUBULIFERA

Haliday, 1836

[Tubulifera HALIDAY, 1836, p. 459]

Terminal abdominal segments of both sexes almost always tubular; female without ovipositor; fore wing without definite costal vein and with only a vestige of another longitudinal vein, long fringe present. *Oligo.*—*Holo.*

Family PHLAEOTHRIPIDAE

Uzel, 1895

[Phlaeothripidae UZEL, 1895, p. 42]

Characteristics of suborder. *Oligo.*—*Holo.*

Phlaeothrips HALIDAY, 1836, p. 441. SCHLECHTENDAL, 1887; BAGNALL, 1924a, 1929. *Oligo.*, Europe (Baltic)—*Holo.*

Cephenothrips PRIESNER, 1930, p. 135 [**C. laticeps*; OD]. Similar to *Pygidothrips* (recent), but wing bristles short and knobbed. USINGER, 1942; PRIESNER, 1949. *Oligo.*, Europe (Baltic).

Hoplothrips AMYOT & SERVILLE, 1843, p. 640. BAGNALL, 1929; PRIESNER, 1949. *Oligo.*, Europe (Baltic)—*Holo.*

Liotrichothrips BAGNALL, 1929, p. 97 [**L. bystrix*; OD]. Head longer than pronotum, broader than long; cheeks with few prominent setae; antennae long, with third and fourth segments subequal. Similar to *Ethirothrips* (recent), but legs as in *Liothrips* (recent). PRIESNER, 1949. *Oligo.*, Europe (Baltic).

Necrothrips PRIESNER, 1924, p. 147 [**N. nanus*;

- OD]. Similar to *Austrothrips* (recent), but eyes very large, protruding, and consisting of many facets. USINGER, 1942; PRIESNER, 1949. *Oligo.*, Europe (Baltic).
- Proleuwenia** PRIESNER, 1924, p. 148 [**P. succini*; OD]. Wings reduced (female); similar to *Idiothrips* (recent), but antennae with 8 segments. USINGER, 1942; PRIESNER, 1949. *Oligo.*, Europe (Baltic).
- Schlechtendalia** BAGNALL, 1929, p. 96 [**S. longitubus*; OD]. Similar to *Pblaeothrips*, but tenth abdominal segment substantially longer than head; fifth antennal segment with a projection; wing bristles blunt. PRIESNER, 1949. *Oligo.*, Europe (Baltic).
- Symphiothrips** HOOD & WILLIAMS, 1915, p. 131. PRIESNER, 1924, 1949. *Oligo.*, Europe (Baltic)—*Holo.*
- Treherniella** WATSON, 1923, p. 81. PRIESNER, 1930, 1949. *Oligo.*, Europe (Baltic)—*Holo.*

Suborder UNCERTAIN

The genus described below, apparently belonging to the order Thysanoptera, is too poorly known to permit assignment to suborders.

Family LIASSOTHRIPIDAE Priesner, 1949

[Liassothripidae PRIESNER, 1949, p. 34] [=Mesothripidae MARTYNOV, 1927b, p. 768]

Antennae thin, with at least 7 segments; head narrow; anterior femora very broad; wings unknown. *Jur.*

Liassothrips PRIESNER, 1949, p. 34, *nom. subst. pro Mesothrips* MARTYNOV, 1927b, p. 768, *non* ZIMMERMANN, 1900 [**Mesothrips crassipes* MARTYNOV, 1927b, p. 768; OD]. Little-known thysanopteron; abdomen apparently constricted basally. *Jur.*, USSR (Kazakh).—FIG. 133,3. **L. crassipes* (MARTYNOV); body, $\times 16$ (Martynov, 1927b).

HEMIPTEROID EXOPTERYGOTES

Order HEMIPTERA Linné, 1758

[Hemiptera LINNÉ, 1758, p. 434] [=Hemipsocoptera ZALESSKY, 1937e, p. 51; Palaeohemiptera HANDLIRSCH, 1904b, p. 2]

Exopterygote Neoptera, mostly small to very small, with much morphological diversity. Head opisthognathous or prognathous; compound eyes usually present but diverse

in size; two ocelli commonly present, rarely three or none; antennae typically with five segments or less, rarely with as many as ten; mouthparts haustellate, consisting of two pairs of maxillary stylets in a segmented, rostrate labium. Pronotum of moderate size, often diversely modified; meso- and metathorax well developed. Legs usually cursorial, but forelegs of some genera raptorial, vestigial, or absent; tarsi commonly with three segments, rarely with two or one. Wings usually present, but very different in the two suborders. Wing venation quite generalized in primitive forms but much reduced in most families; fore wings of suborder Homoptera usually of uniform texture, those of suborder Heteroptera partly membranous and partly coriaceous. Abdomen well developed; ovipositor usually present. Nymphs resembling adults in basic body structure. *Perm.—Holo.*

This is the largest of the exopterygote orders, and it has apparently been a major order at least since the Triassic. All available evidence suggests that the Hemiptera are most closely related to the Psocoptera, which were well represented in the Permian. The order Hemiptera has traditionally been divided into two suborders, Homoptera and Heteroptera, the members of both groups having the same distinctive, haustellate mouthparts. Both suborders are also represented in the Permian, but the Homoptera have by far the more extensive record in that period.

The wings provide the best means of distinguishing the members of the two suborders. The homologies of the main veins are clear throughout both suborders, even in those in which the venation is much reduced. However, there has been much convergence in the reduction process. In part because of this, the family and generic classifications of the Hemiptera, especially of the Homoptera, have been based mainly on body features, such as the detailed structure of the rostrum, number and size of ocelli, tarsal segmentation, and integumentary details. Since fossils do not usually show such structures, the family position of many of the extinct genera is uncertain.




FIG. 134. Hemiptera; wing structure of the suborder Heteroptera. —1. *Nazara* sp., Pentatomidae. (adapted from CSIRO, 1970). —2. *Megacoelum* sp., Miridae (adapted from CSIRO, 1970).

The fore wings of the two suborders, except for a few primitive families in each one, differ mainly in texture. Those of the Homoptera are usually of a uniform or nearly uniform texture. Exceptions are found in a few primitive families, such as the Palaeontinidae and Prosbolidae, in which the fore wing is partly divided into two areas by an irregular nodal line (see Figs. 137,1 and 139,4a). The proximal area is usually more coriaceous than the distal part. A slight break (costal break or indentation) in the costal margin appears to be functionally associated with the nodal line. The anal area, commonly termed the clavus in the Hemiptera, is usually a distinct region of the fore wing. The fore wings of the Heteroptera, usually termed hemelytra, have a more complex structure. The distal portion of the wing is membranous, but the proximal part is coriaceous, consisting of a large, anterior, triangular section (corium) and a relatively small region (clavus), separated from the corium by the claval suture (Fig. 134,1). These areas are diverse in form and size in families and genera. The costal area of the corium may be separated from the rest of it by vein M+R, forming the embolium; and a more distal part (cuneus) of the corium may be separated by a costal fracture (Fig. 134,2). The hind wings of the Hemiptera

are generally more specialized than the fore pair, often with a very different shape. Polymorphism of wings, including aptery, occurs commonly in the order.

The great majority of the Hemiptera are phytophagous, but some Heteroptera are active predators. Immature stages have essentially the same feeding habits as the adults.

The geological record of the Hemiptera is very extensive, including almost a hundred families, two-thirds of which are Homoptera. A surprisingly large number of entomologists, specialists on the systematics of existing Hemiptera, have contributed to our knowledge of this fossil record and of the phylogeny of the order. There is, however, much difference of opinion among them about the systematic position of many of the extinct genera.

Suborder HOMOPTERA

Leach, 1815

[Homoptera LEACH, 1815, p. 124]

Fore wing of uniform texture or nearly so, not sharply differentiated into membranous and coriaceous areas; wings typically held sloping over the sides of the body at rest. *Perm.—Holo.*

Family DUNSTANIIDAE

Tillyard, 1916

[Dunstaniidae TILLYARD in TILLYARD & DUNSTAN, 1916, p. 31]

Fore wing sharply separated into tegminous basal part and membranous distal area; nodal break prominent; vein SC long, terminating on costal margin; R and RS curved; RS unbranched; clavus broad, triangular; 1A and 2A long, extending to hind margin. Hind wing little known, smaller than fore wing, with rounded anal area. Head, compound eyes, and pronotum relatively large. Relatively large insects. Affinities uncertain, but apparently closely related to the Palaeontinidae. TILLYARD, 1918d; BECKER-MIGDISOVA, 1949b; EVANS, 1956; BECKER-MIGDISOVA & WOOTTON, 1965; RIEK, 1976b. *Trias.*




FIG. 135. Dunstaniidae (p. 214).

Dunstania TILLYARD in TILLYARD & DUNSTAN, 1916, p. 31 [**D. pulchra*; OD]. M3+4 not forked, RS apparently joined to M by a short, oblique crossvein. [The genus has been reported from South Africa (RIEK, 1976b), but the generic position of the species described is very uncertain.] *Trias.*, Australia (Queensland).—FIG. 135,1. **D. pulchra*; fore wing as preserved, $\times 1.5$ (Evans, 1956).

Dunstaniodes BECKER-MIGDISOVA & WOOTTON, 1965, p. 64 [**D. elongatus*; OD]. Fore wing elongate; costal margin of basal half of wing strongly convex. *Trias.*, USSR (Kirghiz).—FIG. 135,3. **D. elongatus*; a, fore and b, hind wings as preserved, $\times 3.0$ (Becker-Migdisova & Wootton, 1965).

Dunstaniopsis TILLYARD, 1918d, p. 584 [**D. triassica*; OD]. Little-known genus, based on incomplete fore wing; apex apparently more pointed than in *Dunstania*. EVANS, 1956; BECKER-MIGDISOVA & WOOTTON, 1965. *Trias.*, Australia (Queensland).

Paradunstania TILLYARD, 1918d, p. 585 [**P. affinis*; OD]. Little-known genus, based on fragment of fore wing; probably a synonym of *Dunstania*. EVANS, 1956; BECKER-MIGDISOVA & WOOTTON,

1965. *Trias.*, Australia (Queensland).—FIG. 135,2. **P. affinis*; fore wing as preserved, $\times 1.5$ (Evans, 1956).

Siksteliana BECKER-MIGDISOVA & WOOTTON, 1965, p. 68 [**S. popovi*; OD]. Little-known genus, based on fore wing. Similar to *Dunstaniodes*, but costal margin of basal half nearly straight. *Trias.*, USSR (Kirghiz).—FIG. 135,4. **S. popovi*; fore wing, $\times 3$ (Becker-Migdisova & Wootton, 1965).

Family PALAEONTINIDAE Handlirsch, 1906

[Palaeontinidae HANDLIRSCH, 1906b, p. 618] [=Cicadomorphae EVANS, 1956, p. 222]

Fore wing as in Dunstaniidae, with membranous, distal part of wing broader and longer than basal, tegminous part; vein SC usually weakly developed, commonly with branches or suggestions of branches; R and M separating before or close to midwing; R and RS nearly straight. Hind wing with a prominent indentation on costal margin; M1 commonly coalesced for short interval with RS; M with 4 branches. Head small, narrow, pronotum wide; body generally with numerous hairs. *Perm.*—*Jur.*

Palaeontina BUTLER, 1873, p. 126 [**P. oolitica*; OD]. Little-known genus, based on fore wing. M with 4 branches, M1+2 and M3+4 forking at about same level. [The genus was excluded from Homoptera by EVANS (1956) but included here by BECKER-MIGDISOVA (1962b) and POPOV (1980b).] *Jur.*, England.—FIG. 136,1. **P. oolitica*; fore wing, $\times 0.8$ (Handlirsch, 1906b).

Asiocossus BECKER-MIGDISOVA, 1962a, p. 89 [**A. subcostalis*; OD]. Little-known genus, based on fragment of fore wing. SC free from R+M except for very base, branched; R+M and stem of R very short. *Trias.*, USSR (Kirghiz).—FIG. 136,3. **A. subcostalis*; fore wing base, $\times 2.5$ (Becker-Migdisova, 1962b).

Cicadomorpha MARTYNOV, 1926b, p. 1357 [**C. punctulata*; OD]. SC coalesced with R+M at base; area between M and CUA very broad, without crossveins; CU slightly arched at base. *Jur.*, USSR (Kazakh).—FIG. 136,7. **C. punctulata*; fore wing, $\times 1.0$ (Becker-Migdisova, 1962b).

Fletcheriana EVANS, 1956, p. 224 [**F. triassica*; OD]. Fore wing as in *Pseudocossus*, but costal area much broader; SC lying alongside R basally; RS arising from R remote from wing base. [The assignment of a species from the Triassic of South Africa (RIEK, 1976b) to this genus is very uncertain.] *Trias.*, Australia (New South Wales).—FIG. 136,2. **F. triassica*; a, fore wing; b, hind wing, $\times 1.0$ (Evans, 1956).




FIG. 136. Palaeontinidae (p. 214–216).

Ijacossus BECKER-MIGDISOVA, 1950, p. 1106 [*I. suchanovae*; OD]. Little-known genus, based on fore wing. Similar to *Palaeontinodes*, but SC with several branches. [Family assignment uncertain.] *Jur.*, USSR (Asian RSFSR).—FIG. 136, 8. *I. suchanovae*; fore wing, $\times 1$ (Becker-Migdisova, 1962b).

Palaeocicadopsis T'AN, 1980, p. 161 [**P. chinensis*; OD]. Fore wing similar to that of *Cicadomorpha*, but M branching near wing base. *Perm.*, China (Inner Mongolia).

Palaeocossus OPPENHEIM, 1885, p. 333 [**P. jurassicus*; OD]. Fore wing without nodal indentation; wing broadly oval; distal margin of basal median




FIG. 137. Palaeontinidae (p. 216–217).

cell (between M and CUA) smoothly curved; hind margin strongly convex. *Jur.*, USSR (Asian RSFSR).—FIG. 136,6. **P. jurassicus*; fore wing, $\times 1.5$ (Evans, 1956).

Palaeontinodes MARTYNOV, 1937a, p. 166 [**P. shabarovi*; OD]. Fore wing triangular; costal indentation weak; SC without branches; crossvein m-cua long; basal median cell divided by crossveins (not shown in figure). BECKER-MIGDISOVA, 1949b; EVANS, 1956. *Jur.*, USSR (Tadzhik, Asian RSFSR).—FIG. 136,5. *P. angarensis* BECKER-MIGDISOVA & WOOTTON; fore wing, $\times 1$ (Becker-Migdisova & Wootton, 1965).

Palaeontinopsis MARTYNOV, 1937a, p. 167 [**P. latipennis*; OD]. Little-known genus. Fore wing apparently oval and with rounded apex. EVANS, 1956; BECKER-MIGDISOVA & WOOTTON, 1965. *Jur.*, USSR (Tadzhik).—FIG. 136,4. **P. latipennis*; fore wing, $\times 1.5$ (Becker-Migdisova, 1962b).

Phragmatocites OPPENHEIM, 1885, p. 333 [**P. damesi*; OD]. Fore wing with costal margin straight or only slightly curved; nodal indentation weak. EVANS, 1956; BECKER-MIGDISOVA & WOOTTON, 1965. *Jur.*, USSR (Asian RSFSR).—FIG. 136,9. **P. damesi*; fore wing, $\times 2.5$ (Becker-Migdisova, 1962b).

Plachutella BECKER-MIGDISOVA, 1949b, p. 11 [**P. rotundata*; OD]. Little-known genus, based on hind wing. M2 close to M3+4 at one point but not coalesced with it. BECKER-MIGDISOVA, 1950; BECKER-MIGDISOVA & WOOTTON, 1965. *Jur.*, USSR (Kazakh, Tadzhik).—FIG. 136,10. **P. rotundata*; hind wing, $\times 2.5$ (Becker-Migdisova, 1949b).

Pseudocossus MARTYNOV, 1931d, p. 94 [**P. zemcuznicovi*; OD]. Fore wing triangular, with pronounced indentation of costal margin at nodal break; SC free from R+M at base, branched; RS arising from R near wing base; distinct bands of coloration. Hind wing rounded, much smaller than fore wing. *Jur.*, USSR (Asian RSFSR, Kazakh).—FIG. 137,2. *P. turgaiensis* BECKER-MIGDISOVA & WOOTTON, Kazakh; a, fore and b, hind wings, $\times 1.5$ (Becker-Migdisova & Wootton, 1965).

Shurabocossus BECKER-MIGDISOVA, 1949b, p. 15 [**S. gigas*; OD]. Hind wing similar to that of *Plachutella*, but M2 coalesced with M3+4 for a considerable interval before separating. *Jur.*, USSR (Tadzhik).—FIG. 137,5. **S. gigas*; hind wing, $\times 1.5$ (Becker-Migdisova, 1962b).

Suljuktaja BECKER-MIGDISOVA, 1949b, p. 17 [**S. turkestanensis*; OD]. Hind wing as in *Shurabocossus* but with the coalesced parts of 1A and 2A at least as long as the free portions. *Jur.*, USSR (Kirghiz).—FIG. 137,4. **S. turkestanensis*; hind wing, $\times 2$ (Becker-Migdisova, 1962b).

Suljuktocossus BECKER-MIGDISOVA, 1949b, p. 8 [**S.*

prosboloides; OD]. Fore wing as in *Phragma-toectites* but more nearly triangular and with apex nearly pointed. *Jur.*, USSR (Kirghiz).—FIG. 137,1. **S. prosboloides*; fore wing, $\times 1.5$ (Becker-Migdisova, 1962b).

Turgaiella BECKER-MIGDISOVA & WOOTTON, 1965, p. 70 [**T. pomerantsevae*; OD]. Fore wing as in *Palaeontinodes*, but wing oval and basal median cell not divided by crossveins; crossvein m4-cua very short. *Jur.*, USSR (Kazakh).—FIG. 137,3. **T. pomerantsevae*; fore wing, $\times 1.5$ (Becker-Migdisova & Wootton, 1965).

Family MESOGEREONIDAE Tillyard, 1921b

[Mesogereonidae TILLYARD, 1921b, p. 272]

Fore wing slender, with well-developed submarginal (ambient) vein and coriaceous border; veins SC and R close together and to costal margin; RS arising before fork of M1 + 2; crossvein m4-cua near wing base and almost longitudinal in position. Hind wing little known, much smaller than fore wing. Body structure unknown. EVANS, 1956; BECKER-MIGDISOVA & WOOTTON, 1965. *Trias.*

Mesogereon TILLYARD in TILLYARD & DUNSTAN, 1916, p. 33 [**M. neuropunctatum*; OD]. RS joined to M1 by a short crossvein; M3 + 4 forking more basally than M1 + 2. *Trias.*, Australia (New South Wales).—FIG. 138,1a. *M. superbum* TILLYARD; fore wing, $\times 1.2$ (Evans, 1956).—FIG. 138,1b. *M. shepherdii* TILLYARD; hind wing, $\times 1.2$ (Evans, 1956).

Triassogereon RIEK, 1976b, p. 808 [**T. distinctum*; OD]. Fore wing as in *Mesogereon*, but fork of M3 + 4 close to fork of M1 + 2. *Trias.*, South Africa.—FIG. 138,2. **T. distinctum*; fore wing, $\times 1.6$ (Riek, 1976b).

Family PROSBOLIDAE Handlirsch, 1906

[Prosbolidae HANDLIRSCH, 1906b, p. 390 [=Sojaneuridae BECKER-MIGDISOVA, 1946, p. 730]

Fore wing: distal part commonly membranous; costal area broad; vein SC usually forming an anterior branch submarginal to costal margin and more rarely an indistinct, short branch that parallels R + M and even part of R; forks of M and CUA usually shallow. Hind wing: costal margin usually deeply excised near middle, convex basally and dis-




FIG. 138. Mesogereonidae (p. 217).

tally; anal region extended posteriorly. Body structure unknown. BECKER-MIGDISOVA, 1940, 1947, 1962b; EVANS, 1956. *Perm.*—*Trias.*

Prosbolole HANDLIRSCH, 1904b, p. 2 [**P. hirsuta*; OD] [=*Prosbolina* HANDLIRSCH, 1937, p. 132 (type, *Prosbolole biexcisa* MARTYNOV, 1928b, p. 7)]. Fore wing: nodal break and nodal line present; R, M, and CUA dividing at about same level. Hind wing: M with at least 4 branches. *Perm.*, USSR (European and Asian RSFSR).—FIG. 139,4a. **P. hirsuta*; fore wing, $\times 1.6$ (Evans, 1956).—FIG. 139,4b. *P. reducta* MARTYNOV; fore wing, $\times 3.5$ (Becker-Migdisova, 1940).—FIG. 139,4c. *P. breviata* BECKER-MIGDISOVA; hind wing, $\times 2.6$ (Becker-Migdisova, 1940).

Austroprosbolole EVANS, 1943b, p. 181 [**A. maculata*; OD]. Fore wing with nodal break and nodal line; RS curved posteriorly, touching M1 + 2 at point of fork; CUA with a shallow, distal fork. EVANS, 1956. *Perm.*, Australia (New South Wales).—FIG. 139,5. **A. maculata*; fore wing, $\times 4$ (Evans, 1943b).

Austroprosboloides RIEK, 1973, p. 527 [**A. vandijki*; OD]. Little-known genus; fore wing similar to *Austroprosbolole*, but RS touching M1 beyond fork and M3 + 4 connected to CUA distally. RIEK, 1976a. *Perm.*, South Africa.—FIG. 140,6. **A. vandijki*; fore wing, $\times 4$ (Riek, 1973).

Beaufortiscus RIEK, 1976a, p. 779 [**B. dixi*; OD]. Fore wing very similar to that of *Prosbolole*; anal




FIG. 139. Prosbolidae (p. 217–220).

area with 3 veins. [Probably a synonym of *Prosbole*.] *Perm.*, South Africa.
Dictyoprosbole MARTYNOV, 1935d, p. 443 [**D. membranosa*; OD]. Fore wing membranous, covered with a network of crossveins; M and CUA dividing at level of origin of RS; RS, M, and CUA with branching as in *Orthoprosbole*. EVANS, 1956. *Perm.*, USSR (Asian RSFSR). —FIG. 139,6. **D. membranosa*; fore wing, $\times 1.5$ (Becker-Migdisova, 1960).
Evanscicada BECKER-MIGDISOVA, 1962b, p. 170,

nom. subst. pro Evansia BECKER-MIGDISOVA, 1961c, p. 323, non CAMBRIDGE, 1900 [**Evansia speciosa* BECKER-MIGDISOVA, 1961c, p. 323; OD]. Fore wing narrow; basal part tegminous; RS arising at level of forking of M; numerous crossveins distally and indication of network near basal-central part of wing. *Perm.*, USSR (Asian RSFSR). —FIG. 139,1. **E. speciosa*; fore wing, $\times 2.5$ (Becker-Migdisova, 1962b).

Falsia BECKER-MIGDISOVA, 1946, p. 750 [**F. chimaera*; OD]. Similar to *Sojanoneura*, but first




FIG. 140. Prosbolidae (p. 217–221).

and second tarsal segments of same size. BECKER-MIGDISOVA, 1946. *Perm.*, USSR (European RSFSR).

Kaltanetta BECKER-MIGDISOVA, 1961c, p. 303 [**K. nigra*; OD]. Fore wing slender, apex symmetrically curved; RS arising well before level of forking of M and of CUA; M with 3 branches. Hind wing slender distally; marginal indentation deep and wide; M with 3 branches. *Perm.*, USSR (Asian RSFSR).—FIG. 139,2. **K. nigra*; a,

fore and b, hind wings, $\times 6.5$ (Becker-Migdisova, 1961c).

Kaltanopsis BECKER-MIGDISOVA, 1961c, p. 300 [**K. ornata*; OD]. Fore wing similar to that of *Kaltanetta*, but costal margin strongly curved and R continuing in a straight line from its stem; longitudinal veins unusually thick. *Perm.*, USSR (Asian RSFSR).—FIG. 139,3. **K. ornata*; fore wing, $\times 8$ (Becker-Migdisova, 1961c).

Kondomoprosbole BECKER-MIGDISOVA, 1961c, p.




FIG. 141. Prosboldidae (p. 220).

315 [**K. pictata*; OD]. Hind wing: anterior margin with prominent bulge distally; R a straight continuation of stem R; M with 3 short branches. *Perm.*, USSR (Asian RSFSR).—FIG. 139,7. **K. pictata*; hind wing, $\times 8.5$ (Becker-Migdisova, 1962b).

Leptoprosbole RIEK, 1976b, p. 812 [**L. lepida*; OD]. Fore wing elongate; M with 8 terminal branches. [Family assignment doubtful.] *Trias.*, South Africa.—FIG. 140,2. **L. lepida*; fore wing, $\times 1.5$ (Riek, 1976b).

Mesocicada BECKER-MIGDISOVA, 1962a, p. 90 [**M.*

verrucosa; OD]. Little-known fore wing; nodal break absent; M with 4 branches; CUA with small fork. [Family assignment doubtful.] *Trias.*, USSR (Kirghiz).—FIG. 140,1. **M. verrucosa*; fore wing, $\times 14$ (Becker-Migdisova, 1962a).

Mitchelloneura TILLYARD, 1921c, p. 414 [**M. permiana*; OD]. Little-known hind wing; RS with irregular distal branches; M with M1, M2, and M3+4; CUA deeply forked. EVANS, 1956. *Perm.*, Australia (New South Wales).—FIG. 140,7. **M. permiana*; hind wing, $\times 3.2$ (Tillyard, 1921c).

Neurobole RIEK, 1976a, p. 779 [**N. ramosa*; OD]. Little-known genus, based on small apical fragment of wing. [Family assignment doubtful.] *Perm.*, South Africa.

Orthoprosbole MARTYNOV, 1935d, p. 445 [**O. congesta*; OD]. Fore wing strongly narrowed in distal half; RS and M with numerous branches; nodal break prominent. Hind wing little known; distal part elongate; M and CUA with numerous branches. BECKER-MIGDISOVA, 1961c. *Perm.*, USSR (Asian RSFSR).—FIG. 140,4a. *O. triangularis* (MARTYNOV); fore wing, $\times 2.5$ (Becker-Migdisova, 1962b).—FIG. 140,4b. **O. congesta*; hind wing, $\times 3.5$ (Becker-Migdisova, 1962b).

Orthoscytina TILLYARD, 1926a, p. 9 [**O. mitchelli*; OD]. Fore wing slender, oval; anal area long; RS arising just before midwing; M and CUA forked at distal third of wing; R with several oblique branches to costal margin. Hind wing little known. EVANS, 1956; RIEK, 1976a. *Perm.*, Australia (New South Wales), Africa (South Africa), USSR (Asian RSFSR).—FIG. 141,1a. **O. mitchelli*, Australia; fore wing, $\times 6$ (Tillyard, 1926a).—FIG. 141,1b. *O. suchovi* BECKER-MIGDISOVA, USSR; fore wing, $\times 6$ (Becker-Migdisova, 1961c).

Permocicada MARTYNOV, 1928b, p. 19 [**P. umbrata*; SD BECKER-MIGDISOVA, 1940, p. 29] [= *Permocicadopsis* BECKER-MIGDISOVA, 1940, p. 54 (type, *Permocicada angusta* MARTYNOV, 1935c, p. 15)]. Fore wing with weak venation; RS arising before forking of M and CUA; M with 3 or 4 branches. Hind wing with deeply indented costal margin having nearly symmetrical slopes. ZALESSKY, 1929, 1932b; EVANS, 1956; BECKER-MIGDISOVA, 1961c, 1962b. *Perm.*, USSR (European and Asian RSFSR).—FIG. 141,2. *P. integra* BECKER-MIGDISOVA; a, fore wing, $\times 4$; b, hind wing, $\times 4$; c, reconstruction, $\times 3$ (Becker-Migdisova, 1940).

Permodiphthera TILLYARD, 1926a, p. 24 [**P. robusta*; OD]. Little-known genus. Fore wing with RS unbranched; branches of M apparently strongly curved. *Perm.*, Australia (New South Wales).—FIG. 140,5. **P. robusta*; fore wing, $\times 6$ (Evans, 1956).

Pervestigia BECKER-MIGDISOVA, 1961c, p. 318 [**P.*

veteris; OD]. Hind wing: anterior margin without distal hump; M with 3 branches; CUA with narrow fork distally. *Perm.*, USSR (Asian RSFSR).—FIG. 140,8. **P. veteris*; hind wing, $\times 3$ (Becker-Migdisova, 1961c).

Prosbolomorpha RIEK, 1974c, p. 21 [**P. clara*; OD]. Fore wing as in *Austroprosbolus*, but RS not coalesced with M; M3+4 forking at its point of origin. [Probably a synonym of *Austroprosbolus*.] *Trias.*, South Africa.

Prosboloneura BECKER-MIGDISOVA, 1961c, p. 305 [**P. colorata*; OD]. Fore wing shaped as in *Sojanoneura*, but CUA more deeply forked and M with 3 branches. *Perm.*, USSR (Asian RSFSR).—FIG. 140,3. *P. kondonensis* BECKER-MIGDISOVA; fore wing, $\times 8$ (Becker-Migdisova, 1961c).

Sojanoneura MARTYNOV, 1928b, p. 22 [**S. edemskii*; SD BECKER-MIGDISOVA, 1940, p. 44]. Fore wing oval, bluntly rounded; RS arising nearer wing apex than in *Dictyoprosbole*; M with 3 or 4 branches. Hind wing little known, with only a slight bulging of the costal margin basally; M with 2 or 3 branches. MARTYNOV, 1935c; EVANS, 1956. *Perm.*, USSR (European and Asian RSFSR).—FIG. 140,9. *S. stigmata* MARTYNOV; fore wing, $\times 4$ (Becker-Migdisova, 1962b).

Family CICADOPROSBOLIDAE

Evans, 1956

[Cicadoprobsolidae EVANS, 1956, p. 222]

Apparently related to Prosbolidae. Fore wing with vein M forking at midwing; RS arising before midwing; short, supplementary veins between branches of R; nodal line distinct, crossing RS remote from origin of RS and crossing M beyond its first fork. *Trias.*

Cicadoprobsolus BECKER-MIGDISOVA, 1947, p. 445 [**C. sogutensis*; OD]. Fore wing oval, apex slightly asymmetrical; branches of M and CUA slightly curved and parallel. [Originally placed in the family Prosbolidae but transferred to a new family, Cicadoprobsolidae, by EVANS (1956) and later to the Tettigarctidae by BECKER-MIGDISOVA, 1962b.] *Trias.*, USSR (Kirghiz).—FIG. 142,4. **C. sogutensis*; fore wing, $\times 3.5$ (Becker-Migdisova, 1947).

Family TETTIGARCTIDAE

Distant, 1905

[Tettigarctidae DISTANT, 1905, p. 280]

Fore wing with transparent membranous area; costa broadly sclerotized; apical border narrow; venation much as in *Cicadidae*; vein SC with a short, hook-shaped anterior branch




FIG. 142. Cicadoprobsolidae, Tettigarctidae, and Cicadidae (p. 221–222).

basally and a posterior branch coalesced with R+M and R. Hind wing: front margin with shallow indentation and convex area. Body structure: stridulatory organ present in both sexes. [This family is known by only one living genus, *Tettigarcta*, occurring in Australia. There is a reliable Tertiary record of the family, but the several Mesozoic genera that have been placed here are very poorly known and are assigned in this account to the category, family uncertain.] EVANS, 1956; WHALLEY, 1983. *Eoc.*—*Holo.*

Tettigarcta WHITE, 1845, p. 412. *Holo.*

Eotettigarcta ZEUNER, 1944a, p. 110 [**E. scotica*; OD]. Hind wing similar to that of *Tettigarcta* (recent), but indentation of costal margin much longer; origin of posterior branch of RS more remote from base of wing. Fore wing unknown.

WHALLEY, 1983. *Eoc.*, Scotland.—FIG. 142,3.
**E. scotica*; hind wing, $\times 2.5$ (Zeuner, 1944a).

Family CICADIDAE Leach, 1815

[Cicadidae LEACH, 1815, p. 124]

Fore wing with costal area reduced to a narrow strip or absent; apical parts of distal forks of veins aligned to form a submarginal vein along the outer and hind margins; anal area narrow and short. Hind wing much smaller than fore wing; anterior margin smooth; submarginal vein formed as in fore wing; anal-jugal area slightly broader than in fore wing. Stridulatory organs (tymbals) present on dorsum of first abdominal segment, at least in males. [A fragmented specimen from the Eocene of France was described as *beauchampi* by Piton (1940a); this species was placed in the existing genus *Chemsitica* STÅL (= *Ribana* DISTANT). However, the fossil does not show enough structural detail for family assignment. See also *Liassocicada* under Homoptera, family Uncertain.] COOPER 1941; WHALLEY, 1983. *Paleoc.*—*Holo.*

Cicada LINNÉ, 1758, p. 434. COOPER, 1941; WHALLEY, 1983. *Oligo.*, USA (Colorado); *Mio.*, Europe (Yugoslavia, Germany)—*Holo.*

Davisia COOPER, 1941, p. 288 [**D. bearcreekensis*; OD]. Similar to *Tibicen*; cell *cua2* broad but slightly more than twice as long as wide; apical margin of cell *cua1* evenly and shallowly curving into cell *cua2*. WHALLEY, 1983. *Paleoc.*, USA (Montana).—FIG. 142,2. **D. bearcreekensis*; fore wing, $\times 1.0$ (Cooper, 1941).

Lithocicada COCKERELL, 1906c, p. 457 [**L. perita*; OD]. Similar to *Cicada*, but cubital cell of fore wing with pointed or narrowly truncate apex. COOPER, 1941. *Oligo.*, USA (Colorado).

Platypedia UHLER, 1888, p. 23. COCKERELL, 1908a; COOPER, 1941. *Oligo.*, USA (Colorado)—*Holo.*

Tibicen LATREILLE, 1825, p. 426. SCUDDER, 1892; COOPER, 1941. *Oligo.*, USA (Colorado)—*Holo.*

Tymocicada BECKER-MIGDISOVA, 1954, p. 799 [**T. gorbunovi*; OD]. Fore wing similar to that of *Cosmopsaltivia* (recent), but CUA with longer anterior branch; cell between R and RS slightly broader. *Mio.*, USSR (Asian RSFSR).—FIG. 142,1. **T. gorbunovi*; fore wing, $\times 1.4$ (Becker-Migdisova, 1954).

Family SCYTINOPTERIDAE Handlirsch, 1906b

[Scytinopteridae HANDLIRSCH, 1906b, p. 391]

Fore wing tegminous; costal margin commonly thickened basally; veins usually thin;

vein SC obsolescent; branches of M and CUA short; crossveins few; only a few closed cells between R, M, and CUA. Hind wing with costal margin with at most a shallow convexity at base of wing; RS unbranched; M and CUA distally branched. Body little known, apparently as in Cicadellidae. *Perm.*—*Trias.*

Scytinoptera HANDLIRSCH, 1904b, p. 3 [**S. kokeni*; OD] [= *Anomoscyta* MARTYNOV, 1928b, p. 34 (type, *A. reducta*); *Permocixius* MARTYNOV, 1928b, p. 36 (type, *P. kazanensis*); *Scytinoptera curta* ZALESKY, 1929, p. 28]. Fore wing with posterior branch of SC short, forming a sharp curve at level of R+M, or absent; M and CUA with distal forks, forming series of small, marginal cells, usually subequal; anal-jugal region strongly widened. Hind wing with costal margin with conspicuous but gradual convexity near base; no prominent marginal concavity or excision. Pronotum with lateral projections. [RIEK (1976b) has described a late Triassic species (*distorta*) in the genus *Scytinoptera*, but there is really no evidence to justify that placement.] *Perm.*, USSR (European and Asian RSFSR).—FIG. 143,5a,b. *S. kaltanica* BECKER-MIGDISOVA; a, fore and b, hind wings, $\times 10$ (Becker-Migdisova, 1962b).—FIG. 143,5c. *S. picturata* BECKER-MIGDISOVA; fore wing, $\times 8$ (Becker-Migdisova, 1961c).

Anaprosbole BECKER-MIGDISOVA, 1960, p. 28 [**A. ivensis*; OD]. Fore wing with costal margin relatively broad basally; RS arising well beyond midwing; branches of M1+2 much longer than branches of M3+4; CUA with 3 terminal branches. [Family assignment uncertain.] *Perm.*, USSR (European RSFSR).—FIG. 143,6. **A. ivensis*; fore wing, $\times 5.0$ (Becker-Migdisova, 1960).

Anomaloscytina DAVIS, 1942, p. 112 [**A. metapteryx*; OD]. Hind wing with costal margin with distinct but gentle concavity; SC short but distinct; anal area extensive. [Family position uncertain.] *Perm.*, Australia (New South Wales).—FIG. 143,7. **A. metapteryx*; hind wing, $\times 6.5$ (Davis, 1942).

Elliptoscarta TILLYARD, 1926a, p. 16 [**E. ovalis*; OD]. Fore wing oval, with apex evenly rounded; costal area (between C and R) broad; R dichotomously forked; M with 5 branches; CUA forked. *Perm.*, Australia (New South Wales).—FIG. 143,1. **E. ovalis*; fore wing, $\times 8.2$ (Tillyard, 1926a).

Homaloscytina TILLYARD, 1926a, p. 16 [**H. plana*; OD]. Fore wing as in *Anaprosbole*, but CUA with only 2 terminal branches and connected to M by a crossvein; apex of wing bluntly rounded. EVANS, 1943b. *Trias.*, Australia (New South Wales).—FIG. 143,4. **H. plana*; fore wing, $\times 8$ (Evans,




FIG. 143. Scytinopteridae (p. 222–223).

Ingruo BECKER-MIGDISOVA, 1960, p. 19 [*I. lanceolata*; OD]. Fore wing very narrow; posterior branch of SC short, merging with R; CUA dividing at level of origin of RS; fork of CUA large. [Family assignment doubtful]. *Perm.*, USSR (European RSFSR).—FIG. 143,2. *I. lanceolata*; fore wing, $\times 16$ (Becker-Migdisova, 1960).

Ivaia BECKER-MIGDISOVA, 1960, p. 25 [*I. indistincta*; OD]. Fore wing moderately broad; costal area (between C and R) broad; R straight; CUA in brief contact with M, then diverging; M apparently unbranched. *Perm.*, USSR (European RSFSR).—FIG. 143,3a. *I. procucopoides* BECKER-MIGDISOVA; fore wing, $\times 6.5$ (Becker-

Migdisova, 1960).—FIG. 143,3b. *I. indistincta*; fore wing, $\times 8$ (Becker-Migdisova, 1962b).

Kaltanospes BECKER-MIGDISOVA, 1961c, p. 344 [*K. kuznetskiensis*; OD]. Fore wing as in *Ingruo*, but CUA dividing much further distally of origin of RS. *Perm.*, USSR (Asian RSFSR).—FIG. 144,6. *K. kuznetskiensis*; fore wing and body, $\times 10$ (Becker-Migdisova, 1961c).

Mesonirvana EVANS, 1956, p. 191 [*M. abrupta*; OD]. Fore wing: R with several branches; cross-vein m-cu joined to CUA1; RS unbranched. *Trias.*, Australia (Queensland).—FIG. 144,7. *M. abrupta*; fore wing, $\times 5$ (Evans, 1956).

Mesothymbris EVANS, 1956, p. 191 [*M. perkinsi*; OD]. Fore wing: R with several branches; cross-vein m-cu joined to CUA1; RS unbranched. *Trias.*, Australia (Queensland).—FIG. 144,8. *M. perkinsi*; fore wing, $\times 5$ (Evans, 1956).




FIG. 144. Scytinopteridae (p. 223–225).

OD]. Fore wing as in *Triassoscytinopsis*, but M1+2 forming almost straight continuation of M; M3+4 bent towards CUA; crossvein m-cu joined to CUA. *Trias.*, Australia (Queensland). —FIG. 144,3. **M. perkinsi*; fore wing, $\times 6$ (Evans, 1956).

Permododa BECKER-MIGDISOVA, 1961c, p. 347 [**P. membracoides*; OD]. Fore wing very slender; several closed cells between RS and M and between M and CUA. *Perm.*, USSR (Asian RSFSR). —FIG. 144,2. **P. membracoides*; fore wing, $\times 10$ (Becker-Migdisova, 1961c).

Permojassus TILLYARD, 1926a, p. 7 [**P. australis*; OD] [= *Permojassula* HANDLIRSCH, 1937, p. 115, obj.] Fore wing similar to that of *Homaloscytina*, but anal area apparently much narrower. EVANS,

1956. *Perm.*, Australia (New South Wales). —FIG. 144,8. **P. australis*; fore wing, $\times 8$ (Evans, 1956).

Permolamproptera BECKER-MIGDISOVA, 1961c, p. 340 [**P. grandis*; OD]. Hind wing similar to that of *Scytinoptera* but with R2 curved distally; anal area extended. *Perm.*, USSR (Asian RSFSR). —FIG. 144,9. **P. grandis*; hind wing, $\times 5.7$ (Becker-Migdisova, 1961c).

Sarbaloptera BECKER-MIGDISOVA, 1961c, p. 328 [**S. sarbalensis*; OD]. Fore wing with asymmetrical apex; costal area (between C and R) very broad. *Perm.*, USSR (Asian RSFSR). —FIG. 144,11. **S. sarbalensis*; fore wing, $\times 6.5$ (Becker-Migdisova, 1961c).

Stenoscytina TILLYARD, 1926a, p. 15 [**S. aus-*

traliensis; OD]. Little-known genus. Fore wing narrow; M with 4 branches; CUA curving abruptly posteriorly after diverging from M. [Family assignment uncertain.] *Perm.*, Australia (New South Wales).—FIG. 144,4. **S. australiensis*; fore wing, $\times 6.5$ (Tillyard, 1926a).

Surijokovia BECKER-MIGDISOVA, 1961c, p. 342 [**S. lata*; OD]. Little-known fore wing; posterior branch of SC long, mostly parallel to R, then diverging anteriorly to termination on costal margin; RS apparently unbranched. *Perm.*, USSR (Asian RSFSR).—FIG. 144,10. **S. lata*; fore wing, $\times 16$ (Becker-Migdisova, 1961c).

Triassoscelis EVANS, 1956, p. 192 [**T. anomala*; OD]. Fore wing as in *Mesonirvana*, but RS forked. *Trias.*, Australia (Queensland).—FIG. 144,5. **T. anomala*; fore wing, $\times 5$ (Evans, 1956).

Triassoscytina EVANS, 1956, p. 179 [**T. incompleta*; OD]. Fore wing as in *Homaloscytina*, but M forking just beyond level of origin of RS. [Family position uncertain.] *Trias.*, Australia (Queensland).

Triassoscytinopsis EVANS, 1956, p. 190 [**T. stenulata*; OD]. Fore wing with apex evenly rounded; R with at least 4 parallel branches distally; RS with from 2 to 4 branches; M with 4 branches. *Trias.*, Australia (Queensland).—FIG. 144,1. *T. aberrans* EVANS; fore wing, $\times 6$ (Evans, 1956).

Tychoscytina BECKER-MIGDISOVA, 1952, p. 179 [**T. kuznetskiensis*; OD]. Fore wing little known, with wide costal area; R1 straight. *Perm.*, USSR (Asian RSFSR).

Family BITURRITIIDAE Metcalf, 1951

[Biturritiidae METCALF, 1951, p. 11]

Fore wing sclerotized; no marginal border; vein M unbranched; radial cell divided by a crossvein. Hind wing nearly of uniform width, with very slight concavity of costal margin. *Trias.—Holo.*

Biturritia GODING, 1930, p. 39. *Holo.*

Absoluta BECKER-MIGDISOVA, 1962a, p. 92 [**A. distincta*; OD]. Hind wing with base of CUA nearly or completely coalesced with stem of M. [Family assignment doubtful.] *Trias.*, USSR (Kirghiz).—FIG. 145,1. **A. distincta*; hind wing, $\times 12$ (Becker-Migdisova, 1962b).

Family CICAPELLIDAE Latreille, 1802

[Cicadellidae LATREILLE, 1802a, p. 257] [=Jasscopidae HAMILTON, 1971, p. 943]

Fore wing tegminous; several to many closed cells; CUA usually with wide distal fork. Hind wing narrowed distally; submar-




FIG. 145. Biturritiidae and Cicadellidae (p. 225–227).

ginal vein conspicuous, with relatively wide margin; ano-jugal area broad. *Trias.—Holo.*

Cicadella DUMÉRIl, 1806, p. 266. BERVOETS, 1910; COCKERELL, 1920c; BECKER-MIGDISOVA, 1951. *Eoc.*, USA (Colorado); *Oligo.*, Europe (Baltic); *Mio.*, USSR (European RSFSR)—*Holo.*

Acocephalites MEUNIER, 1904e, p. 119 [**A. bred dini*; OD]. Little-known genus, based on fore wing with strongly arched costal margin and a venation similar to that of *Mesojassoides*; M with distal fork. *Jur.*, Europe (Spain).—FIG. 145,5. **A. bred dini*; fore wing, $\times 14$ (Handlirsch, 1907).

Agallia CURTIS, 1833, p. 193. SCUDDER, 1890. *Oligo.*, USA (Colorado)—*Holo.*

Aphrodes CURTIS, 1833, p. 193. SCUDDER, 1890;




FIG. 146. Eurymelidae and Membracidae (p. 227–228).

- STATZ, 1950a. *Eoc.*, USA (Wyoming); *Oligo.*–*Mio.*, Europe (Germany)–*Holo.*
- Batrachomorphus** LEWIS, 1834, p. 51, *nom. correct.* AGASSIZ, 1846, *ex Batrachomorphus* LEWIS, 1834. STATZ, 1950a. *Oligo.*, Europe (Germany)–*Holo.*
- Cicadula** ZETTERSTEDT, 1838, p. 296. SCUDDER, 1890. *Eoc.*, USA (Wyoming)–*Holo.*
- Coelidia** GERMAR, 1821, p. 75. SCUDDER, 1890, 1895b. *Eoc.*, USA (Wyoming), Canada (British Columbia); *Oligo.*, USA (Colorado)–*Holo.*

- Deltocephalus** BURMEISTER, 1838, p. 5. STATZ, 1950a. *Oligo.*, Europe (Germany)–*Holo.*
- Durgades** DISTANT, 1912, p. 608. BECKER-MIGDISOVA & MARTYNOVA, 1951. *Mio.*, USSR (Kirghiz)–*Holo.*
- Eurymelidium** TILLYARD, 1919c, p. 884 [**E. australe*; OD]. SC apparently absent; M1 + 2 anastomosed with part of RS. EVANS, 1956. *Trias.*, Australia (Queensland).—FIG. 145, 3. **E. australe*; fore wing, $\times 10$ (Evans, 1956).
- Euscelis** BRULLÉ, 1832, p. 109. STATZ, 1950a; PIERCE, 1963. *Oligo.*, Europe (Germany); *Mio.*, USA (California)–*Holo.*
- Gypona** GERMAR, 1821, p. 73. [Generic assignment of fossil doubtful.] SCUDDER, 1890. *Oligo.*, USA (Colorado)–*Holo.*
- Gyponites** STATZ, 1950a, p. 10 [**G. pronota*; OD]. Little-known cicadellid; head short and broad; ocelli large; pronotum long, with parallel sides; scutellum shorter than pronotum. *Oligo.*, Europe (Germany).
- Homopterulum** HANDLIRSCH, 1907, p. 641 [**Cercopidium signoreti* WESTWOOD, 1854, p. 396; SD CARPENTER, herein]. Little-known genus, based on fore wing. [Family assignment doubtful.] EVANS, 1956. *Jur.*, England.
- Idiocerus** LEWIS, 1834, p. 47. STATZ, 1950a. *Oligo.*, Europe (Germany)–*Holo.*
- Jascopus** HAMILTON, 1971, p. 944 [**J. notabilis*; OD]. Little-known genus, based on nymph. [Type of family *Jascopidae* HAMILTON.] EVANS, 1972. *Cret.*, Canada (Manitoba).
- Jassites** HANDLIRSCH, 1907, p. 642 [**Cicada punctatus* BRODIE, 1845, p. 33; OD]. Little-known genus, based on fore wing. CUA with very short branches. EVANS, 1956. *Jur.*, England.—FIG. 145, 2. **J. punctatus*; fore wing, $\times 5.5$ (Evans, 1956).
- Jassopsis** SCUDDER, 1890, p. 312 [**J. evidens*; OD]. Little-known genus, similar to *Thamnotettix*. Scutellum not more than half the length of thorax. *Oligo.*, USA (Colorado).
- Jassus** FABRICIUS, 1803, p. 85. BERVOETS, 1910; MEUNIER, 1920c; PITON, 1940a; STATZ, 1950a. *Eoc.*, Europe (France); *Oligo.*, Europe (Baltic, Germany)–*Holo.*
- Lavrushinia** COCKERELL, 1925g, p. 10 [**L. elegantula*; OD]. Little-known genus, based on long and narrow fore wing; marginal vein very close to wing margin. *Mio.*, USSR (Asian RSFSR).
- Macropsis** LEWIS, 1834, p. 49. BERVOETS, 1910; STATZ, 1950a. *Oligo.*, Europe (Baltic, Germany)–*Holo.*
- Maleojassus** ZEUNER, 1941a, p. 90 [**M. primitivus*; OD]. Fore wing as in *Stonastla* (recent), but RS smoothly curved, not bent at junction with M1; M almost straight. *Eoc.*, Scotland.
- Megophthalmus** CURTIS, 1833, p. 193. STATZ, 1950a. *Oligo.*, Europe (Germany)–*Holo.*
- Mesojassoides** OMAN, 1937, p. 38 [**M. gigantea*;

OD]. Fore wing as in *Coelidia* but with additional crossveins; M unbranched. *Cret.*, USA (Colorado).—FIG. 145,4. **M. gigantea*; fore wing, $\times 4.5$ (Evans, 1956).

Miochlorotettix CARPENTER, herein [**M. gibroni* PIERCE, 1963, p. 73; OD]. Similar to *Chlorotettix* (recent), but prothorax strongly arched forward and scutellum extending back between wings about as far as forwards. The original generic name, *Miochlorotettix*, was a *nomen nudum* (PIERCE, 1963.) *Mio.*, USA (California).

Miomesamia PIERCE, 1963, p. 81 [**M. juliae*; OD]. Similar to *Ulope* (recent). Face wide, eyes prominent; antennae at sides of front sutures, opposite outer corners of eyes. *Mio.*, USA (California).

Oligogypona STATZ, 1950a, p. 8 [**O. haupiti*; OD]. Similar to *Gypona*, but head broad, somewhat narrower than pronotum; costal margin of fore wing strongly arched. *Oligo.*, Europe (Germany).

Oligoidiocerus STATZ, 1950a, p. 15 [**O. pronotalis*; OD]. Similar to *Idiocerus* but with richer venation and unmarked fore wing. *Oligo.*, Europe (Germany).

Oligopenthimia STATZ, 1950a, p. 9 [**O. ovalis*; OD]. Similar to *Penthimia* (recent). Head short, as wide as pronotum; scutellum long, reaching to the middle of the abdomen. *Oligo.*, Europe (Germany).

Phlepsius FIEBER, 1866, p. 503. PIERCE, 1963. *Mio.*, USA (California)—*Holo.*

Protochlorotettix PIERCE, 1963, p. 78 [**P. calico*; OD]. Similar to *Chlorotettix* (recent), but with last sternum completely divided. *Mio.*, USA (California).

Tetigonia BLANCHARD, 1852, p. 282. STATZ, 1950a. *Oligo.*, Europe (Germany)—*Holo.*

Tettigella CHINA & FENNAH, 1945, p. 711. SCUDDER, 1890; STATZ, 1950a. *Eoc.*, USA (Wyoming); *Oligo.*, USA (Colorado), Europe (Germany)—*Holo.*

Thamnotettix ZETTERSTEDT, 1838, p. 292. COCKERELL, 1920c, 1924a, 1925a; STATZ, 1950a. *Eoc.*, USA (Colorado); *Oligo.*, Europe (Germany); *Mio.*, USA (Colorado)—*Holo.*

Typhlocyba GERMAR, 1833, p. 180. GERMAR & BERENDT, 1856. *Oligo.*, Europe (Baltic)—*Holo.*

Family EURYMELIDAE

Amyot & Serville, 1843

[Eurymelidae AMYOT & SERVILLE, 1843, p. 554]

Fore wing hyaline or opaque and coriaceous; venation often reticulate; vein RS absent; M1+2 retained as separate vein, extending to apex; M3+4 usually unbranched; CUA forked. *Trias.*—*Holo.*

Eurymela LE PELETIER & SERVILLE, 1828, p. 603. *Holo.*

Mesojassus TILLYARD in TILLYARD & DUNSTAN, 1916,

p. 34 [**M. ipswiciensis*; OD]. Little-known genus, based on fore wing. Fork of CUA marginal, very shallow. EVANS, 1956. *Trias.*, Australia (Queensland).—FIG. 146,1. **M. ipswiciensis*; fore wing, $\times 8.4$ (Evans, 1956).

Family MEMBRACIDAE

Rafinesque, 1815

[Membracidae RAFINESQUE, 1815, p. 121]

Fore wing usually membranous, except for basal region; clavus distinct, claval suture along vein 1A; ends of veins usually forming a scalloped submarginal line, the terminal marginal membrane (limbus) extending beyond the veins; veins usually clear and marked by punctures; M either free basally or coalesced in part with stem of R or CUA; cells usually irregular; venation highly diverse. Hind wing well developed, but usually shorter than fore wing; limbus usually present; venation usually similar to that of fore wing. Pronotum extensively developed, often prolonged posteriorly and concealing the scutellum, the wings, and even the entire abdomen; antennae minute, bristlelike; tarsi with 3 segments. *Trias.*—*Holo.*

This is a very large and diversified family. Fossil forms, which are usually known only from wings, are often difficult to classify because of the variability in the venation, especially that of the fore wings. Much difference of opinion exists among specialists in Homoptera about the generic lines. The taxonomic groups used here are essentially those employed in the *General Catalogue of the Homoptera* (METCALF & WADE, 1966).

Membracis FABRICIUS, 1775, p. 675. *Holo.*

Ceresopsis BECKER-MIGDISOVA, 1958, p. 66 [**C. costalis*; OD]. Fore wing broader than in *Darniopsis*, with conspicuous sclerotized area between costal margin and R+M basally; 3 apical cells. *Trias.*, USSR (Kirghiz).—FIG. 146,4. **C. costalis*; fore wing, $\times 10$ (Becker-Migdisova, 1958).

Darniopsis BECKER-MIGDISOVA, 1958, p. 65. [**D. tragopea*; OD]. Fore wing elongate, with very wide limbus; costal margin only slightly convex; 4 apical cells; M and CUA with common stem; anal area large, triangular. *Trias.*, USSR (Kirghiz).—FIG. 146,3. **D. tragopea*; fore wing, $\times 10$ (Becker-Migdisova, 1958).

Maguviopsis BECKER-MIGDISOVA, 1953c, p. 463 [**M. kotchnevi*; OD]. Fore wing with costal-




FIG. 147. Eoscarterellidae (p. 228).

distal margin broadly curved; anterior half of wing sclerotized; R and M not coalesced with CUA; M unbranched, straight. *Jur.*, USSR (Asian RSFSR).—FIG. 146,5. **M. kotchnevi*; fore wing, $\times 16$ (Becker-Migdisova, 1953c).

Minuta BECKER-MIGDISOVA, 1958, p. 64 [**M. heteropterata*; OD]. Little-known genus; fore wing short, sclerotized from midwing to front margin; R, M, and CUA merged with CUP basally; CUA long; no apical cells. *Trias.*, USSR (Kirghiz).—FIG. 146,6. **M. heteropterata*; fore wing, $\times 20$ (Becker-Migdisova, 1958).

Sphongophoriella BECKER-MIGDISOVA, 1958, p. 63 [**S. reticulata*; OD]. Fore wing elongate, reticulate; M and CUA not coalesced to form a common stem; venation reduced; cells elongate-oval; anal area narrow, 3 apical cells. *Trias.*, USSR (Kirghiz).—FIG. 146,2. **S. reticulata*; fore wing, $\times 34$ (Becker-Migdisova, 1958).

Family AETALIONIDAE Spinola, 1850

[Aetalionidae SPINOLA, 1850, p. 53]

Similar to Cicadellidae; fore wing with vein RS absent; M1+2 (or M1 and M2) extending to apex of wing; M3+4 usually forked; CUA unbranched. Hind wing with RS absent. *Oligo.*—*Holo.*

Aetalion LATREILLE, 1810, p. 263. STATZ, 1950a. *Oligo.*, Europe (Germany)—*Holo.*

Family EOSCARTERELLIDAE

Evans, 1956

[Eoscarterellidae EVANS, 1956, p. 220]

Fore wing with vein RS arising from R about one-third wing length from base; R with at least 2 branches; CUA separate from M. *Perm.*—*Trias.*

Eoscarterella EVANS, 1956, p. 220 [**E. media*; OD] [= *Prosbolopsites* BECKER-MIGDISOVA, 1960, p. 90 (type, *P. illyardi*)]. Fore wing tegminous and rugose, broadest in distal half; RS and M parallel for most of their lengths; M with 4 branches. EVANS, 1961. *Trias.*, Australia (Queensland).—FIG. 147,3. **E. media*; fore wing, $\times 5.5$ (Evans, 1956).

Belmontocarta EVANS, 1958, p. 112 [**B. perfecta*; OD]. Fore wing with SC very short, curving distally towards R+M; M1 and M2 longer than M3 and M4; CUA curved and joined to base of M by short crossvein. *Perm.*, Australia (New South Wales).—FIG. 147,1. **B. perfecta*; fore wing, $\times 4.5$ (Evans, 1958).

Eoscartoides EVANS, 1956, p. 220 [**E. bryani*; OD]. Fore wing with complete marginal border; R and M arched basally; M1+2 forked. EVANS, 1961. *Trias.*, Australia (Queensland).—FIG. 147,2. **E. bryani*; fore wing, $\times 4.5$ (Evans, 1961).

Family PROCERCOPIDAE

Handlirsch, 1906

[Procercopidae HANDLIRSCH, 1906b, p. 500]

Fore wing slender, at least three times as long as wide; vein RS arising in basal third of wing; M and CUA branching in distal third of wing, their branches short. Hind wing very little known. EVANS, 1956. *Trias.*—*Jur.*

Procercopis HANDLIRSCH, 1906b, p. 500 [**P. alutacea*; SD BECKER-MIGDISOVA, 1962b, p. 180]. Fore wing elongate, about 4 times as long as broad; M with at least 3 branches; several crossveins in distal part of wing. *Trias.*, USSR (Kirghiz); *Jur.*, Europe (Germany).—FIG. 148,3. *P. longipennis* BECKER-MIGDISOVA, *Trias.*; fore wing, $\times 4$ (Becker-Migdisova, 1962b).

Procercopina MARTYNOV, 1937a, p. 99 [**P. asiatica*; OD]. Fore wing as in *Procercopis* but relatively broader; only one crossvein between adjacent veins. EVANS, 1956. *Jur.*, USSR (Kirghiz).—FIG. 148,5. **P. asiatica*; fore wing, $\times 4.6$ (Becker-Migdisova, 1962b).




FIG. 148. Procercopidae, Cercopidae, and Archijassidae (p. 228-233).

Family DYSMORPHOPTILIDAE

Handlirsch, 1906

[Dysmorphoptilidae HANDLIRSCH, 1906b, p. 492]

Tegmen of irregular form, abruptly narrowed distally, strongly sclerotized; vein SC apparently fused with R; several short

branches from R to costal margin; RS arising before midwing. EVANS, 1956. *Trias.-Jur.*

Dysmorphoptila HANDLIRSCH, 1906b, p. 492 [**Belostoma liasina* GIEBEL, 1856, p. 371; OD]. Broad portion of tegmen extending only to about midwing; M with only one distal fork. EVANS, 1956. *Jur.*, Europe (Germany).




FIG. 149. Dymorphoptilidae and Ipsviciidae (p. 230–233).

Dymorphoptiloides EVANS, 1956, p. 218 [**D. elongata*; OD]. Tegmen as in *Dymorphoptila*, but broad portion extending nearer to apex; M with 2 distal forks. RIEK, 1974b. *Trias.*, Australia (Queensland), South Africa.—FIG. 149,9. **D. elongata*; tegmen, $\times 3.4$ (Evans, 1956).

Mesoatraxis BECKER-MIGDISOVA, 1949b, p. 40 [**M. reducta*; OD]. Tegmen as in *Dymorphoptiloides* but with shorter distal area; M with 3 terminal

branches. BECKER-MIGDISOVA, 1962b. *Jur.*, USSR (Tadzhik).

Tennentsia RIEK, 1976b, p. 813 [**Dymorphoptiloides protuberans* RIEK, 1974c, p. 22; OD]. Fore wing similar to that of *Dymorphoptiloides*, but SC with several distal branches and RS unbranched; M and CU apparently connected basally by a crossvein. *Trias.*, South Africa.—FIG. 149,10. **T. protuberans*; fore wing, $\times 2.3$ (Riek, 1976b).

Family CERCOPIDAE

Westwood, 1838

[Cercopidae WESTWOOD, 1838, p. 39]

Head narrower than pronotum, usually as wide as anterior margin of scutellum; ocelli on disc of crown, each at posterior end of sulcus; length and width of eyes almost equal; antennae originating in cavities below anterior margin of head; postclypeus commonly protuberant. Fore wings usually coriaceous. [The Aphrophoridae are included here, as a subfamily, because of the difficulty of recognizing the distinguishing features in the fossils.] EVANS, 1956. *Trias.*—*Holo.*

Cercopis FABRICIUS, 1775, p. 688. [Numerous extinct species from Tertiary deposits and described before 1900 were placed in the genus, but their assignment to *Cercopis* has not been generally accepted (see HANDLIRSCH, 1907, p. 1072–1074). However, a few, well-preserved specimens appear to justify at least tentative placement in the genus.] SCUDDER, 1890; COCKERELL, 1920a, 1927b; EVANS, 1956. *Eoc.*, Canada (British Columbia), USA (Colorado, Wyoming), USSR (Asian RSFSR)—*Holo.*

Alotritifidus EVANS, 1956, p. 216 [**A. interruptus*; OD]. Fore wing as in *Trifidella*, but costal margin arching basally and RS arising further distally. *Trias.*, Australia (Queensland).—FIG. 148, 10. **A. interruptus*; fore wing, $\times 10$ (Evans, 1956).

Aphrophora GERMAR, 1821, p. 48. COCKERELL, 1922f, 1925g; PONGRÁZ, 1928; PITON, 1936c; THÉOBALD, 1937a; BECKER-MIGDISOVA, 1964. *Eoc.*, Europe (Baltic, France), Canada (British Columbia); *Oligo.*, England, Europe (France); *Mio.*, USSR (Asian RSFSR)—*Holo.*

Cercopites SCUDDER, 1890, p. 316 [**C. calliscens* SCUDDER, 1890, p. 316; SD CARPENTER, herein]. Head relatively small; thorax hexagonal; fore wing more than twice as long as broad. *Eoc.*, USA (Wyoming), Canada (British Columbia).

Clastoptera GERMAR, 1839, p. 187. SCUDDER, 1890. *Oligo.*, USA (Colorado)—*Holo.*

Dawsonites SCUDDER, 1895b, p. 18 [**D. veter.*; OD]. Similar to *Palecphora*, but RS arising at mid-wing. *Mio.*, Canada (British Columbia).

Eocercopidium ZEUNER, 1944a, p. 116, *nom. subst. pro Eocercopis* ZEUNER, 1941a, p. 88, *non* HANDLIRSCH, 1939 [**Eocercopis maculata* ZEUNER, 1941a, p. 88; OD]. Fore wing similar to that of *Aphrophora*, but R strongly bent anteriorly near base; preradial part of wing very wide, crossed by pectinate branches from R; radial-median area very broad; M separating from CUA very near to base. *Eoc.*, Scotland.—FIG. 148, 4. **E.*

maculata (ZEUNER); fore wing, $\times 6.4$ (Zeuner, 1944a).

Megacercopis COCKERELL, 1925g, p. 9 [**M. optima*; OD]. Little-known fore wing with venation similar to that of *Stenecphora*, but apex much more pointed. *Mio.*, USSR (Asian RSFSR).—FIG. 148, 2. **M. optima*; fore wing, $\times 2.5$ (Cockerell, 1925g).

Palaeoptysma SCUDDER, 1895b, p. 21 [**P. venosa*; OD]. Little-known fore wing, related to *Aphrophora* but very slender. *Eoc.*, Canada (British Columbia).

Palaphrodes SCUDDER, 1890, p. 333 [**P. irregularis* SCUDDER, 1890, p. 333; SD CARPENTER, herein]. Fore wing as in *Cercopis*, but head very obtuse and rounded in front, narrower distally than thorax. COCKERELL, 1908k. *Oligo.*, USA (Colorado).

Palecphora SCUDDER, 1890, p. 324 [**P. communis* SCUDDER, 1890, p. 324; SD CARPENTER, herein]. Fore wing longer and more slender than that of *Palaphrodes*; costal margin less arched. COCKERELL, 1908k. *Oligo.*, USA (Colorado).

Petrolystra SCUDDER, 1878a, p. 530 [**P. gigantea* SCUDDER, 1878a, p. 530; SD CARPENTER, herein]. Large insects; head large, flat dorsally, twice as broad as long, the front broadly convex; scutellum very small, about half as long as thorax. SCUDDER, 1890. *Oligo.*, USA (Colorado).

Philagra STÅL, 1863, p. 593. COCKERELL, 1925g. *Mio.*, USSR (Asian RSFSR)—*Holo.*

Ptyelus LE PELETIER & SERVILLE, 1828, p. 608. THÉOBALD, 1937a. *Oligo.*, Europe (France)—*Holo.*

Ptysmaphora SCUDDER, 1895b, p. 21 [**P. fleischeri*; OD]. Fore wing as in *Palaeoptysma* but with costal margin straighter. *Eoc.*, Canada (British Columbia).

Sinophora MELICHAR, 1902, p. 113 [**S. maculosa*; OD]. BECKER-MIGDISOVA, 1964. *Mio.*, USSR (Asian RSFSR)—*Holo.*

Stenecphora SCUDDER, 1895b, p. 17 [**S. punctulata*; OD]. Fore wing with very broad apex, slender clavus; RS arising near base. *Eoc.*, Canada (British Columbia).

Stenolocris SCUDDER, 1895b, p. 19 [**S. venosa*; OD]. Little-known fore wing, with very strong costal vein and RS arising at wing base. [Family assignment doubtful.] *Mio.*, Canada (British Columbia).

Triassoscarta TILLYARD, 1919c, p. 874 [**T. subcostalis*; OD]. Little-known genus, based on incomplete tegmen. SC apparently absent; R long, nearly parallel with costal margin and connected to costal margin by about 8 subequal crossveins. [Originally placed in the *Scytinopteridae* but transferred to *Cercopidae* by EVANS (1956).] *Trias.*, Australia (Queensland).—FIG. 148, 7. **T. subcostalis*; fore wing, $\times 6$ (Evans, 1956).

Triecphora AMYOT & SERVILLE, 1843, p. 561. WOODWARD, 1879. *Eoc.*, England—*Holo.*

Trifidella EVANS, 1956, p. 215 [**T. perfecta*; OD]. Fore wing tegminous, coarsely rugose; several long veinlets between wing margin and R; M and CUA fused basally; CUA forked. *Trias.*, Australia (Queensland).—FIG. 148,6. **T. perfecta*; fore wing, $\times 10$ (Evans, 1961).

Family IPSVICIIDAE Tillyard, 1919

[Ipsviciidae TILLYARD, 1919c, p. 878] [=Stenoviciidae EVANS, 1956, p. 205]

Fore wing uniformly sclerotized; costal margin thick and flattened; vein R consisting usually of R and less commonly of RS; R joined to M by a prominent crossvein; M and CUA usually arising from a common basal stem; M typically branched; CUA and CUP apparently unbranched. Hind wing (known only in *Ipsvicia*) strongly curved anteriorly in distal area; CUA branched. Body unknown. *Perm.*—*Trias.*

The systematic position of this family is obscure. TILLYARD (1919c) originally assigned it to the Homoptera, close to the extinct family Syntonopteridae, but later (1926d) transferred it to the Fulgoroidea of the Homoptera. Subsequently, it has been placed in the Heteroptera by EVANS (1956), in the Homoptera (Auchenorrhyncha) by BECKER-MIGDISOVA (1962b), in the Homoptera (Peloridioidea) by CHINA (1962), and in the Homoptera (Cercopoidea) by EVANS (1963). Also, eight of the genera discussed below (*Stenovicia*, *Permocentrus*, *Permagra*, *Permonia*, *Stanleyana*, *Palaeovicicia*, *Apheloscya*, and *Permoscarta*) were placed in a new family, Stenoviciidae, by EVANS (1956), although most of these were previously assigned to the Ipsviciidae (EVANS, 1943b). BECKER-MIGDISOVA (1962b) concluded that the new family is unnecessary, and I have followed her treatment in retaining these genera in the Ipsviciidae.

Ipsvicia TILLYARD, 1919c, p. 878 [**I. jonesi*; OD]. R with several anterior branches to costa near middle of tegmen. TILLYARD, 1923b. *Trias.*, Australia (Queensland).—FIG. 149,2. **I. jonesi*; *a*, tegmen; *b*, hind wing, $\times 4$ (Evans, 1956).

Apheloscya TILLYARD, 1922b, p. 458 [**A. mesocampa*; OD]. Branches of all veins of tegmen very short. [Family assignment doubtful.] EVANS,

1956; BECKER-MIGDISOVA, 1962b. *Perm.*, Australia (New South Wales).

Ipsviciella BECKER-MIGDISOVA, 1962a, p. 100 [**I. asiatica*; OD]. Tegmen with rounded apex; R nearly straight, with several parallel branches to costal margin; CUA unbranched, merging with M basally. *Trias.*, USSR (Kirghiz).—FIG. 149,5. **I. asiatica*; tegmen, $\times 6.5$ (Becker-Migdisova, 1962b).

Ipsviciopsis TILLYARD, 1922b, p. 464 [**I. elegans*; OD]. RS separating from R near base of tegmen. EVANS, 1963. *Trias.*, Australia (Queensland).

Palaeovicicia EVANS, 1943b, p. 189 [**P. incerta*; OD]. Tegmen: RS short; M with 3 branches. EVANS, 1956; BECKER-MIGDISOVA, 1962b. *Perm.*, Australia (New South Wales).—FIG. 149,6. **P. incerta*; tegmen, $\times 8$ (Evans, 1943b).

Permagra EVANS, 1943a, p. 7 [**P. distincta*; OD]. Tegmen as in *Tomioscarta* but lacking closed cells. EVANS, 1956; BECKER-MIGDISOVA, 1962b. *Perm.*, Australia (New South Wales).

Permocentrus EVANS, 1956, p. 207 [**Permoscarta trivenulata* TILLYARD, 1926a, p. 19; OD]. Tegmen with M and CUA independent basally. BECKER-MIGDISOVA, 1962b. *Perm.*, Australia (New South Wales).

Permoscarta TILLYARD, 1918b, p. 726 [**P. Mitchellii*; OD]. Little-known genus. Tegmen as in *Permocentrus* but with 2 crossveins between M and CUA. EVANS, 1943a, 1956; BECKER-MIGDISOVA, 1962b. *Trias.*, Australia (Queensland).

Permovicicia EVANS, 1943b, p. 189 [**P. obscura*; OD]. Tegmen with RS broadly curved. EVANS, 1956. *Perm.*, Australia (New South Wales).—FIG. 149,3. **P. obscura*; $\times 10$ (Evans, 1943b).

Stanleyana EVANS, 1943b, p. 188 [**S. pulchra*; OD]. Tegmen with RS apparently absent; M and CUA coalesced basally; M with 3 branches. EVANS, 1956. *Perm.*, Australia (New South Wales).—FIG. 149,8. **S. pulchra*; tegmen, $\times 6.5$ (Evans, 1943b).

Stenovicia EVANS, 1943b, p. 188 [**S. angustata*; OD]. Tegmen as in *Ipsvicia* but much more slender; R long, arising at about midwing; M with 2 very short branches; CUA and M coalesced basally. [Type of family Stenoviciidae EVANS, 1956.] *Perm.*, Australia (New South Wales).—FIG. 149,1. **S. angustata*; fore wing, $\times 8$ (Evans, 1943b).

Tomioscarta BECKER-MIGDISOVA, 1961c, p. 350 [**T. surijokovensii*; OD]. Tegmen with R branched at point of origin of RS; several closed cells between M, CUA, and RS. BECKER-MIGDISOVA, 1962b. *Perm.*, USSR (Asian RSFSR).—FIG. 149,7. **T. surijokovensii*; fore wing, $\times 6.5$ (Becker-Migdisova, 1961c).

Tychoscarta BECKER-MIGDISOVA, 1961c, p. 350 [**T. sokolovensii*; OD]. Little-known genus.

Tegmen long and narrow; RS unbranched and continuing in a straight line from stem of R; M unbranched; CUA forked distally. BECKER-MIGDISOVA, 1962b. *Perm.*, USSR (Asian RSFSR).—FIG. 149,4. **T. sokolovensis*; fore wing, $\times 8$ (Becker-Migdisova, 1961c).

Family ARCHIJASSIDAE Becker-Migdisova, 1962

[Archijassidae BECKER-MIGDISOVA, 1962a, p. 95]

Fore wing very wide, in some species with triangular costal area traversed by vein SC; SC usually divided into 2 long branches; RS present; numerous crossveins between branches of R and M; anal area wide, triangular. *Jur.*

Archijassus HANDLIRSCH, 1906b, p. 501 [**Cercopidium beeri* GEINITZ, 1880, p. 529; SD CARPENTER, herein]. Fore wing with costal margin strongly angular; RS arising beyond midwing; M with 4 branches. EVANS, 1956. *Jur.*, Europe (Germany).—FIG. 148,8. **A. beeri* (GEINITZ); fore wing, $\times 8$ (Evans, 1956).

Atitizon HANDLIRSCH, 1939, p. 144 [**A. jassoides*; OD]. Fore wing very broad; costal margin strongly curved but not angular basally; RS arising at midwing. *Jur.*, Europe (Germany).—FIG. 148,11. **A. jassoides*; fore wing, $\times 8$ (Handlirsch, 1939).

Eojassus HANDLIRSCH, 1939, p. 145 [**E. indistinctus*; OD]. Little-known genus, based on fore wing; costal margin smoothly curved. *Jur.*, Europe (Germany).—FIG. 148,9. **E. indistinctus*; fore wing, $\times 6.5$ (Handlirsch, 1939).

Liojassus HANDLIRSCH, 1939, p. 146 [**L. affinis*; OD]. Fore wing; SC with 2 long branches; RS arising at midwing; costal margin smoothly curved; M with 3 branches. [Family assignment doubtful.] *Jur.*, Europe (Germany).—FIG. 148,1. **L. affinis*; fore wing, $\times 6.5$ (Handlirsch, 1939).

Family HYLICELLIDAE Evans, 1956

[Hylcellidae EVANS, 1956, p. 195]

Fore wing as in Hylcidae (recent), with M coalesced basally with CUA, but CUA1 present and coalesced with part of M3+4 distally. *Trias.*

Hylcella EVANS, 1956, p. 195 [**H. colorata*; OD] [= *Hylcellites* BECKER-MIGDISOVA, 1962a, p. 95, (type, *Hylcella reducta* EVANS)]. CUA with abrupt basal bend; 2 crossveins between RS and M1+2; 1 crossvein between M1+2 and M3+4. *Trias.*, Australia (Queensland).—FIG. 150,6. **H. colorata*; fore wing, $\times 5$ (Evans, 1956).

Family MUNDIDAE Becker-Migdisova, 1960

[Mundidae BECKER-MIGDISOVA, 1960, p. 31]

Fore wing weakly tegminous, without pits; veins thick; RS, M, and CUA with prominent projections; costal area and anal area broad. *Perm.*

Mundus BECKER-MIGDISOVA, 1960, p. 31 [**M. nodosus*; OD]. Fore wing relatively broad, with asymmetrical, blunt apex; R diverging abruptly at midwing toward costal margin, forking; R2 parallel to RS. *Perm.*, USSR (European RSFSR).—FIG. 150,5. **M. nodosus*; fore wing, $\times 8$ (Becker-Migdisova, 1960).

Family PEREBORIIDAE Zalessky, 1930

[*nom. correct.* BRUES, MELANDER, & CARPENTER, 1954, p. 813 (pro Pereboridae ZALESSKY, 1930, p. 1026)] [= Permoglyphidae EVANS, 1943b, p. 183]

Fore wing membranous; veins R, RS, and CUA with extensive branching. BECKER-MIGDISOVA, 1962b. *Perm.*—*Trias.*

Pereboria ZALESSKY, 1930, p. 1021 [**P. bella*; OD]. Little-known genus, based on fore wing. R with close pectinate branching; crossveins numerous, irregular; wing large, about 40 mm long. EVANS, 1956; BECKER-MIGDISOVA, 1962b. *Perm.*, USSR (Asian RSFSR).—FIG. 150,9. **P. bella*; fore wing, $\times 1.5$ (Becker-Migdisova, 1962b).

Crosbella EVANS, 1956, p. 192 [**C. elongata*; OD]. Fore wing as in *Permobrachus*, but M more extensively branched. *Trias.*, Australia (Queensland).—FIG. 150,1. **C. elongata*; fore wing, $\times 4.5$ (Evans, 1956).

Kaltanopibrocha BECKER-MIGDISOVA, 1961c, p. 357 [**K. boreoscytinoides*; OD]. Little-known genus, based on hind wing fragment. Costal margin almost straight; R directed posteriorly in apical region, pectinately branched; M forking before RS. [Family assignment doubtful.] *Perm.*, USSR (Asian RSFSR).—FIG. 150,10. **K. boreoscytinoides*; hind wing, $\times 4.5$ (Becker-Migdisova, 1961c).

Neuropibrocha BECKER-MIGDISOVA, 1961c, p. 356 [**N. ramisubcostalis*; OD]. Fore wing as in *Pereboria*, but R with fewer pectinate branches and less dense reticulation of branches of RS, M, and CUA; area between stems R and M with few crossveins. *Perm.*, USSR (Asian RSFSR).—FIG. 150,7. **N. ramisubcostalis*; fore wing, $\times 2.0$ (Becker-Migdisova, 1961c).

Permobrachus EVANS, 1943b, p. 183 [**Permopdipthera dubia* TILLYARD, 1926a, p. 24; OD]. Fore wing shaped as in *Scytophara*, but R1 curv-




FIG. 150. Hylicellidae, Mundidae, and Pereboriidae (p. 233–235).

ing abruptly to anterior margin; branching of R2 pectinate; M branching well beyond midwing. *Perm.*, Australia (New South Wales).—FIG. 150.4. *P. magnus* EVANS; fore wing, $\times 3.5$ (Evans, 1943b).

Permoglyphis TILLYARD, 1926a, p. 22 [**P. belmontensis*; OD]. Little-known genus, based on fore wing; similar to *Permopibrocha* but apparently with less branching of R, RS, and M; costal margin nearly straight. *Trias.*, Australia (New

South Wales).—FIG. 150.8. **P. belmontensis*; fore wing, $\times 4.5$ (Evans, 1956).

Permopibrocha MARTYNOV, 1935c, p. 18 [**P. ramosa*; OD]. Fore wing as in *Pereboria*, but R with fewer branches; M more deeply forked than CUA; fore wing small. *Perm.*, USSR (European RSFSR).—FIG. 150.3. **P. ramosa*; fore wing, $\times 3.5$ (Martynov, 1935c).

Scytophara MARTYNOV, 1937b, p. 36 [**S. extensa*; OD]. Fore wing more slender than in *Permopi-*

brocha; costal margin straight beyond base; M forking at about level of origin of RS. *Perm.*, USSR (European RSFSR).—FIG. 150,2. **S. extensa*; fore wing, $\times 6.5$ (Martynov, 1937b).

Family FULGORIDIIDAE

Handlirsch, 1939

[*nom. transl.* BECKER-MIGDISOVA, 1962b, p. 184, ex Fulgoridiinae HANDLIRSCH, 1939, p. 122]

Fore wing tegminous; costal margin only slightly arched; vein SC long, without branches; RS arising at about midwing; CUA forking well before origin of RS; crossveins few. Hind wing a little shorter than fore wing; anal area very broad; RS simple or with short fork; 1A arched away from CUP. EVANS, 1956; BECKER-MIGDISOVA, 1962b. *Jur.*

Fulgoridium HANDLIRSCH, 1906b, p. 496 [**Pbyrganidium balticum* GEINITZ, 1880, p. 527; OD] [= *Fulgoridulum* HANDLIRSCH, 1939, p. 140 (type, *F. egens*)]. Fore wing slender; usually with maculations; SC close to margin; R with a series of short branches distally; CUA with several long branches. BODE, 1953; EVANS, 1956. *Jur.*, Europe (Germany).—FIG. 151,2a. *F. punctatum* HANDLIRSCH; fore wing, $\times 10$ (Handlirsch, 1939).—FIG. 151,2b. *F. reductum* HANDLIRSCH; hind wing, $\times 10$ (Handlirsch, 1939).

Metafulgoridium CARPENTER, herein [**M. spilotum* HANDLIRSCH, 1939, p. 139; OD]. Fore wing as in *Fulgoridium*, but CUA2 unbranched. [The original generic name, *Metafulgoridium*, was a *nomen nudum* (HANDLIRSCH, 1939).] *Jur.*, Europe (Germany).—FIG. 151,1. **M. spilotum*; fore wing, $\times 6.5$ (Handlirsch, 1939).

Family LOPHOPIDAE Stål, 1866

[Lophopidae Stål, 1866, p. 130]

Head markedly narrower than pronotum; vertex usually narrow; pronotum short and broad, tricarinate. Fore wing coriaceous, with conspicuous venation and supernumerary longitudinal veins and crossveins; wing usually elongate; apical margin broadly rounded; claval veins united before apex. Fore and middle tibiae usually compressed. *Jur.*—*Holo.*

Lophops SPINOLA, 1838, p. 205. *Holo.*

Eofulgoridium MARTYNOV, 1937a, p. 164 [**E. kirsylkiense*; OD]. Fore wing with SC about midway between C and R; M dividing at midwing; M with 3 branches. Hind wing little known; costal margin concave; RS arising beyond midwing; M and CUA dividing beyond midwing. EVANS, 1956; BECKER-MIGDISOVA, 1962b. *Jur.*, USSR




FIG. 151. Fulgoridiidae and Lophopidae (p. 235).

(Kirghiz).—FIG. 151,3a. **E. kirsylkiense*; fore wing, $\times 4.2$.—FIG. 151,3b. *E. proximum* MARTYNOV; hind wing, $\times 5$ (Martynov, 1937a). **Scoparidea** COCKERELL, 1920c, p. 243 [**S. nebulosa*; OD]. Fore wing with RS parallel to R; apical region with numerous, parallel veins; no regular gradate series of veins. [Family assignment doubtful.] *Eoc.*, USA (Colorado).

Family CIXIIDAE Spinola, 1838

[Cixiidae SPINOLA, 1838, p. 204]

Head not elongate; antennae with 2 segments, bearing flagella; wings unusually well developed. In fore wing, veins SC, R, and M with common stem; claval suture distinct; claval veins united into a claval stem. *Perm.*—*Holo.*

- Cixius** LATREILLE, 1804, p. 168. [The assignment of a Jurassic species from England to this genus is very doubtful (FENNAH, 1961), as is that of the several species in Baltic amber (GERMAR & BERENDT, 1856).] SCUDDER, 1890. *Eoc.*, USA (Wyoming); *Oligo.*, USA (Colorado)—*Holo.*
- Asiocixius** BECKER-MIGDISOVA, 1962a, p. 97 [**A. fulgoroides*; OD]. Fore wing membranous, except at base; costal margin smoothly rounded; R2 curved toward RS and giving rise to several veinlets; RS forked distally; M forking beyond midwing and with extensive pectinate branching; CUA with a long fork. *Trias.*, USSR (Kirghiz).—FIG. 152,6. **A. fulgoroides*; fore wing, $\times 5$ (Becker-Migdisova, 1962b).
- Boreocixius** BECKER-MIGDISOVA, 1955, p. 1100 [**B. sibiricus*; OD]. Fore wing with costal margin strongly thickened; RS arising very near wing base; R and RS with very short branches; fork of CUA long and curved. *Trias.*, USSR (Asian RSFSR).—FIG. 152,12. **B. sibiricus*; fore wing, $\times 10$ (Becker-Migdisova, 1962b).
- Cixiella** BECKER-MIGDISOVA, 1962a, p. 98 [**C. reducta*; OD]. Fore wing weakly tegminous, distal portion membranous; RS arising near midwing; M forking beyond level of origin of RS, with 3 terminal branches, and forming a large, closed cell; CUA curved basally. *Trias.*, USSR (Kirghiz).—FIG. 152,8. **C. reducta*; fore wing, $\times 10$ (Becker-Migdisova, 1962b).
- Cyclocyrtina** MARTYNOV, 1926b, p. 1349 [**C. delutinervis*; OD]. Fore wing tegminous, elongate; costal margin only slightly curved; R with a series of branches as in *Mesocixiella* but shorter; M joined to RS distally by a recurved branch. Hind wing little known; M with 2 long branches, arising before midwing. EVANS, 1956. *Trias.*, USSR (Kirghiz); *Jur.*, USSR (Kazakh, Tadzhik).—FIG. 152,3. **C. delutinervis*, *Jur.*, Kazakh; fore wing, $\times 6$ (Becker-Migdisova, 1962b).
- Diaplegma** SCUDDER, 1890, p. 288 [**D. abductum* SCUDDER, 1890, p. 290; SD CARPENTER, herein]. Similar to *Cixius*, but RS arising near midwing, each of its forks dividing into 2 or 3 distal, curved branches. *Oligo.*, USA (Colorado).
- Eofulgorella** COCKERELL, 1909j, p. 172 [**E. bradburyi*; OD]. Fore wing resembling that in *Oliarius* but elongate and with costal margin concave; crossveins forming a very regular series. [Family assignment doubtful.] *Eoc.*, USA (Colorado).
- Eoliarius** COCKERELL, 1925a, p. 10 [**E. quadri-stictus*; OD]. Similar to *Oliarius*, but RS arising well before the pterostigmal area and giving rise to 4 very oblique branches anteriorly. *Eoc.*, USA (Colorado).
- Hyaesthes** SIGNORET, 1865, p. 128 [**H. obsoletus*; OD]. STATZ, 1950a. *Oligo.*, Europe (Germany)—*Holo.*
- Mesocixiella** BECKER-MIGDISOVA, 1949b, p. 38 [**M. asiatica*; OD]. Fore wing with costal margin only slightly curved; R with a series of parallel branches leading to margin; RS arising before midwing with 3 or 4 terminal branches; M forked beyond midwing. EVANS, 1956. *Trias.*, USSR (Kirghiz); *Jur.*, USSR (Kazakh).—FIG. 152,7. **M. asiatica*; fore wing, $\times 6.5$ (Becker-Migdisova, 1962b).
- Mesocixius** TILLYARD, 1919c, p. 876 [**M. trias-sicus*; OD]. Fore wing with RS forking about halfway between origin of RS and wing apex; fork of M less distal. EVANS, 1956. *Trias.*, Australia (Queensland).—FIG. 152,10. **M. trias-sicus*; fore wing, $\times 5.4$ (Tillyard, 1919c).
- Mundopoides** COCKERELL, 1925g, p. 11 [**M. cisthenaria*; OD]. Similar to *Mundopa* (recent), having nearly straight costal and outer margins, the apex being obliquely truncate; SC terminating at midwing. *Mio.*, USSR (Asian RSFSR).
- Myndus** STÅL, 1862, p. 307. COCKERELL, 1926b. *Oligo.*, England—*Holo.*
- Oeclixius** FENNAH, 1963, p. 43 [**O. amphion*; OD]. Similar to *Oecleus* (recent) but with long, slender tibiae; pterostigma only moderately developed; tegminal veins distinctly granulate. *Mio.*, Mexico (Chiapas).—FIG. 152,5. **O. amphion*; fore wing, $\times 13$ (Fennah, 1963).
- Oliarites** SCUDDER, 1890, p. 293 [**Mnemosyne terrentula* SCUDDER, 1878b, p. 773; OD]. Little-known genus, with head less than half as broad as thorax; veins forming a weak reticulation distally. [Family assignment doubtful.] *Eoc.*, USA (Wyoming).
- Oliarius** STÅL, 1862, p. 306. COCKERELL, 1910b. *Oligo.*, Europe (Baltic)—*Holo.*
- Oligonila** CARPENTER, herein [**O. defectuosa* THÉOBALD, 1937a, p. 258; OD]. Fore wing as in *Anila* (recent) but lacking the oblique vein in the costal area. [The original generic name, *Oligonila*, was a *nomen nudum* (THÉOBALD, 1937a).] *Oligo.*, Europe (France).
- Permcixiella** BECKER-MIGDISOVA, 1961c, p. 361 [**P. venosa*; OD]. Fore wing elongate, costal margin nearly straight; R2 straight; branches of CUA nearly straight. *Perm.*, USSR (Asian RSFSR).—FIG. 152,4. **P. venosa*; fore wing, $\times 5.4$ (Becker-Migdisova, 1961c).
- Protoliarius** COCKERELL, 1920c, p. 243 [**P. hamatus*; OD]. Similar to *Oliarius* but without a stigmatic spot on wings. COCKERELL, 1924a; COCKERELL & LEVEQUE, 1931. *Eoc.*, USA (Colorado).
- Scytocixius** MARTYNOV, 1937b, p. 34 [**S. mendax*; OD]. Fore wing broader distally than basally; costal margin smoothly curved; R2 strongly arched away from margin; RS similarly arched but less strongly; M with 3 distal branches; CUA forking at the level of origin of RS. *Perm.*, USSR (Asian RSFSR).—FIG. 152,1. **S. mendax*; fore wing, $\times 10$ (Becker-Migdisova, 1962b).
- Suriokocixius** BECKER-MIGDISOVA, 1961c, p. 359




FIG. 152. Cixiidae (p. 236-238).

[**S. tomiensis*; OD]. Fore wing broad, with broadly rounded apex; costal margin thickened; R2 strongly curved; RS unbranched; branches of CUA long and curved. EVANS, 1956. *Perm.*, USSR (Asian RSFSR).—FIG. 152,11. **S. tomiensis*; fore wing, $\times 15$ (Becker-Migdisova, 1961c). **Triassocixius** TILLYARD, 1919c, p. 878 [**T. aus-*

tralicus; OD]. Little-known genus, based on fragment of fore wing; R forked close to the origin of RS; oblique crossveins from R to costal margin. [Family position uncertain.] *Trias.*, Australia (Queensland).—FIG. 152,2. **T. australicus*; fore wing, $\times 5.5$ (Evans, 1956). **Vitreacixius** BECKER-MIGDISOVA, 1962a, p. 99 [**V.*

ellipticus; OD]. Fore wing weakly tegminous; similar to *Cixiella*, but RS arising slightly more basally; M with 4 branches, closed cell smaller than in *Cixiella*. *Trias.*, USSR (Kirghiz).—FIG. 152,9. **V. ellipticus*; fore wing, $\times 6$ (Becker-Migdisova, 1962b).

Family ACHILIDAE Stål, 1866

[Achilidae STÅL, 1866, p. 130]

Head usually small; frons and clypeus large. Hind tibiae elongate; second segment of hind tarsus large. Fore wing well developed, basal two-thirds thickened; veins SC and R united for a short interval basally; SC with 2 or more short branches leading to costal margin, forming stigmatic area; R branched only apically, connected to M by 2 or more crossveins; M with at least 3 branches; clavus short, claval veins united to form claval stem. Hind wing moderately large. *Oligo.*—*Holo.*

Achilus KIRBY, 1819, p. 474. *Holo.*

Elidiptera SPINOLA, 1839, p. 304. SCUDDER, 1890. *Oligo.*, USA (Colorado)—*Holo.*

Proteiptera USINGER, 1939, p. 66 [**P. kaweckii*; OD]. Similar to *Eiptera* (recent) but with vertex distinctly in front of eyes; posterior margin of vertex concavely arcuate. *Oligo.*, Europe (Baltic).

Family RICANIIDAE

Amyot & Serville, 1843

[Ricaniidae AMYOT & SERVILLE, 1843, p. 527]

Head usually as wide as the pronotum; vertex short and broad; clypeus much narrower than frons. Fore wing large, broadly triangular; costal margin usually nearly straight; costal area broad with numerous crossveins; basal area of clavus without pustules; venation diverse; veins R, M, and CU typically with numerous branches, with 1 or 2 subapical lines of gradate crossveins. Hind wing smaller than fore wing and with reduced venation. Basal segment of hind tarsus very small, without lateral spines. *Trias.*—*Holo.*

Ricania GERMAR, 1818, p. 221. DALMAN, 1826; GIEBEL, 1862; SCUDDER, 1890. *Eoc.*, Canada (British Columbia); *Oligo.*, Europe (Baltic)—*Holo.* *Cotradechites* FENNAH, 1968, p. 144 [**C. litbinus*; OD]. Similar to *Cotrades* (recent), but tegmen twice as long as broad; costal area broad, with dense venation. *Paleoc.*, USA (North Dakota).

Dilaropsis COCKERELL, 1920c, p. 244 [**D. ornatus*; OD]. Fore wing broad, triangular; costal margin slightly convex; SC ending about two-thirds wing length from base; M diverging abruptly from R near origin of RS. *Eoc.*, USA (Colorado).

Eobladina HAAPT, 1956, p. 13 [**E. antiqua*; OD]. Little-known genus, based on fore wing; costal area wide distally; SC joined to R at base by curved crossvein, forming a very short basal cell; RS arising well before midwing. *Eoc.*, Europe (Germany).—FIG. 153,2. **E. antiqua*; fore wing, $\times 6$ (Haupt, 1956).

Eoricania HENRIKSEN, 1922b, p. 24 [**E. danica*; OD]. Fore wing as in *Ricania* (recent), but 1A and 2A joined proximally beyond wing base. *Eoc.*, Europe (Denmark).—FIG. 153,4. **E. danica*; fore wing, $\times 2.5$ (Henriksen, 1922b).

Hammapteryx SCUDDER, 1890, p. 298 [**H. reticulata*; OD]. Fore wing subtriangular; costal margin arched at base; numerous crossveins from SC to margin; R with at least 2 arcuate branches distally; RS arising well before midwing. COCKERELL, 1920a, 1920b; COCKERELL & SANDHOUSE, 1921; HENRIKSEN, 1922b; PITON, 1940a. *Eoc.*, USA (Colorado, Wyoming), Europe (Denmark, France), England.—FIG. 153,3. *H. paucistriata* HENRIKSEN, Denmark; fore wing, $\times 4$ (Henriksen, 1922b).

Ludibrium BECKER-MIGDISOVA, 1962a, p. 100 [**L. ludus*; OD]. Hind wing little known; RS apparently arising distally as a continuation of stem R; M forked to about midwing. *Trias.*, USSR (Kirghiz).—FIG. 153,5. **L. ludus*; hind wing, $\times 6$ (Becker-Migdisova, 1962a).

Neoricania CARPENTER, 1990, p. 131, *nom. subst. pro Eoricania* HAAPT, 1956, p. 12, *non* HENRIKSEN, 1922b [**Eoricania reticulata* HAAPT; OD]. Fore wing with costal space much narrower than in *Eoricania*; SC much closer to C. *Eoc.*, Europe (Germany).

Scolypopites TILLYARD, 1923a, p. 17 [**S. bryani*; OD]. Fore wing as in *Scolypopa* (recent), but SC shorter, reaching only to a little beyond midwing; only one series of gradate veins. *Mio.*, Australia (Queensland).—FIG. 153,6. **S. bryani*; fore wing, $\times 3.5$ (Tillyard, 1923a).

Family NOGODINIDAE

Melichar, 1898

[Nogodinidae MELICHAR, 1898, p. 204]

Head about as wide as pronotum; frons longer than wide. Fore wing large, usually broadest towards apex, coriaceous or hyaline, with numerous veins and crossveins; costal area with several crossveins; basal cell usually large; clavus not punctulate; claval stem reaching apex of fore wing. Hind tibiae with

lateral spines; second segment of hind tarsus small, with a pair of spines distally. *Eoc.*—*Holo.*

Nogodina STÅL, 1859, p. 326. *Holo.*

Detyopsis COCKERELL, 1920c, p. 242 [**D. scud-deri*; OD]. Fore wing much as in *Detya* (recent); veinlets from SC to costal margin numerous; RS forking well before midwing. *Eoc.*, USA (Colorado).

Tritophania JACOBI, 1937, p. 188 [**T. patruelis*; OD]. Similar to *Gaetulia* (recent), but frons without a keel; pterostigma absent. *Oligo.*, Europe (Baltic).—FIG. 153, 1. **T. patruelis*; whole insect, $\times 3.4$ (Jacobi, 1937).

Family FULGORIDAE Latreille, 1807

[Fulgoridae LATREILLE, 1807, p. 163]

Head usually large and simple, but often with prominent, cephalic process; postclypeus large, triangular; compound eyes large. Fore wing well developed, with numerous supernumerary veins and crossveins; hind wing with the anal and jugal areas reticulate. *Eoc.*—*Holo.*

Fulgora LINNÉ, 1767, p. 703. *Holo.*

Callospilopteron COCKERELL, 1920c, p. 245 [**C. ocellatum*; OD]. Fore wing broad, with obtuse apex; costal area much reduced; SC short; anterior veinlets from SC and R very oblique; ocelliform spots near outer margin. [Family assignment doubtful.] *Eoc.*, USA (Wyoming).

Eucophora SPINOLA, 1839, p. 200. SCUDDER, 1895b. *Eoc.*, Canada (British Columbia)—*Holo.*

Lystra FABRICIUS, 1803, p. 56. SCUDDER, 1890. [Generic assignment of fossil doubtful.] *Eoc.*, USA (Wyoming)—*Holo.*


Nyktalos METCALF, 1952, p. 230, *nom. subst. pro Nyctophylax* SCUDDER, 1890, p. 279, *non FITZINGER*, 1860 [**Nyctophylax ubleri* SCUDDER; OD]. Large species of uncertain affinities; head with a stout, recurved process; legs stout; femora and tibiae carinate. *Oligo.*, USA (Colorado).

Poiocera LAPORTE, 1832, p. 221. GERMAR & BERENDT, 1856. *Oligo.*, Europe (Baltic)—*Holo.*


Family FLATIDAE Spinola, 1838

[Flatidae SPINOLA, 1838, p. 205]


Head narrower than thorax; lateral edges of face not angular. Fore wing with costal area having crossveins; basal area of clavus granulate; clavus often open, claval veins separate or joined apically. Hind tibiae without a movable spur; first hind tarsomere short,




1
Tritophania




2
Eobladina




3
Hammapteryx



4
Eoricania



5
Ludibrum



6
Scolyopites

FIG. 153. Ricaniidae and Nogodinidae (p. 238–239).

second one small with a spine on each side. *Eoc.*—*Holo.*

Flata FABRICIUS, 1798, p. 511. *Holo.*

Aphaena GUÉRIN & MÉNEVILLE, 1833, p. 452. SCUDDER, 1890; COCKERELL, 1920c. *Eoc.*, USA (Wyoming); *Oligo.*, USA (Colorado)—*Holo.*




FIG. 154. Issidae and Dictyopharidae (p. 240–241).

Ficarasites SCUDDER, 1890, p. 301 [**F. stigmatiscum*; OD]. Little-known genus; costal area narrow, with oblique veinlets; few crossveins. *Eoc.*, USA (Wyoming).

Giselia HAUPT, 1956, p. 14 [**G. multifurcata*; OD]. Fore wing as in *Uxantis* (recent); SC curved away from margin as it approaches midwing; R and M fused basally, separating early, with RS arising about one-sixth wing length from base; CUA apparently with a deep fork. *Eoc.*, Europe (Germany).

Lechaea STÅL, 1866, p. 236. HENRIKSEN, 1922b. *Eoc.*, Europe (Denmark)—*Holo.*

Ormenis STÅL, 1862, p. 68. COCKERELL, 1926a. *Tert.* (epoch unknown), Argentina. *Holo.*

Poekilloptera LATREILLE, 1796, p. 90. COCKERELL, 1921d. *Oligo.*, England—*Holo.*

Thaumastocladus COCKERELL & SANDHOUSE, 1921, p. 456 [**T. simplex*; OD]. Fore wing as in *Gaga* (recent); costal area broad, with numerous oblique veinlets; R branching apically; M and CUA coalesced to about midwing; CUP distinctly forked. [Family position doubtful.] COCKERELL, 1924a. *Eoc.*, USA (Wyoming).

Family ARAEOPIDAE Metcalf, 1938

[*Araeopidae* METCALF, 1938, p. 281]

Head usually small; antennae short, usually not longer than head and thorax combined. Fore wing diverse, ranging from brachypterous, with reduced venation, to fully developed, with normal venation; vein SC typically with 2 branches; R coalesced with SC for about half its length, then coalesced with part of M; M usually with 3 branches; CU with 3 branches. Hind wing usually present, sometimes reduced; SC and R coalesced for more than half their lengths; M

unbranched. Hind femora and tibiae elongate; spur well developed at apex of tibia, either spinelike or much enlarged and complex. *Eoc.*—*Holo.*

Araeopus SPINOLA, 1839, p. 336. COCKERELL, 1924a; STATZ, 1950a. *Eoc.*, USA (Colorado); *Oligo.*, Europe (Germany)—*Holo.*

Amagua COCKERELL, 1924a, p. 3 [**A. fortis*; OD]. Fore wing as in *Stenocranus* (recent); wing of uniform width, narrow; crossveins m-cu long. *Mio.*, USSR (Asian RSFSR).

Chloriona FIEBER, 1866, p. 519. BECKER-MIGDISOVA, 1964. *Mio.*, USSR (Asian RSFSR)—*Holo.*

Liburnia STÅL, 1866, p. 179. COCKERELL, 1917h. *Mio.*, Burma—*Holo.*

Family ISSIDAE Spinola, 1838

[*Issidae* SPINOLA, 1838, p. 158]

Head usually at least as wide as thorax; lateral margins of thorax not keeled; anterior margin of pronotum rounded and extended. Fore wing usually with reduced venation and often small; costal area small, without crossveins, or absent; base of costal margin not strongly curved; clavus not granulate. Hind tibiae with 2 to 4 spines; second hind tarsomere with a spine on each side. *Jur.*—*Holo.*

Issus FABRICIUS, 1803, p. 99. BERVOETS, 1910. *Oligo.*, Europe (Baltic)—*Holo.*

Elasmocelidium MARTYNOV, 1926b, p. 1355 [**E. rotundatum*; OD]. Fore wing short, much broadened distally; SC nearly parallel to costal margin; costal margin thickened; RS arising well before midwing; RS and M forked distally; anal area extending only to about midwing. BODE, 1953; EVANS, 1956; BECKER-MIGDISOVA, 1962b. *Jur.*, USSR (Kazakh); Europe (Germany).—FIG. 154, 1. **E. rotundatum*, Kazakh; fore wing, $\times 6.3$ (Becker-Migdisova, 1962b).

Issites HAUPT, 1956, p. 16 [**I. glaber*; OD]. Fore wing as in *Issus* (recent) but without the dense reticulation. *Eoc.*, Europe (Germany).

Mesotubilistrum BECKER-MIGDISOVA, 1949b, p. 35 [**M. asiaticum*; OD]. Similar to *Elasmocelidium*, but RS arising near midwing. *Jur.*, USSR (Kazakh).

Tetragonidium BODE, 1953, p. 194 [**T. paralelogramma*; OD]. Fore wing as in *Elasmocelidium*, but M with more branches. *Jur.*, Europe (Germany).

Family DICTYOPHARIDAE Spinola, 1838

[*Dictyopharidae* SPINOLA, 1838, p. 202]

Head relatively large; structural details of vertex and of frons diverse. Legs usually slender

and elongate; hind tibiae commonly with 3 to 5 stout spines; second hind tarsal segment large, with a row of small spines at apex. Fore wing either normal or reduced; vein SC and R coalesced beyond basal area of wing; R branching irregularly distally; an irregular transverse line commonly formed by series of crossveins in apical third of wing. Hind wing usually large, with irregular venation. EMELJANOV, 1983. *Cret.—Holo.*

Dictyophara GERMAR, 1833, p. 175. [The family assignment of "*Dictyophara scudderi* PITON (1940a), from the Eocene of France, is uncertain.] SCUDDER, 1890; BECKER-MIGDISOVA, 1964; EMELJANOV, 1983. *Mio.*, USSR (Asian RSFSR)—*Holo.*

Chanithus AMYOT, 1847, p. 160. BECKER-MIGDISOVA, 1964; EMELJANOV, 1983. *Mio.*, USSR (Asian RSFSR)—*Holo.*

Florissantia SCUDDER, 1890, p. 293 [**F. elegans*; OD]. Little-known genus, apparently related to *Dictyophara*. [Originally placed in Cixiidae by SCUDDER (1890); transferred to Dictyopharidae by EMELJANOV (1983).] COCKERELL, 1909a. *Oligo.*, USA (Colorado).

Netutela EMELJANOV, 1983, p. 84 [**N. annunciator*; OD]. Similar to *Cladodiptera* (recent), but clavus of fore wing without crossveins; M forking distally of origin of RS. *Cret.*, USSR (Asian RSFSR).—FIG. 154,2. **N. annunciator*; fore wing, $\times 6.5$ (Emeljanov, 1983).

Family ARCHESCYTINIDAE Tillyard, 1926

[Archescytinidae TILLYARD, 1926g, p. 385] [=Permopsyllidae TILLYARD, 1926g, p. 390; Lithoscytinidae CARPENTER, 1933a, p. 436; Maueriidae ZALESSKY, 1937e, p. 54; Permopsycynopsidae ZALESSKY, 1939, p. 36; Uraloscytinidae ZALESSKY, 1939, p. 40; Maripsoctidae ZALESSKY, 1939, p. 44; Kaltanaphididae SZELEGIEWICZ, 1971, p. 63]

Fore and hind wings membranous, similar in size and almost alike in venation. Fore wing with vein SC very close and parallel to R+M, R, and R1; R forming a pterostigma; RS originating at about midwing; M usually with at least 3 branches; CUA arising from stem CU, then directed towards R+M, which it touches at the point of separation of M; CUA forked; anal area small. Hind wing similar to fore wing except that CUA arises as an independent vein from the wing base and is not directed towards R+M. Head hypognathous; beak long; antennae long, multisegmented; ovipositor prominent in some genera at least. SZELEGIEWICZ & POPOV, 1978. *Perm.*

Archescytina TILLYARD, 1926g, p. 385 [**A. permiana*; OD] [=*Maueria* ZALESSKY, 1937e, p. 54 (type, *M. sylvensis*); *Permopsycynopsis* ZALESSKY, 1939, p. 36 (type, *P. maueriaeformis*)]. Fore wing with costal margin nearly straight except near base; SC close and parallel to R; R+M arched anteriorly; R2 parallel to RS; M usually with 3 branches. Antennae long and slender, with about 25 segments; beak long; forelegs with thickened femora; female with long, retractible ovipositor. CARPENTER, 1931b, 1939; ZALESSKY, 1937e, 1939; BECKER-MIGDISOVA, 1961c, 1961d, 1962b. *Perm.*, USA (Kansas), USSR (European and Asian RSFSR).—FIG. 155,1a. *Archescytina* sp., USSR; lateral view of body, $\times 6$ (Becker-Migdisova, 1961d).—FIG. 155,1b,c. **A. permiana*, Kansas; b, fore wing; c, hind wing, $\times 6.5$ (Carpenter, 1939).

Bekkerscytina EVANS, 1958, p. 111 [**B. primitiva*; OD]. Similar to *Eoscytina*, but RS arising nearer to origin of M. *Perm.*, Australia (New South Wales).—FIG. 155,10. **B. primitiva*; fore wing, $\times 6.3$ (Evans, 1958).

Eoscytina EVANS, 1958, p. 109 [**E. migdisovae*; OD]. Similar to *Archescytina*, but fork of CUA very deep and broad and stem of CUA, as it leaves CUP, sigmoidally curved. *Perm.*, Australia (New South Wales).—FIG. 155,9. **E. migdisovae*; fore wing, $\times 6$ (Evans, 1958).

Kaltanaphis BECKER-MIGDISOVA, 1959a, p. 107 [**K. permiansis*; OD]. Little-known genus, based on fragment of hind wing. [Originally assigned to Permaphidopseidae; placed in new family, Kaltanaphididae, by SZELEGIEWICZ, 1971; transferred to Archescytinidae by SZELEGIEWICZ & POPOV, 1978.] *Perm.*, USSR (Asian RSFSR).

Kaltanoscytina BECKER-MIGDISOVA, 1959a, p. 105 [**K. nigra*; OD]. Wings as in *Archescytina*, but R longer and straighter in both pairs. BECKER-MIGDISOVA, 1961c; SZELEGIEWICZ & POPOV, 1978. *Perm.*, USSR (Asian RSFSR).—FIG. 155,8. **K. nigra*; fore wing, $\times 7$ (Becker-Migdisova, 1961c).

Maripsocus ZALESSKY, 1939, p. 44 [**M. ambiguus*; OD]. Little-known fore wing; venation as in *Archescytina*, but M apparently with 2 branches. EVANS, 1956. *Perm.*, USSR (European RSFSR).

Paleoscytina CARPENTER, 1931b, p. 118 [**P. brevistigma*; OD]. Similar to *Archescytina*, but CUA of fore wing unbranched. BECKER-MIGDISOVA, 1961c. *Perm.*, USA (Kansas), USSR (Asian RSFSR).—FIG. 155,3. **P. brevistigma*; fore wing, $\times 18$ (Carpenter, 1933a).

Permopsylla TILLYARD, 1926g, p. 390 [**P. americana*; OD] [=*Lithoscytina* CARPENTER, 1933a, p. 436 (type, *L. cubitalis*)]. Fore wing as in *Archescytina* but relatively broader; costal margin slightly concave at level of origin of M. BECKER-MIGDISOVA, 1960, 1961c, 1962b. *Perm.*, USA (Kansas), USSR (European and Asian RSFSR).—FIG. 155,7. **P. americana*; fore wing, $\times 16$ (Carpenter, 1931b).




FIG. 155. Archescytinidae (p. 241–243).

Permopsyllopsis ZALESSKY, 1939, p. 38 [**P. rosica*; OD]. Little-known fore wing; venation as in *Archescytina*, but RS straight. BECKER-MIGDISOVA, 1960. *Perm.*, USSR (Asian RSFSR).
Permoscytina TILLYARD, 1926g, p. 387 [**P. kansasensis*; OD]. Similar to *Archescytina*, but SC and R nearly straight basally; proximal branch

of M arising at about level of origin of RS. CARPENTER, 1939. *Perm.*, USA (Kansas). — FIG. 155,2. **P. kansasensis*; fore wing, $\times 4.2$ (Carpenter, 1939).

Sarbaloscytina BECKER-MIGDISOVA, 1959a, p. 104 [**S. angustipennis*; OD]. Similar to *Archescytina*, but stem R+M short and nearly straight.

BECKER-MIGDISOVA, 1961c. *Perm.*, USSR (Asian RSFSR).—FIG. 155,5. **S. angustipennis*; fore wing, $\times 4.5$ (Becker-Migdisova, 1961c).

Sojanoscytina MARTYNOV, 1933c, p. 885 [**S. grandis*; OD] [= *Ivascytina* MARTYNOV, 1933c, p. 888 (type, *I. difficilis*). Fore wing similar to that of *Archescytina*, but M with 4 or more branches. *Perm.*, USSR (European RSFSR).—FIG. 155,6. **S. grandis*; fore wing, $\times 3.4$ (Becker-Migdisova, 1961c).

Tshekardaella BECKER-MIGDISOVA, 1960, p. 59 [**T. tshekardaensis*; OD; = *Tshekardaella tshekardaensis* BECKER-MIGDISOVA, 1948a, p. 130, *nom. nud.*]. Little-known genus, based on wing and body fragments. Fore wing as in *Archescytina* but shorter and more nearly oval. BECKER-MIGDISOVA, 1962b; SZELEGIEWICZ & POPOV, 1978. *Perm.*, USSR (Asian RSFSR).

Uraloscytina ZALESSKY, 1939, p. 40 [**U. prosbolicoides*; OD]. Fore wing as in *Archescytina*, but M more extensively branched and with proximal branch arising about the level of origin of RS. [Type of family Uraloscytinidae ZALESSKY, 1939.] *Perm.*, USSR (Asian RSFSR).—FIG. 155,4. *U. multinervosa* BECKER-MIGDISOVA; fore wing, $\times 4$ (Becker-Migdisova, 1962b).

Family BOREOSCYTIDAE

Becker-Migdisova, 1949

[Boreoscytidae BECKER-MIGDISOVA, 1949a, p. 171]

Little-known family. Fore wing much broader distally than basally; vein M with at least 3 branches. Hind wing and body unknown. *Perm.*

Boreoscyta BECKER-MIGDISOVA, 1949a, p. 172 [**B. nefasta*; OD]. Fore wing triangular; RS with pectinate branches directed to costal margin. ROHDENDORF, 1957. *Perm.*, USSR (European RSFSR).—FIG. 156,4. *B. mirabilis* BECKER-MIGDISOVA; fore wing, $\times 6.5$ (Becker-Migdisova, 1949a).

Archescytinopsis BECKER-MIGDISOVA, 1949a, p. 175 [**Sojanoscytina latipennis* MARTYNOV, 1933c, p. 887; OD]. Fore wing not so markedly triangular as in *Boreoscyta*; RS without pectinate branches. *Perm.*, USSR (European RSFSR).—FIG. 156,3. **A. latipennis* (MARTYNOV); fore wing, $\times 6.5$ (Becker-Migdisova, 1949a).

Family PINCOMBEIDAE

Tillyard, 1922

[Pincombeidae TILLYARD, 1922a, p. 282]

Little-known family of uncertain affinities. Fore(?) wing triangular; veins M and CUA originating at same point on R; anal area apparently very narrow. Hind wing appar-




FIG. 156. Boreoscytidae and Pincombeidae (p. 243).

ently smaller than fore; R, M, and CUA diverging from same place. Body unknown. *Perm.*

Pincombea TILLYARD, 1922a, p. 282 [**P. mirabilis*; OD]. Fore(?) wing: M with 3 branches; CUA forked to half its length; one crossvein between M and RS, none between CUA and M. EVANS, 1956. *Perm.*, Australia (New South Wales).—FIG. 156,2. **P. mirabilis*; fore(?) wing, $\times 16$ (Tillyard, 1922a).

Eupincombea DAVIS, 1942, p. 114 [**E. postica*; OD]. Hind wing: RS, M, and CUA unbranched; costal area triangular. EVANS, 1956. *Perm.*, Australia (New South Wales).—FIG. 156,1. **E. postica*; hind wing, $\times 20$ (Davis, 1942).

Protopincombea EVANS, 1943b, p. 193 [**P. obscura*; OD]. Fore wing as in *Pincombea*, but 2 crossveins between RS and M and one between

M and CUA. EVANS, 1956. *Perm.*, Australia (New South Wales).

Family PROTOPSYLLIDIIDAE
Carpenter, 1931

[Protopsyllidiidae CARPENTER, 1931b, p. 119] [=Permaphidopsidae BECKER-MIGDISOVA, 1960, p. 57]

Fore wing variable in shape; vein SC not a distinct vein; RS typically unbranched; stem of M fused with CUA; anal area small but distinct and coriaceous; CUP straight. Hind wing smaller than fore wing. Body structure little known; legs slender. *Perm.-Jur.*

Protopsyllidium TILLYARD, 1926a, p. 26 [**P. australe*; OD]. Fore wing with RS arising well before midwing; M with 2 branches. EVANS, 1956. *Perm.*, Australia (New South Wales).—FIG. 157,9. **P. australe*; fore wing, $\times 16$ (Tillyard, 1926a).

Asiopsyllidium BECKER-MIGDISOVA, 1959a, p. 113 [**A. unicum*; OD]. Fore wing much wider distally than basally; RS arising well before midwing; M with 2 branches; CUA with a narrow fork. *Trias.*, USSR (Kirghiz).—FIG. 157,6. **A. unicum*; fore wing, $\times 10$ (Becker-Migdisova, 1959a).

Belpsylla EVANS, 1943b, p. 192 [**B. reticulata*; OD]. Fore wing broad distally; M with 3 straight branches; one crossvein between RS and M1+2 and another between RS and M1; CUA with small fork; anal area with Y-shaped vein. *Perm.*, Australia (New South Wales).—FIG. 157,10. **B. reticulata*; fore wing, $\times 12$ (Evans, 1943b).

Cicadellopsis MARTYNOV, 1937a, p. 107 [**C. incerta*; OD]. Fore wing with costal margin strongly convex; RS arising near wing base; M forked; CUA with small distal fork. EVANS, 1956; BECKER-MIGDISOVA, 1962b. *Trias.-Jur.*, USSR (Kirghiz).—FIG. 157,8. **C. incerta*, *Jur.*; fore wing, $\times 13$ (Martynov, 1937a).

Cicadopsyllidium BECKER-MIGDISOVA, 1959a, p. 112 [**C. elongatum*; OD]. Little-known genus. Fore wing narrow; pterostigma apparently absent; RS arising well before midwing; M and CUA apparently fused basally. [Family assignment doubtful.] *Trias.*, USSR (Kirghiz).

Clavopsyllidium DAVIS, 1942, p. 117 [**C. minutum*; OD]. Fore wing as in *Protopsyllidium*, but M with 3 branches; CUA1 arched. EVANS, 1943b, 1956. *Perm.*, Australia (New South Wales).—FIG. 157,7. **C. minutum*; fore wing, $\times 18$ (Davis, 1942).

Permaphidopsis BECKER-MIGDISOVA, 1960, p. 58 [**P. sojanensis*; OD]. Little-known genus, based on hind wing. Wing broad distally; M coalesced

basally with CUA; CUA forked distally with strongly curved CUA1. SZELEGIEWICZ & POPOV, 1978. *Perm.*, USSR (European RSFSR).

Permapsyllidium TILLYARD, 1926a, p. 27 [**P. mitchelli*; OD]. RS arising near midwing; M with 3 branches. CARPENTER, 1931b. *Perm.*, Australia (New South Wales).—FIG. 157,5. **P. mitchelli*; fore wing, $\times 14$ (Tillyard, 1926a).

Permapsyllidops DAVIS, 1942, p. 116 [**P. stanleyi*; OD]. Fore wing similar to *Protopsyllidium*, but CUP absent or poorly developed; M with 3 branches. EVANS, 1956. *Perm.*, Australia (New South Wales).—FIG. 157,1. **P. stanleyi*; fore wing, $\times 15$ (Davis, 1942).

Permapsylloides EVANS, 1943b, p. 193 [**P. insolita*; OD]. Fore wing of uniform width; costal area wide; RS arising before midwing, curved; M apparently with 2 branches; CUA sinuate; anal area with Y-shaped vein. EVANS, 1956. *Perm.*, Australia (New South Wales).—FIG. 157,4. **P. insolita*; fore wing, $\times 12$ (Evans, 1943b).

Permothea TILLYARD, 1926a, p. 28 [**P. latipennis*; OD]. Fore wing much as in *Protopsyllidium*, but M with 3 branches. CARPENTER, 1931b; EVANS, 1956. *Perm.*, Australia (New South Wales).

Permotheella DAVIS, 1942, p. 116 [**P. scytinopteroides*; OD]. RS strongly curved; M with 3 branches; anal veins forming Y-shaped vein. EVANS, 1943b, 1956. *Perm.*, Australia (New South Wales).—FIG. 157,3. **P. scytinopteroides*; fore wing, $\times 14$ (Davis, 1942).

Propatrix BECKER-MIGDISOVA, 1960, p. 55 [**P. psylloides*; OD; =*P. psylloides* BECKER-MIGDISOVA, 1948a, p. 130, *nom. nud.*]. Fore wing with long pterostigmal area. RS arising at midwing; M with 3 branches; CUA with wide fork. BECKER-MIGDISOVA, 1962b; SZELEGIEWICZ & POPOV, 1978. *Perm.*, USSR (European RSFSR).—FIG. 157,2. **P. psylloides*; fore wing and body, $\times 8$ (Becker-Migdisova, 1960).

Psocopsyllidium DAVIS, 1942, p. 115 [**P. media*; OD]. Fore wing as in *Protopsyllidium* but more slender. EVANS, 1943b, 1956. *Perm.*, Australia (New South Wales).

Psococytina DAVIS, 1942, p. 112 [**P. bifida*; OD]. Similar to *Protopsyllidium*, but M with 3 branches; RS arising at midwing with distal fork. EVANS, 1956. *Perm.*, Australia (New South Wales).—FIG. 158,2. **P. bifida*; fore wing, $\times 12$ (Davis, 1942).

Psyllidella EVANS, 1943b, p. 192 [**P. magna*; OD]. Fore wing with RS arising beyond midwing; M with 3 long branches; costal margin sinuate. *Perm.*, Australia (New South Wales).—FIG. 158,5. **P. magna*; fore wing, $\times 10$ (Evans, 1943b).

Psyllidiana EVANS, 1943b, p. 192 [**P. davisia*; OD] [=*Protopsyllidops* EVANS, 1943b, p. 192 (type, *P. minuta*)]. Fore wing as in *Protopsyll-*




FIG. 157. Protosylliidae (p. 244).

lidium, but RS arising near midwing and very straight; CUA deeply forked. EVANS, 1956. *Perm.*, Australia (New South Wales).—FIG. 158, 1. **P. davisia*; fore wing, $\times 22$ (Evans, 1943b).

Tomiopsyllidium BECKER-MIGDISOVA, 1959a, p. 112 [**T. iljinskiense*; OD]. Fore wing slender, triangular; RS arising just before midwing, curving away from R distally. BECKER-MIGDISOVA, 1961c. *Perm.*, USSR (Asian RSFSR).—FIG. 158, 4.

**T. iljinskiense*; fore wing, $\times 22$ (Becker-Migdisova, 1960).

Triassoptylla TILLYARD, 1918b, p. 753 [**T. pleiocoides*; OD]. Little-known genus, based on wing fragment; RS curved; M with 3 branches. EVANS, 1956. *Trias.*, Australia (New South Wales).

Triassothea EVANS, 1956, p. 236 [**T. analis*; OD]. Fore wing as in *Protosyllidium*, but RS arising near wing base; M+CUA very short; M with distal fork. *Trias.*, Australia (Queensland).




FIG. 158. Protopsyllidiidae (p. 244–246).

— FIG. 158.3. **T. analis*; fore wing, $\times 14$ (Evans, 1956).

Family GENAPHIDIDAE Handlirsch, 1907

[Genaphididae HANDLIRSCH, 1907, p. 643]

Little-known family. Antennae with 7 segments, bearing annular, secondary sense organs (rhinaria). Fore wing with all veins of nearly same thickness; M arising at level of base of pterostigma, with 3 terminal branches; CUA with short base, arising from stem R+M. HEIE, 1967, 1985; SHAPOSHNIKOV, 1979b, 1980. *Jur.*

Genaphis HANDLIRSCH, 1907, p. 643 [**Apbis valdensis*; OD]. Little-known genus. RS arising near middle of pterostigma. HEIE, 1967. *Jur.*, England. — FIG. 159.1. **G. valdensis*; fore wing, $\times 18$ (Heie, 1967).

Juraphis SHAPOSHNIKOV, 1979b, p. 66 [**J. crassipes*; OD]. Fore wing with RS arising slightly distally of middle of pterostigma. Antennae and legs stout. HEIE, 1985. *Jur.*, USSR (Kazakh). — FIG. 159.4. **J. crassipes*; fore and hind wings, $\times 18$ (Shaposhnikov, 1979b).

Family CANADAPHIDIDAE Richards, 1966

[*nom. transl.* KONONOVA, 1976, p. 119, ex Canadaphidinae RICHARDS, 1966, p. 757]

Head dorsoventrally flattened, prolonged anteriorly; antennal bases ventral, in front of compound eyes; antennae with 5 to 6 segments; rostrum apparently very short; tarsi long; ovipositor well developed; siphuncles and cauda apparently not present. Fore wing with vein M with two forks. Hind wing relatively large. *Cret.*

Canadaphis ESSIG in CARPENTER & others, 1937, p. 19 [**C. carpenteri*; OD]. M of fore wing arising near origin of CUA1; CUA1 slightly sinuate; tarsi with 2 segments. HEIE, 1967, 1981; KONONOVA, 1976. [A record of this genus (*C. mordvilko*) KONONOVA, 1976, p. 120] from the Cretaceous of USSR (Asian RSFSR) is very questionable. See KONONOVA, 1976, and HEIE, 1985.] *Cret.*, Canada (Manitoba). — FIG. 159.2. **C. carpenteri*; dorsal view, $\times 35$ (Essig in Carpenter & others, 1937).

Alloambria RICHARDS, 1966, p. 756 [**A. caudata*; OD]. Antennae with at least 5 segments. Fore wing with CUA1 and CUA2 arising independently from stem SC+R+M; CUA1 sinuate. Tarsi with 2 segments. *Cret.*, Canada (Manitoba). — FIG. 159.3. **A. caudata*; dorsal view, $\times 50$ (Richards, 1966).

Pseudambria RICHARDS, 1966, p. 758 [**P. longirostris*; OD]. Antennae with 6 segments. Fore wing with CUA1 sinuate; CUA2 very weakly developed. HEIE, 1981, 1985. *Cret.*, Canada (Manitoba).

Family PALAEOAPHIDIDAE Richards, 1966

[*nom. transl.* KONONOVA, 1976, p. 121, ex Palaeoaphidinae RICHARDS, 1966, p. 750]

Similar to Canadaphididae, but antennae with 7 segments; ovipositor well developed. Fore wing with vein RS arising from proxi-




FIG. 159. Genaphididae and Canadaphididae (p. 246).

mal third of pterostigma; hind wing relatively shorter than in Canadaphididae. HEIE, 1985. *Cret.*

Palaeoaphis RICHARDS, 1966, p. 750 [**P. archimeditia*; OD]. Little-known genus. Media of fore wing incomplete basally; legs with short hairs. [The assignment of *P. incognata* KONONOVA, 1976, p. 121 (Cretaceous of USSR) to the family Palaeoaphididae is very uncertain.] HEIE, 1985. *Cret.*, Canada (Manitoba).—FIG. 160, I. **P. archimeditia*; fore wing, $\times 45$ (Richards, 1966).

Ambaraphis RICHARDS, 1966, p. 752 [**A. costalis*; OD]. Similar to *Palaeoaphis*, but apical tarsal segments with long, conspicuous preapical setae. HEIE, 1985. *Cret.*, Canada (Manitoba).

Family SHAPOSHNIKOVIIDAE Kononova, 1976

[Shaposhnikoviidae KONONOVA, 1976, p. 122]

Little-known family. Antennae with 7 segments, its total length only half that of fore wing. Fore wing: vein M with 3 terminal

branches; CUA1 and CUA2 widely separated basally. HEIE, 1981, 1985. *Cret.*

Shaposhnikovia KONONOVA, 1976, p. 122 [**S. electri*; OD]. Fore wing with M arising from base of pterostigma. Second segment of fore tarsus about one-fourth as long as tibia. HEIE, 1981. *Cret.*, USSR (Asian RSFSR).

Family OVIPARASIPHIDAE Shaposhnikov, 1979

[Oviparasiphidae SHAPOSHNIKOV, 1979b, p. 75]

Antennae with annular, secondary sense organs (rhinaria). Fore wing with vein RS arising from middle of pterostigma; M with 3 branches; CUA1 and CUA2 originating separately from a common stem (SC+R+M). Ovipositor large. *Cret.*

Oviparasiphum SHAPOSHNIKOV, 1979b, p. 75 [**O. jakovlevi*; OD]. Rhinaria on antennae forming convex rings. Femora stout. *Cret.*, Mongolia.

Family TAJMYRAPHIDIDAE

Kononova, 1975

[Tajmyraphididae KONONOVA, 1975, p. 795]

Antennae with 4 to 6 segments. Fore wing broadly rounded distally; pterostigma short, vein RS not connected to it; M with one fork; CUA1 about three times as long as CUA2. HEIE, 1985. *Cret.*

Tajmyraphis KONONOVA, 1975, p. 796 [**T. zberichini*; OD]. Antennae with 5 or 6 segments. *Cret.*, USSR (Asian RSFSR).

Jantardakhia KONONOVA, 1975, p. 804 [**J. electri*; OD]. Antennae with 5 segments. Fore wing with bases of CUA1 and CUA2 widely separated. *Cret.*, USSR (Asian RSFSR).

Khatangaphis KONONOVA, 1975, p. 803 [**K. sibirica*; OD]. Similar to *Tajmyraphis*, but antennae with 4 or 5 segments; pterostigma of fore wing very short. *Cret.*, USSR (Asian RSFSR).

Retinaphis KONONOVA, 1975, p. 801 [**R. glandulosa*; OD]. Similar to *Tajmyraphis*, but antennae longer, with 6 segments. *Cret.*, USSR (Asian RSFSR).

Family MINDARIDAE Tullgren, 1909

[Mindaridae TULLGREN, 1909, p. 58]

Cauda subtriangular. Fore wing with pterostigma narrow, pointed, extending to apex of wing; vein RS arising from the proximal part of pterostigma. *Cret.—Holo.*

Mindarus KOCH, 1857, p. 277 [= *Pterostigma* BUCKTON, 1883, p. 178 (type, *P. recurvus*); *Schizoneuroides* BUCKTON, 1883, p. 178 (type, *S. scudderi*); *Sychnobrochus* SCUDDER, 1890, p. 268 (type, *S. reviviscens*)]. BAKER, 1922; HEIE, 1967, 1969b, 1985. *Oligo.*, Europe (Baltic), USA (Colorado)—*Holo.*

Nordaphis KONONOVA, 1977, p. 593 [**N. sukatchevae*; OD]. Little-known genus. Antennae with 6 segments. Fore wing with pterostigma very elongate; RS straight; M with one fork. Legs long. [Placed in Drepanosiphidae by KONONOVA but transferred to Mindaridae by HEIE (1985).] *Cret.*, USSR (Asian RSFSR).

Family HORMAPHIDIDAE

Mordvilko, 1908

[Hormaphididae MORDVILKO, 1908, p. 364]

Antennae with 3 to 5 segments, much shorter than body; antennae of alate form with narrow, ringlike, secondary rhinaria. Fore wing with veins CUA1 and CUA2 arising from same point on SC+R+M. *Oligo.—Holo.*

Hormaphis OSTEN-SACKEN, 1861, p. 422. *Holo.*
Electrocornia HEIE, 1972, p. 249 [**E. antiqua*; OD]. Little-known genus, based on nymph. Antennae with 5 segments; head and pronotum fused; frons with 2 hornlike processes. [Originally placed in Thelaxidae but later transferred to Hormaphididae (HEIE, 1985).] *Oligo.*, Europe (Baltic).

Family ELEKTRAPHIDIDAE

Steffan, 1968

[Elektraphididae STEFFAN, 1968, p. 11]

Antennae with 5 segments. Fore wing with vein RS greatly reduced; M typically without branches; CUA1 and CUA2 arising from stem CUA or originating independently from stem SC+R+M. KONONOVA, 1976. *Cret.—Oligo.*

Schizoneurites COCKERELL, 1915, p. 487 [**S. brevirostris*; OD] [= *Antiquaphis* HEIE, 1967, p. 88 (type, *A. robustus*); *Elektraphis* STEFFAN, 1968, p. 11 (type, *E. polykrypta*)]. Fore wing with CUA1 and CUA2 arising from common stem CUA. Antennae with transverse folds. HEIE, 1967, 1976, 1980, 1985; STEFFAN, 1968. *Oligo.*, Europe (Baltic), England.—FIG. 160, *S. robustus* (HEIE), Baltic; fore wing, $\times 34$ (Heie, 1967).

Antonaphis KONONOVA, 1977, p. 589 [**A. brachycera*; OD]. Antennae short, with 5 segments. Fore wing with RS long, slightly curved; M branched once. [Originally placed in Pemphigidae but transferred to *Elektraphididae* by HEIE (1985).] *Cret.*, USSR (Asian RSFSR).

Tajmyrella KONONOVA, 1976, p. 118 [**T. cretacea*; OD]. Similar to *Schizoneurites*, but CUA1 and CUA2 arising independently from stem SC+R+M. HEIE, 1981. *Cret.*, USSR (Asian RSFSR).

Family THELAXIDAE Baker, 1920

[Thelaxidae BAKER, 1920, p. 21]

Antennae with 5 segments. Media of fore wing with 2 terminal branches. Hind wing with two oblique veins. *Oligo.—Holo.*

Thelaxes WESTWOOD, 1840, p. 118. *Holo.*
Palaeothelaxes HEIE, 1967, p. 42 [**P. setosa*; OD]. Little-known genus. All body segments of apterous form with very thick, large setae; frons of alate form with similar large setae. *Oligo.*, Europe (Baltic).

Family ANOECIIDAE Tullgren, 1909

[Anoeciidae TULLGREN, 1909, p. 186]

Antennae commonly with 6 segments, and in alate forms with oval or subcircular sec-




FIG. 160. Palaeoaphididae, Elektraphididae, Pemphigidae, and Drepanosiphidae (p. 247–251).

ondary rhinaria; marginal tubercles present on prothorax and some abdominal segments. Pterostigma of fore wing not more than four times longer than its width. *Oligo.*—*Holo.*

Anoecia KOCH, 1857, p. 275. *Holo.*

Berentaphis HEIE, 1971, p. 262 [*Lachnus cimicoides* GERMAR & BERENDT, 1856, p. 5; OD]. Little-known genus, based on apterous form. Antennae with 6 segments, distal segments conspicuously thickened; head and pronotum not fused. *Oligo.*, Europe (Baltic).

Family PEMPHIGIDAE Koch, 1857

[Pemphigidae KOCH, 1857 p. viii]

Antennae short, usually with 6 segments and with one very short terminal process. Fore wing with vein M unbranched or with one fork. Hind wing with 1 or 2 oblique veins. *Oligo.*—*Holo.*

Pemphigus HARTIG, 1839, p. 645. *Holo.*

Eriosoma LEACH, 1818, p. 60. HEIE, 1968a, 1969a,

- 1969c, 1985. *Mio./Plioc.*, Europe (Germany)-*Holo*.
- Germaraphis** HEIE, 1967, p. 47 [**Lachnus dryoides* GERMAR & BERENDT, 1856, p. 29; OD]. Little-known genus, based mainly on apterous specimens. Antennae with 5 or 6 segments, the second one at least as long as the fourth. [Originally placed in Phloeomyzidae by HEIE (1967) and BECKER-MIGDISOVA (1973) but transferred to Pemphigidae by HILLE RIS LAMBERS (1980) and HEIE (1985).] HEIE, 1969b, 1972, 1985; BECKER-MIGDISOVA, 1973. *Oligo.*, Europe (Baltic). — FIG. 160,7. **G. dryoides*; apterous specimen, reconstruction, $\times 30$ (Heie, 1967).
- Succinaphis** HEIE, 1967, p. 173 [**S. flauensgaardii*; OD]. Apparently similar to *Pemphigus* (recent). Media of fore wing branched; wing membrane with fine reticulation. HEIE, 1985. *Oligo.*, Europe (Baltic). — FIG. 160,4. **S. flauensgaardii*; fore and hind wings, $\times 45$ (Heie, 1967).
- Family DREPANOSIPHIDAE**
Koch, 1857
- [Drepanosiphidae Koch, 1857, p. vii]
- Closely allied to the Aphididae. Secondary transverse or circular rhinaria usually present on third antennal segment of alate females. Fore wing: vein M with 2 or 3 terminal branches. Hind wing with 2 or 3 oblique veins. HEIE, 1980, 1982. *Cret.-Holo*.
- Drepanosiphum** KOCH, 1855, p. 201. *Holo*.
- Aixaphis** HEIE, 1970b, p. 115 [**Tetraneura oligocenica* THÉOBALD, 1937a, p. 16; OD]. Antennae about half body length, with 6 segments. Fore wing: M with 3 terminal branches; CUA1 and CUA2 arising independently from stem SC+R+M, their bases relatively remote. [Family assignment doubtful.] HEIE, 1985. *Oligo.*, Europe (France).
- Aniferella** RICHARDS, 1966, p. 759 [**A. bostoni*; OD]. Antennae with 5 segments. Fore wing with well developed pterostigma; RS nearly straight; M with 2 forks; CUA1 and CUA2 arising separately from stem SC+R+M. HEIE, 1981, 1985. *Cret.*, Canada (Alberta). — FIG. 160,3. **A. bostoni*; fore and hind wings, $\times 35$ (Richards, 1966).
- Balticaphis** HEIE, 1967, p. 160 [**B. exsiccata*; OD]. Little-known genus, based on apterous form. Antennae with 5 or 6 segments. Fore femora thickened. HEIE, 1985. *Oligo.*, Europe (Baltic).
- Balticomaraphis** HEIE, 1967, p. 167 [**B. latens*; OD]. Little-known genus, based on cast cuticle of nymph. Antennae with 6 segments; ocular tubercles well developed. *Oligo.*, Europe (Baltic).
- Conicaudus** HEIE, 1972, p. 255 [**C. longipes*; OD]. Little-known genus, based on alate form. Antennae about as long as body. M of fore wing with 3 or 4 terminal branches. Tarsi very long. *Oligo.*, Europe (Baltic).
- Cretacallis** SHAPOSHNIKOV, 1979a, p. 730, footnote [**C. polysensoria*; OD]. Antennae with 6 segments. Fore wing: M with 3 terminal branches; CUA1 and CUA2 originating independently from stem SC+R+M. [Family position doubtful.] *Cret.*, Mongolia.
- Electrocallis** HEIE, 1967, p. 147 [**E. bakeri*; OD] [= *Dimeraphis* BECKER-MIGDISOVA, 1973, p. 87 (type, *D. arnoldii*)]. Antennae of alate form much longer than body and composed of 6 segments. Fore wing with pterostigma short; M with 3 terminal branches; CUA1 and CUA2 arising separately from stem SC+R+M. Fore femora thicker than the others. *Oligo.*, Europe (Baltic).
- Megantennaphis** HEIE, 1967, p. 142 [**M. haurienseis*; OD]. Antennae with 6 segments and much longer than body. Fore wing with pterostigma long, pointed; RS almost straight; M with 3 terminal branches. Fore and hind femora large and strong. BECKER-MIGDISOVA, 1973. *Oligo.*, Europe (Baltic).
- Megapodaphis** HEIE, 1967, p. 155 [**M. monstrabilis*; OD]. Antennae with 6 segments and at least as long as body. Fore wing: M with 2 terminal branches. Fore femora strongly thickened. HEIE, 1972, 1985. *Oligo.*, Europe (Baltic).
- Mengeaphis** HEIE, 1967, p. 113 [**Lachnus glandulosus*; OD]. Little-known genus, based on immature nymphs. Antennae with 4 segments; rostrum at least twice the length of body. BECKER-MIGDISOVA, 1973. *Oligo.*, Europe (Baltic). — FIG. 160,6. **M. glandulosus*; dorsal view, $\times 85$ (Becker-Migdisova, 1973).
- Oligocallis** HEIE, 1967, p. 133 [**O. larssoni*; OD]. Little-known genus, based on alate form. Similar to *Pterasthenica* (recent), but venation of fore wing less reduced in *Oligocallis*. HEIE, 1972. *Oligo.*, Europe (Baltic).
- Oryctaphis** SCUDDER, 1890, p. 266 [**O. lesueurii*; OD]. Little-known genus, possibly a synonym of *Siphonophoroides*. HEIE, 1985. *Oligo.*, USA (Colorado).
- Palaeophyllaphis** HEIE, 1967, p. 97 [**P. longirostris*; OD]. Antennae with 6 segments. Fore wing: M with 2 or 3 terminal branches; pterostigma slightly pointed but short. GERMAR & BERENDT, 1856; HEIE, 1972, 1985. *Oligo.*, Europe (Baltic).
- Palaeosiphon** HEIE, 1967, p. 119 [**Aphis hirsuta* GERMAR & BERENDT, 1856, p. 6; OD]. Little-known genus. Antennae of apterous form with 5 segments. Fore wing: M with 3 terminal branches. Hind wing with only one oblique vein.

- Head and first two thoracic segments of alate form with long, curved, hornlike projections. HEIE, 1971. *Oligo.*, Europe (Baltic).
- Siphonophoroides** BUCKTON, 1883, p. 176 [**S. antiqua*; OD][=*Archilachus* BUCKTON, 1883, p. 177 (type, *A. pennata*); *Aphantaphis* SCUDDER, 1890, p. 253 (type, *S. exsuca*); *Cataneura* SCUDDER, 1890, p. 245 (type, *C. absens*); *Amalancon* SCUDDER, 1890, p. 270 (type, *A. lutosus*)]. Antennae slender, longer than body. Fore wing with RS very long, relatively straight, arising from proximal half of pterostigma; M with 3 terminal branches. COCKERELL, 1908u, 1909b; HEIE 1967, 1985. *Eoc.*, Europe (Denmark); *Oligo.*, USA (Colorado).—FIG. 160,2. **S. antiqua*; fore wing, $\times 14$ (Heie, 1967).
- Sternaphis** HEIE, 1972, p. 257 [**S. electricola*; OD]. Fore wing with RS short and straight; M with 2 terminal branches. *Oligo.*, Europe (Baltic).
- Succaphis** HEIE, 1967, p. 110 [**S. bolgeri*; OD]. Little-known genus, based on apterous form. Head and pronotum separated; antennae with 4 segments; rostrum longer than body. [Family assignment doubtful.] HEIE, 1985. *Oligo.*, Europe (Baltic).
- Tertiaphis** HEIE, 1969b, p. 144 [**T. haentzscheli*; OD]. Antennae with 6 segments and shorter than body. Fore wing: M with 2 terminal branches; CUA1 and CUA2 arising separately from stem SC+R+M. HEIE, 1985. *Oligo.*, Europe (Baltic).
- Zymus** HEIE, 1972, p. 254 [**Z. succincola*; OD]. Little-known genus, based on nymph. Antennae with 4 segments and with long, filamentous terminal segment; head and pronotum fused; strong bristles on head and posterior part of abdomen. *Oligo.*, Europe (Baltic). — FIG. 160,8. **Z. succincola*; dorsal view of nymph, $\times 24$ (Heie, 1972).

Family APHIDIDAE Latreille, 1802

[Aphididae LATREILLE, 1802a, p. 263]

Compound eyes large in all instars; antennae commonly with 6 segments (rarely with 5), at least half length of body. Fore wing: vein RS with 2 or 3 terminal branches; CUA and CUP arising independently from stem R+M+CU. Hind wing commonly with 2 oblique veins, rarely only one. Wings slanted at rest. *Cret.—Holo.*

Aphis LINNÉ, 1758, p. 451. *Holo.*

Aphidocallis KONONOVA, 1977, p. 595 [**A. caudatus*; OD]. Antennae with 5 segments. Fore wing with pterostigma short, extending only to about level of midwing; M with 3 terminal branches. *Cret.*, USSR (Asian RSFSR).

Baltichaitophorus HEIE, 1967, p. 180 [**B. jurtlandicus*; OD]. Little-known genus, based on apterous forms. Antennae with 6 segments, about as long as body. HEIE, 1980. *Oligo.*, Europe (Baltic).

Diatomyzus HEIE, 1970a, p. 163 [**D. eocaenicus*; OD]. Little-known genus, based on alate specimens. Similar to several existing genera, but RS of fore wing unusually long. *Eoc.*, Europe (Denmark).

Pseudamphorophora HEIE, 1967, p. 175 [**P. succini*; OD]. Little-known genus, based on apterous forms. [Family assignment doubtful.] HEIE, 1971, 1980. *Oligo.*, Europe (Baltic).

Family LACHNIDAE Koch, 1857

[Lachnidae KOCH, 1857, p. vii]

Similar to Anoeciidae, but prothorax and abdominal segments lacking marginal tubercles. Pterostigma of fore wing commonly much longer than 4 times its width. *Mio./Plio.—Holo.*

Lachnus BURMEISTER, 1835, p. 92. *Holo.*

Longistigma WILSON, 1909, p. 385. HEIE & FRIEDRICH, 1971; HEIE, 1985. *Mio./Plio.*, Iceland—*Holo.*

Family ALEYRODIDAE Westwood, 1840

[Aleyrodidae WESTWOOD, 1840, p. 442]

Wings slightly thickened, commonly covered with a powdery wax. Fore wing venation weakly formed, only veins R and M extending to distal part of wing. Antennae with 7 segments; terminal abdominal segment with a large, dorsal opening, associated with storage of honey dew. *Oligo.—Holo.*

Aleyrodes LATREILLE, 1796, p. 93. [Generic assignment of fossil doubtful.] MENGE, 1856; SCHLEE, 1970. *Oligo.*, Europe (Baltic)—*Holo.*

Aleurodicus DOUGLAS, 1892, p. 32. [Generic assignment of fossil doubtful.] COCKERELL, 1919e; SCHLEE, 1970. *Mio.*, Burma—*Holo.*

Family COLEOSCYTIDAE Martynov, 1935

[Coleoscytidae MARTYNOV, 1935c, p. 24]

Fore wing oval, weakly coriaceous, membranous distally; subcostal area abruptly widened at base; costal margin at right angles to wing axis at this point; vein SC marginal;




FIG. 161. Coleoscytidae, Cicadopsyllidae, Psyllidae, and Margarodidae (p. 252-254).

R long, with a branch to costal margin near midwing; M and CUA distally branched; CUP straight, unbranched; 1A and 2A with a common stem. Hind wing membranous, widened distally, more slender than fore wing, with concave anterior margin; subcostal area very narrow; M forked, CUA with a very wide fork; anal area narrow. Head hypognathous; eyes not divided. Hind coxae large. *Perm.*

Coleoscyta CARPENTER, herein [**C. rotundata* MARTYNOV, 1935c, p. 24; OD] [= *Coleoscytodes* CARPENTER, herein (type, *C. venosa* MARTYNOV,

1935c, p. 24; OD)]. Fore wing very broad, costal margin thickened; RS with distal fork. [The original generic names, *Coleoscyta* and *Coleoscytodes*, were *nomina nuda* (MARTYNOV, 1935c).] BECKER-MIGDISOVA, 1962b. *Perm.*, USSR (European RSFSR). — FIG. 161,4a. **C. rotundata*; fore wing, $\times 8$. — FIG. 161,4b. *C. venosa* (MARTYNOV); hind wing, $\times 8$ (Becker-Migdisova, 1960).

Sojanopsylla BECKER-MIGDISOVA, 1960, p. 45 [**S. brevipennis*; OD]. Fore wing as in *Coleoscyta*, but subcostal area gradually widened basally and R and RS longer; M with 3 or 4 branches. *Perm.*, USSR (European and Asian RSFSR). — FIG. 161,6. **S. brevipennis*; fore wing, $\times 4.5$ (Becker-Migdisova, 1960).

Family CICADOPSYLLIDAE

Martynov, 1931

[*nom. transl.* MARTYNOV, 1935c, p. 16, ex Cicadopsyllinae MARTYNOV 1931c, p. 172]

Fore wing elongate oval, commonly membranous; subcosta apparently close to costal margin; RS long, ending near wing apex. Hind wing with M apparently arising from stem of R; CUA originating independently of R + M. Head hypognathous, with protuberances on vertex. Hind coxae conical, elongate. BECKER-MIGDISOVA, 1962b. *Perm.*

Cicadopsylla MARTYNOV, 1931c, p. 173 [**C. permiana*; OD]. Fore wing with M forking near level of midwing. *Perm.*, USSR (European RSFSR).—FIG. 161,5. **C. permiana*; fore wing, $\times 4$ (Becker-Migdisova, 1962b).

Cicadopsis BECKER-MIGDISOVA, 1959a, p. 110 [**C. rugosipenna*; OD]. Similar to *Cicadopsylla*, but R without distal, anterior branch. *Perm.*, USSR (Asian RSFSR).—FIG. 161,7. **C. rugosipenna*; fore wing, $\times 8$ (Becker-Migdisova, 1962b).

Scytoneura MARTYNOV, 1935c, p. 16 [**S. elliptica*; OD]. Fore wing similar to *Cicadopsylla*, but M dividing more distally. BECKER-MIGDISOVA, 1962b. *Perm.*, USSR (Asian RSFSR).—FIG. 161,3. **S. elliptica*; fore wing, $\times 3$ (Becker-Migdisova, 1962b).

Scytoneurella ZALESSKY, 1939, p. 39 [**S. major*; OD]. Fore wing membranous, costal margin slightly convex; M dividing distally of fork of CUA, with 3 short branches. BECKER-MIGDISOVA, 1962b. *Perm.*, USSR (Asian RSFSR).

Family PSYLLIDAE Latreille, 1807

[Psyllidae LATREILLE, 1807, p. 168]

Fore wing usually coriaceous; costal area broad; veins M and CUA united to form a basal stem; RS arising from R independently; M and CU usually arising as a common stem; RS unbranched; M and CUA forked. Hind wing smaller and more slender, with R and M unbranched. Antennae with 9 to 10 segments. *Jur.*—*Holo.*

Psylla GEOFFREY, 1762, p. 482. BECKER-MIGDISOVA, 1964. *Oligo.*, England; *Mio.*, USSR (European RSFSR)—*Holo.*

Agonoscena ENDERLEIN, 1914, p. 234. BECKER-MIGDISOVA, 1964. *Mio.*, USSR (European RSFSR)—*Holo.*

Catopsylla SCUDDER, 1890, p. 277 [**C. prima*; OD].

Little-known genus. Fore wing as in *Prylla*, but cell of CU much longer. *Oligo.*, USA (Colorado). *Liadopsylla* HANDLIRSCH, 1920, p. 213 [**L. geinitszi*; OD]. Fore wing oval, membranous; R and RS long, parallel; stem of R short; fork of M long. MARTYNOV, 1926b; BECKER-MIGDISOVA, 1949b. *Jur.*, Europe (Germany), USSR (Asian RSFSR).—FIG. 161,2. *L. tenuicornis* MARTYNOV, USSR; fore wing, $\times 20$ (Martynov, 1926b).

Livilla CURTIS, 1836, p. 625. COCKERELL, 1921d. *Oligo.*, England—*Holo.*

Necropsylla SCUDDER, 1890, p. 276 [**N. rigida*; OD]. Little-known genus; fore wing as in *Psyllopsis* (recent) but subtriangular. COCKERELL, 1911b, 1915. *Oligo.*, USA (Colorado), England.

Psyllites COCKERELL, 1914f, p. 636 [**P. crawfordi*; OD]. Little-known genus, probably a synonym of *Catopsylla*. *Oligo.*, USA (Colorado).

Retrocizzia HESLOP-HARRISON, 1961, p. 504. BECKER-MIGDISOVA, 1964. *Mio.*, USSR (European RSFSR)—*Holo.*

Strophingia ENDERLEIN, 1914, p. 233. *Oligo.*, Europe (Baltic)—*Holo.*

Trioza FÖRSTER, 1848, p. 67. BECKER-MIGDISOVA, 1964. *Mio.*, USSR (European RSFSR)—*Holo.*

Family COCCIDAE Fallén, 1814

[Coccidae FALLÉN, 1814, p. 23]

Adults with marked sexual dimorphism. Males with fore wings normally developed; hind wings reduced or halterlike. Females apterous; antennae diverse, commonly much reduced; abdominal spiracles absent. *Oligo.*—*Holo.*

Coccus LINNÉ, 1758, p. 455. MENGE, 1856; COCKERELL, 1906b; BECKER-MIGDISOVA, 1962b. *Oligo.*, Europe (Baltic)—*Holo.*

Family ORTHEZIIDAE

Amyot & Serville, 1843

[Ortheziidae AMYOT & SERVILLE, 1843, p. 619]

Similar to Coccidae. Females with body clearly segmented; antennae with distinct segmentation; abdominal spiracles present. *Oligo.*—*Holo.*

Orthezia BOSC, 1784, p. 173. *Holo.*

Ochyrocoris MENGE, 1856, p. 17 [**O. electrina*; OD]. Little-known genus, probably a synonym of *Orthezia* (recent). COCKERELL, 1906a; BECKER-MIGDISOVA, 1962b. *Oligo.*, Europe (Baltic).

Family MARGARODIDAE

Cockerell, 1899

[Margarodidae COCKERELL, 1899, p. 390]

Males commonly winged, with few unbranched veins. Females with convex body, strongly sclerotized, with clear segmentation; abdomen with an anal tube or a sclerotized ring, lacking setae. *Cret.*—*Holo.*

Margarodes GUILDING, 1829, p. 118. *Holo.*

Acreegris KOCH in KOCH & BERENDT, 1854, p. 123 [**A. crenata*; OD]. Female adult: antennae with 9 segments; body entirely or nearly devoid of setae; tarsi two-segmented. Male adult: compound eyes; wings with a single vein paralleling the costal margin to wing apex; M delicate, bisecting the wing diagonally; hind wing reduced to slender halteres; antennae with at least 8 segments; tarsi one-segmented; abdomen with long threads of wax arising from clusters of dorsal ducts. FERRIS, 1941. *Oligo.*, Europe (Baltic). —FIG. 161, 1. **A. crenata*; fore wing of male, $\times 6$ (Ferris, 1941).

Electrococcus BEARDSLEY, 1969, p. 271 [**E. canadensis*; OD]. Male small; antennae with 10 segments, pedicel conspicuously enlarged; legs long and slender; compound eye reduced to a single row of ommatidia. Fore wing well developed, with R and M distinct. *Cret.*, Canada (Manitoba).

Family PSEUDOCOCCIDAE

Cockerell, 1905

[Pseudococcidae COCKERELL, 1905, p. 193]

Similar to the Coccidae. Females typically covered with a mealy or filamentous, waxy secretion, commonly protruding as short lateral and long anal filaments; legs well developed. Males apterous or winged, typically with two long caudal wax filaments. *Oligo.*—*Holo.*

Pseudococcus WESTWOOD, 1840, p. 118. *Holo.*

Puto SIGNORET, 1875, p. 394. COCKERELL, 1908g. *Oligo.*, Europe (Baltic)—*Holo.*

Family UNCERTAIN

The following genera, apparently belonging to the suborder Homoptera, are too poorly known to permit assignment to families.

Anconatus BUCKTON, 1883, p. 177 [**A. dorsuosus*; OD]. Little-known aphidoid of uncertain affinities. HEIE, 1967, 1985. *Oligo.*, USA (Colorado).

Annulaphis SHAPOSHNIKOV, 1979b, p. 73 [**A. ras-*

nitsyni; OD]. Little-known genus, based on incomplete specimens; apparently related to *Ellinaphis*. [Originally placed in Palaeoaphididae, but transferred by HEIE (1985) to family uncertain.] *Cret.*, USSR (Asian RSFSR).

Aphidioides MOTSCHULSKY, 1856, p. 29 [**A. succifera*; OD]. Little-known aphidoid genus, based on apterous form. HEIE, 1967, 1985. *Oligo.*, Europe (Baltic).

Aphidulum HANDLIRSCH, 1939, p. 163 [**A. pusillum*; OD]. Little-known genus. HEIE, 1967. *Jur.*, England.

Archeglyphis MARTYNOV, 1931a, p. 89 [**A. cras-sinervis*; OD]. Little-known wing fragment. BECKER-MIGDISOVA, 1961c; ROHDENDORF & RASNITSYN, 1980. *Perm.*, USSR (Asian RSFSR).

Archipsyche HANDLIRSCH, 1906b, p. 624 [**A. eichstattensis*; OD]. Little-known genus, apparently similar to *Limacodites*. *Jur.*, Europe (Germany).

Austrocystina EVANS, 1943b, p. 181 [**A. imperfecta*; OD]. Little-known wing, possibly related to *Archescytinidae*. *Perm.*, Australia (New South Wales).

Beaconiella EVANS, 1963, p. 21 [**B. fennabi*; OD]. All principal veins of fore and hind wings multibranching; possibly a fulgoroid. RIEK, 1973. *Trias.*, Australia (New South Wales).

Beloptesis HANDLIRSCH, 1906b, p. 625 [**B. oppenbeimi*; OD]. Fore wing markedly triangular, nearly as broad as long; venation apparently as in *Limacodites*. Hind wing small, oval. EVANS, 1956. *Jur.*, Europe (Germany).

Bernaea SCHLEE, 1970, p. 18 [**B. neocomica*; OD]. Female with head wider than pronotum; median ocellus present; antennae with 7 segments, the third segment much longer than distal segments. Veins absent on hind wing, represented by lines of pigment. [Placed by SCHLEE in "Aleyrodina *sensu lato*," without family assignment.] *Cret.*, Lebanon.

Borisrohdendorfia BECKER-MIGDISOVA, 1959b, p. 138 [**B. picturata*; OD]. Based on distal fragment of wing. BECKER-MIGDISOVA, 1961c. *Perm.*, USSR (Asian RSFSR).

Cercopidium WESTWOOD, 1854, p. 394 [**C. habni* WESTWOOD, 1854, p. 394; SD CARPENTER, herein]. Little-known genus, based on wing fragment. HEER, 1870a; HENRICKSEN, 1922b. *Jur.*, England; *Eoc.*, Greenland.

Chiliocycla TILLYARD, 1919c, p. 868 [**C. scolopoides*; OD]. Fore wing with strongly thickened costal border; RS present, arising before midwing; closed cell between M1+2 and M3+4; CUA connected to base of M by crossvein. [Type of family Chiliocyclidae EVANS, 1956, p. 209.] EVANS, 1956, 1961. *Trias.*, Australia (Queensland). —FIG. 162, 1. **C. scolopoides*; fore wing, $\times 4.5$ (Evans, 1956).

Cicadellites HEER, 1853a, p. 119 [**C. pallidus* HEER, 1853a, p. 119; SD CARPENTER, herein]. Little-




FIG. 162. Uncertain (p. 254–259).

known homopteron, possibly belonging to the Cercopidae. PITON & THÉOBALD, 1935. *Oligo.*, Europe (France); *Mio.*, Europe (Croatia).
Cicadellium WESTWOOD, 1854, p. 394 [**C. dipsas*; SD HANDLIRSCH, 1907, p. 641] [= *Pseudodelphax* HANDLIRSCH, 1907, p. 641 (type, *Delphax pulcher* BRODIE, 1845, p. 33)]. Little-known genus, based on fore wing. EVANS, 1956. *Jur.*, England.
Cixiites HANDLIRSCH, 1906b, p. 498 [**C. liasinus*; OD]. Little-known wing, possibly related to Fulgoridiidae. BECKER-MIGDISOVA, 1962b. *Jur.*, Europe (Germany).
Cixioides HANDLIRSCH, 1906b, p. 640 [**Cixius maculatus* BRODIE, 1845, p. 33; OD]. Little-known fore wing, possibly related to Cixiidae. *Jur.*, England.
Diphtheropsis MARTYNOV, 1937a, p. 110 [**D. incerta*; OD]. Little-known genus, based on incomplete fore wing with nearly straight costal

margin and long R+M. EVANS, 1956. *Jur.*, USSR (Kirghiz).

Dysmorphoscirtella RIEK, 1973, p. 527 [**D. lobata*; OD]. Little-known genus, based on distal fragment of wing. RIEK, 1976a. [Originally placed in Eoscartarellidae.] *Perm.*, South Africa.

Echinaphis COCKERELL, 1913f, p. 229 [**E. rohweri*; OD]. Little-known genus, based on hind wing and body fragments; apparently related to Greenideidae (recent) and Drepanosiphidae. HEIE, 1967, 1970b, 1985. *Oligo.*, USA (Colorado).

Electromyzus HEIE, 1972, p. 250 [**E. acutirostris*; OD]. Fore wing with RS very slightly curved; M with 2 terminal branches, arising close to point of origin of CUA1 and CUA2. HEIE, 1985. *Oligo.*, Europe (Baltic).

Ellinaphis SHAFOSHNIKOV, 1979b, p. 71 [**E. incognita*; OD]. Little-known genus, originally placed

- in Palaeoaphididae but transferred by HEIE (1985) to category of family uncertain. *Cret.*, USSR (Asian RSFSR).
- Eochilicycla** DAVIS, 1942, p. 114 [**E. angusta*; OD]. Little-known genus, based on fore wing; possibly fulgoroid. EVANS, 1956. *Perm.*, Australia (New South Wales).
- Eocicada** OPPENHEIM, 1888, p. 229 [**E. microcephala*; OD]. Little-known genus, based on wing fragment. EVANS, 1956. *Jur.*, Europe (Germany).
- Eopsyllidium** DAVIS, 1942, p. 114 [**E. delicatulum*; OD]. Little-known hind wing, with CUA free from M basally. Possibly related to the Protopsyllidiidae. EVANS, 1956. *Perm.*, Australia (New South Wales).
- Fulgoridiella** BECKER-MIGDISOVA, 1962a, p. 96 [**F. raetica*; OD]. Little-known wing fragment, possibly related to Fulgoridiidae. *Trias.*, USSR (Kirghiz).
- Fulgoropsis** MARTYNOV, 1937a, p. 165 [**F. dubiosa*; OD]. Little-known genus, based on wing fragment, possibly related to Fulgoridiidae. BECKER-MIGDISOVA, 1962b. *Jur.*, USSR (Kirghiz).
- Gerancho** SCUDDER, 1890, p. 248 [**Lacchnus petrorum* SCUDDER, 1877b, p. 279; OD]. Little-known genus, possibly belonging to Aphidoidea. HEIE, 1967, 1985. *Eoc.*, Canada (British Columbia).
- Gryllites** GERMAR, 1842, p. 82 [**G. dubius*; OD]. Little-known genus, originally placed in Orthoptera. HAGEN, 1862; ASSMANN, 1877; POPOV, 1971. *Jur.*, Europe (Germany).
- Hasrites** COCKERELL, 1922f, p. 161 [**H. muiri*; OD]. Little-known genus. Fore wing elongate; R apparently with a short distal branch; M dividing distally, with 3 terminal branches; CUA with 3 terminal branches. *Oligo.*, England.
- Heidea** SCHLEE, 1970, p. 9 [**H. cretatica*; OD]. Male with head about same width as pronotum; median ocellus present; third antennal segment about as long as distal segments. Vein present in hind wing. [Considered by SCHLEE to be related to the existing and Tertiary Aleurodidae but differing markedly in several traits.] *Cret.*, Lebanon.
- Hooley** COCKERELL, 1922f, p. 160 [**H. indecisa*; OD]. Little-known fore wing; costal margin broad; SC apparently separating from R before midwing, and giving rise to a series of short, oblique veinlets to costal margin. *Oligo.*, England.
- Homopterites** HANDLIRSCH, 1906b, p. 499 [**H. anglicus*; OD]. Little-known fore wing. *Jur.*, England.
- Hylaeoneura** LAMEERE & SEVERIN, 1897, p. 37 [**H. lignei*; OD]. Little-known genus, based on distal fragment of fore wing. R with several long, pectinate branches to costal margin; M with 3 branches. *Cret.*, Europe (Belgium). — FIG. 162.9. **H. lignei*; fore wing, $\times 2.5$ (Handlirsch, 1907).
- Hypocixius** COCKERELL, 1926a, p. 501 [**H. oblitescens*; OD]. Little-known genus, based on incomplete fore wing. Possibly related to Cixiidae. *Tert.* (*epoch unknown*), Argentina (Jujuy).
- Jurocallis** SHAPOSHNIKOV, 1979b, p. 68 [**J. longipes*; OD]. Antennae tapering from base to apex; RS arising from distal part of prerostigmal area; M arising from base of pterostigma and with 3 terminal branches. [Originally placed in Drepanosiphidae.] HEIE, 1985. *Cret.*, USSR (Asian RSFSR).
- Kaltanocicada** BECKER-MIGDISOVA, 1961c, p. 291 [**K. dunstanoides*; OD]. Little-known hind wing, with broadly rounded apex and wide concavity of front wing margin; CUA with long fork. BECKER-MIGDISOVA, 1962b. *Perm.*, USSR (Asian RSFSR).
- Kaltanoscyta** BECKER-MIGDISOVA, 1959a, p. 110 [**K. reticulata*; OD]. Little-known fragment of fore wing, strongly coriaceous and with dense reticulation over wing. Possibly related to Coleoscytidae. *Perm.*, USSR (Asian RSFSR).
- Karabasia** MARTYNOV, 1926b, p. 1356 [**K. paucineris*; OD]. Little-known insect, possibly related to Jassidae. *Jur.*, USSR (Kazakh).
- Karajassus** MARTYNOV, 1926b, p. 1352 [**K. crassineris*; OD]. Little-known insect, possibly close to Cicadellidae. BECKER-MIGDISOVA, 1962b. *Jur.*, USSR (Kazakh).
- Kisyliya** MARTYNOV, 1937a, p. 109 [**K. pylloides*; OD]. Little-known genus, based on fore wing. Nodus and nodal line absent; stem of R slightly shorter than R + M; CUA not coalesced with M. *Jur.*, USSR (Kirghiz). — FIG. 163.4. **K. pylloides*; fore wing, $\times 3$ (Becker-Migdisova, 1962b).
- Larssonaphis** HEIE, 1967, p. 168 [**L. obnubila*; OD]. Little-known aphidoid genus. HEIE, 1985. *Oligo.*, Europe (Baltic).
- Liassocicada** BODE, 1953, p. 201 [**L. antecedens*; OD]. Little-known genus, based mainly on body structure. Rostrum elongate, extending at least to middle of abdomen. [Liassocicada was redefined by WHALLEY (1983) and provisionally placed in the Cicadidae. However, I doubt that our very slight knowledge of the body structures of these Jurassic and Triassic specimens justifies the extension of the range of the Cicadidae to another 150 million years before the Paleocene. Accordingly, the genus *Liassocicada* is herein provisionally placed in the Homoptera, family uncertain.] WHALLEY, 1983. *Trias.*, England; *Jur.*, Europe (Germany).
- Limacodites** HANDLIRSCH, 1906b, p. 622 [**L. mesozoicus*; OD]. Little-known genus, based on wing fragments. Probably related to *Eocicada*. *Jur.*, Europe (Germany).
- Lithecphora** SCUDDER, 1890, p. 329 [**L. unicolor* SCUDDER, 1890, p. 329; SD CARPENTER, herein]. Little-known insect, with slender fore wing. *Oligo.*, USA (Colorado).

Lithopsis SCUDDER, 1878b, p. 773 [**L. fimbriata*; OD]. Body stout; head not produced between the eyes. Tegmina extending well beyond abdomen. SCUDDER, 1890; COCKERELL, 1921b; PONGRÁČZ, 1935; PITON, 1940a. *Eoc.*, USA (Wyoming), Europe (France, Germany).

Locrites SCUDDER, 1890, p. 323 [**L. copei* SCUDDER, 1890, p. 323; SD CARPENTER, herein]. Little-known homopteron; head large, protuberant; scutellum equiangular. HEER, 1853a. *Oligo.*, USA (Colorado); *Mio.*, Europe (Croatia).

Margaroptilon HANDLIRSCH, 1906b, p. 499 [**M. woodwardi* HANDLIRSCH, 1906b, p. 499; SD CARPENTER, herein]. Little-known wings, with numerous small maculations; possibly a fulgoroïd. BODE, 1953; EVANS, 1956. *Jur.*, England, Europe (Germany).

Mesaleuropis MARTYNOV, 1937a, p. 108 [**M. venosa*; OD]. Little-known wings. Fore wing rounded distally; pterostigma absent; M with 2 branches; CUA apparently unbranched. Hind wing about half as long as fore wing, with unbranched RS and M. *Jur.*, USSR (Tadzhik).

Meshemipteron COCKERELL, 1915, p. 476 [**M. incertum*; OD]. Little-known genus, based on small fragment of wing. *Jur.*, England.

Mesocicadella EVANS, 1956, p. 193 [**M. venosa*; OD]. Little-known genus, based on fragment of fore wing. Several parallel, oblique veins between R and wing margin; M with numerous branches. [Originally placed in the Scytinopteridae but moved to family uncertain by EVANS in 1961.] *Trias.*, Australia (Queensland).—FIG. 163,2. **M. venosa*; fore wing, $\times 3.5$ (Evans, 1956).

Mesocixiodes TILLYARD, 1922b, p. 462 [**M. termioneura*; OD]. Fore wing with SC very close to costal margin; RS present; M forking in distal part of wing, with a small, closed cell between forks. EVANS, 1956. *Trias.*, Australia (Queensland).—FIG. 162,6. **M. termioneura*; fore wing, $\times 5.2$ (Evans, 1956).

Mesococcus BECKER-MIGDISOVA, 1959a, p. 110 [**M. asiaticus*; OD]. Based on wingless form (female?); body oval; legs greatly reduced; abdomen with 9 visible segments. BECKER-MIGDISOVA, 1962b. *Trias.*, USSR (Kirghiz).—FIG. 162,7. **M. asiaticus*; whole insect, $\times 24$ (Becker-Migdisova, 1959a).

Mesodiphthera TILLYARD, 1919c, p. 873 [**M. grandis*; OD]. Little-known genus, based on small fragment of fore wing. CUA anastomosed with M basally. [Placed in Tropiduchidae by TILLYARD (1922b) and in Homoptera, family uncertain, by EVANS (1956).] *Trias.*, Australia (Queensland).—FIG. 163,3. **M. grandis*; fore wing, $\times 3.5$ (Tillyard, 1919c).

Mesojassula EVANS, 1956, p. 203 [**M. marginata*; OD]. Hind wing with costal margin with marked medial depression; M unbranched; CUA with 2 equal branches; marginal vein present. *Trias.*,






1
Tychicola2
Mesocicadella3
Mesodiphthera4
Kisylia5
Shuraboprosbole

FIG. 163. Uncertain (p. 256–259).

Australia (Queensland).—FIG. 162,4. **M. marginata*; hind wing, $\times 4.5$ (Evans, 1956).
Mesoledra EVANS, 1956, p. 211, *nom. subst. pro Mesojassus* HANDLIRSCH, 1939, p. 145, *non* TILLYARD, 1916 [**Mesojassus pachyneurus* HANDLIRSCH, 1939, p. 145; OD]. Little-known

- genus, based on incomplete wing; possibly related to Cicadellidae. *Jur.*, Europe (Germany).
- Mesoscytina** TILLYARD, 1919c, p. 871 [**M. australis*; OD]. Fore wing with SC distinct, long; RS apparently arising in very distal part of wing; M dividing at midwing and forming a closed cell at fork. Possibly related to Scytinopteridae. EVANS, 1956. *Trias.*, Australia (Queensland). —FIG. 162,5. **M. australis*; fore wing, $\times 5.2$ (Evans, 1956).
- Meuniera** PITON, 1936c, p. 1 [**M. haupti*; OD]. Little-known genus, based on fragment of fore wing. RS arising well before midwing; basal stem of M free from R. COOPER, 1941. *Eoc.*, Europe (France).
- Pachypsyche** HANDLIRSCH, 1906b, p. 623 [**Palaontina vidale* MEUNIER, 1902e, p. 9; OD]. Little-known genus. Fore wing rectangular; anterior margin straight, without nodal break; venation as in *Limacodites*. *Jur.*, Europe (Spain).
- Palaeoforda** KONONOVA, 1977, p. 588 [**P. tajmyrensis*; OD]. Little-known genus. Antennae with 6 segments. Fore wing with RS arising from distal part of pterostigma; M unbranched. Legs short. [Placed in Pemphigidae by KONONOVA but transferred to family uncertain by HEIE (1985).] *Cret.*, USSR (Asian RSFSR).
- Parafulgoridium** HANDLIRSCH, 1939, p. 138 [**Fulgoridium simplex* GEINITZ, 1880, p. 528; OD]. Little-known genus, based on poorly preserved fore wing. *Jur.*, Europe (Germany).
- Parajassus** BODE, 1953, p. 200 [**P. hattorfensis*; OD]. Little-known wing. BECKER-MIGDISOVA, 1962b. *Jur.*, Europe (Germany).
- Perissovena** RIEK, 1976a, p. 775 [**P. heidiaae*; OD]. Little-known genus, based on hind wing. *Perm.*, South Africa.
- Permocapitus** EVANS, 1943b, p. 195 [**P. globulus*; OD]. Little-known genus, based on head. Head oval, eyes globular; transverse ridge between eyes. *Perm.*, Australia (New South Wales).
- Permocephalus** EVANS, 1943a, p. 8 [**P. knightii*; OD]. Little-known insects, known only by fragments of head. *Perm.*, Australia (New South Wales).
- Permodunstantia** BECKER-MIGDISOVA, 1961c, p. 290 [**P. proboloides*; OD]. Distal fragment of fore wing; RS forked; M4 free from M3 distally. *Perm.*, USSR (Asian RSFSR). —FIG. 162,8. **P. proboloides*; fore wing, $\times 2.7$ (Becker-Migdisova, 1961c).
- Petroperon** COCKERELL, 1912b, p. 94 [**P. mirandum*; OD]. Little-known genus, based on wing fragment; possibly a fulgoroid. *Cret.*, USA (Colorado).
- Phragmatoecicossus** BECKER-MIGDISOVA, 1949b, p. 11 [**P. shurabensis*; OD]. Little-known genus, based on fragment of fore wing. Probably related to Paleontinidae. *Jur.*, USSR (Asian RSFSR).
- Plecopterus** COCKERELL, 1917h, p. 327 [**P. nebulosus*; OD]. Little-known genus, based on wing and fragments of body. [Originally placed in Trichoptera, but transferred to Homoptera, family uncertain, by BOTOSANEANU, 1981.] *Mio.*, Burma.
- Prolystra** OPPENHEIM, 1888, p. 228 [**P. lithographica*; OD]. Little-known genus, probably close to *Limacodites*. EVANS, 1956. *Jur.*, Europe (Germany).
- Prosbolopsis** MARTYNOV, 1935c, p. 19 [**P. ovalis*; OD]. Little-known insect, with reduced venation in tegmen. [Type of family Prosbolopseidae BECKER-MIGDISOVA, 1946.] EVANS, 1956; BECKER-MIGDISOVA, 1962b. *Perm.*, USSR (European RSFSR).
- Protopsyche** HANDLIRSCH, 1906b, p. 623 [**P. braueri*; OD]. Little-known genus, similar to *Limacodites*. *Jur.*, Europe (Germany).
- Reticulocicada** BECKER-MIGDISOVA, 1961c, p. 362 [**R. brachyptera*; OD]. Little-known tegmen, with coarse reticulation; possibly a fulgoroid. *Perm.*, USSR (Asian RSFSR).
- Sbenaphis** SCUDDER, 1890, p. 250 [**S. quesneli*; OD]. Little-known aphidoid genus. HEIE, 1967, 1985. *Eoc.*, Canada (British Columbia).
- Shuraboprosbole** BECKER-MIGDISOVA, 1949b, p. 23 [**S. plachutai*; OD]. Little-known genus, based on wing fragment. Basal stem of R only about half as long as R+M; RS arising well before midwing; CUA anastomosed with M for a short distance. *Jur.*, USSR (Tadzhik). —FIG. 163,5. **S. plachutai*; fore wing as preserved, $\times 2.5$ (Becker-Migdisova, 1949b).
- Stenoglyphis** EVANS, 1947b, p. 432 [**S. kimblensis*; OD]. Little-known genus, possibly related to Scytinopteridae. EVANS, 1956. *Perm.*, Australia (New South Wales).
- Tingiopsis** BECKER-MIGDISOVA, 1953c, p. 461 [**T. reticulata*; OD]. Little-known genus, based on incomplete fore wing with fine reticulation. [Originally placed in Tingidae (Heteroptera) but transferred to Homoptera, probably Cercopidae, by EVANS (1957).] *Trias.*, USSR (Tadzhik).
- Tipuloidea** WIELAND, 1925, p. 23 [**T. rhaetica*; OD]. Little-known genus, based on fore wing. Costal margin arched; SC apparently absent; RS arising before midwing; closed median cell very small. [Originally placed in order Diptera.] EVANS, 1956. *Trias.*, Argentina. —FIG. 162,2. **T. rhaetica*; fore wing, $\times 2$ (Evans, 1956).
- Triassoaphis** EVANS, 1956, p. 238 [**T. cubitus*; OD]. Little-known genus, based on wing fragment. [Originally placed in Aphididae but transferred to Aphidoidea, family uncertain, by BECKER-MIGDISOVA & AIZENBERG (1962).] RICHARDS, 1966; HEIE, 1967, 1981; SHAPOSHNIKOV, 1979a. *Trias.*, Australia (Queensland).
- Triassocotis** EVANS, 1956, p. 194 [**T. australis*; OD]. Little-known genus, based on distal half of tegmen. Tegmen narrow; R with 4 branches;

- RS unbranched; M with 4 branches and a cell included between M1+2 and M3+4. [Originally placed in Scytinopteridae but transferred to family uncertain by EVANS (1961).] *Trias.*, Australia (Queensland). — FIG. 162,3. **T. australis*; fore wing, $\times 4.5$ (Evans, 1956).
- Triassojassus** TILLYARD, 1919c, p. 887 [**T. proavittus*; OD]. Little-known genus, based on incomplete tegmen. Costal margin unusually convex; RS unbranched; M with 5 branches. [Originally placed in the Jassidae, but EVANS transferred first (1956) to the Chilocyclidae and later (1961) to family uncertain.] *Trias.*, Australia (New South Wales).
- Turatanovia** BECKER-MIGDISOVA, 1949b, p. 21 [**T. karatavia*; OD]. Little-known genus, based on distal fragment of fore wing. BECKER-MIGDISOVA, 1962b. *Jur.*, USSR (Kazakh). — FIG. 162,10. **T. karatavia*; fore wing as preserved, $\times 2$ (Becker-Migdisova, 1962b).
- Tychicola** BECKER-MIGDISOVA, 1952, p. 181 [**T. longipenna*; OD]. Little-known genus, based on incomplete fore wing. Wing apparently long and narrow; RS long and parallel to R2. *Perm.*, USSR (Asian RSFSR). — FIG. 163,1. **T. longipenna*; fore wing, $\times 5$ (Becker-Migdisova, 1962b).
- p. 494 (type, *E. liasinus*). Fore wing with rounded apex; clavus broad, nearly triangular; M with 3 branches. BECKER-MIGDISOVA, 1962b; POPOV & WOOTTON, 1977. *Jur.*, Europe (Germany). — FIG. 164,8. *Progonocimex*; a, **P. jurassicus*, dorsal view; b, *P. liasinus* (HANDLIRSCH), fore wing, both $\times 9$ (Popov & Wootton, 1977).
- Actinoscytina** TILLYARD, 1926a, p. 18 [**A. belmontensis*; OD] [= *Pseudiprivia* HANDLIRSCH, 1939, p. 17 (type, *P. ala*)]. Little-known genus. Tegmen similar to that of *Progonocimex*, but more slender, anterior margin less curved; SC curving directly toward anterior margin of wing. EVANS, 1956; POPOV & WOOTTON, 1977. *Perm.*, Australia (New South Wales). — FIG. 164,5. **A. belmontensis*; tegmen, $\times 8$ (Evans, 1956).
- Archicercopis** HANDLIRSCH, 1939, p. 142 [**A. falcata*; OD]. Anterior margin of fore wing strongly convex basally; precostal area broad; wing apex pointed and directed anteriorly. EVANS, 1956; BECKER-MIGDISOVA, 1962b; POPOV & WOOTTON, 1977. *Jur.*, Europe (Germany). — FIG. 164,6. **A. falcata*; fore wing, $\times 13$ (Popov & Wootton, 1977).
- Cicadocoris** BECKER-MIGDISOVA, 1958, p. 62 [**C. kuliki*; OD]. Tegmen with smoothly curved anterior margin; M with 3 branches; M3+4 unbranched. EVANS, 1961; POPOV, 1982. *Trias.*, USSR (Kirghiz). — FIG. 164,9. **C. kuliki*; respiration, $\times 10$ (Becker-Migdisova, 1958).
- Eocercopis** HANDLIRSCH, 1939, p. 142 [**E. ancyloptera*; OD] [= *Cercoprisca* HANDLIRSCH, 1939, p. 143 (type, *C. similis*); *Cercopinus* HANDLIRSCH, 1939, p. 143 (type, *C. ovalis*)]. Fore wing with very convex and thickened costal margin; apex pointed; clavus broad and nearly triangular. EVANS, 1956; BECKER-MIGDISOVA, 1958; POPOV & WOOTTON, 1977. *Jur.*, Europe (Germany). — FIG. 164,7. **E. ancyloptera*; fore wing, $\times 13$ (Popov & Wootton, 1977).
- Heterojassus** EVANS, 1961, p. 23 [**H. membranaceus*; OD]. Tegmen oval; SC and R terminating on costal margin near level of midwing. *Trias.*, Australia (Queensland). — FIG. 164,4. **H. membranaceus*; fore wing, $\times 19$ (Evans, 1961).
- Heteroscytina** EVANS, 1956, p. 245 [**H. tillyardi*; OD]. Fore wing narrowed apically, much as in *Actinoscytina*, but costal area narrower and crossveins forming a more nearly complete transverse series. WOOTTON, 1963. *Trias.*, Australia (Queensland).
- Hexascytina** WOOTTON, 1963, p. 250 [**H. transecta*; OD]. Little-known genus, apparently similar to *Progonocimex*, based on incomplete tegmen. SC diverging from stem R near midwing at almost a 90° angle; anterior margin of tegmen distinctly convex. *Trias.*, Australia (Queensland).
- Microscytinella** WOOTTON, 1963, p. 251 [**M. radians*; OD]. Little-known genus, based on small

Suborder HETEROPTERA

Latreille, 1810

[Heteroptera LATREILLE, 1810, p. 433]

Fore wing typically with the proximal part strongly coriaceous and the distal part membranous, forming a hemelytron; wings usually held flat over abdomen at rest. *Perm.*—*Holo.*

Family PROGONOCIMICIDAE Handlirsch, 1906

[Progonocimicidae HANDLIRSCH, 1906b, p. 493] [= Eocimicidae HANDLIRSCH, 1906b, p. 494; Actinoscytinidae EVANS, 1956, p. 244; Cicadocoridae BECKER-MIGDISOVA, 1958, p. 60]

Small species, dorsoventrally flattened; pronotum distinctly broader than long; fore wing apparently of uniform texture; veins RS and M coalesced basally; SC apparently coalesced with stem of R basally, diverging toward costal margin near midwing; M with 2 to 4 branches; CUA with 2 to 3 branches. [Placed by POPOV (1980a) in suborder Peloridiina, along with the Peloridiidae (recent).] *Perm.*—*Jur.*

Progonocimex HANDLIRSCH, 1906b, p. 494 [**P. jurassicus*; OD] [= *Eocimex* HANDLIRSCH, 1906b,

Microscytinella WOOTTON, 1963, p. 251 [**M. radians*; OD]. Little-known genus, based on small




FIG. 164. Progonocimicidae (p. 259–260).




FIG. 165. Archegocimicidae, Enicocoridae, and Cuneocoridae (p. 261–262).

fragment of tegmen. *Trias.*, Australia (Queensland).

Olgamartynovia BECKER-MIGDISOVA, 1958, p. 63 [*O. turanica*; OD]. Tegmen as in *Cicadocoris*, but fork of M1+2 longer. POPOV, 1982. *Trias.*, USSR (Kirghiz).—FIG. 164,1. *O. complexa* POPOV; tegmen, $\times 14$ (Popov, 1982).

Platyscytinella EVANS, 1956, p. 245 [**P. paradoxa*; OD]. Tegmen shaped as in *Actinoscytina*; SC absent; M1 continuing the nearly straight line of stem M; clavus unknown. WOOTTON, 1963. *Trias.*, Australia (New South Wales).—FIG. 164,3. **P. paradoxa*; fore wing, $\times 15$ (Wootton, 1963).

Triassocoecus EVANS, 1963, p. 22 [**T. chinai*; OD]. Little-known genus. Tegmen broader than in *Actinoscytina*; pronotum with large lateral lobes. *Trias.*, Australia (New South Wales).—FIG. 164,10. **T. chinai*; dorsal view, $\times 10$ (Evans, 1963).

Triscytina EVANS, 1956, p. 246 [**T. rotundata*; OD]. Similar to *Actinoscytina* but tegmen much broader; costal margin nearly straight; apex evenly rounded. *Trias.*, Australia (New South Wales).—FIG. 164,2. **T. rotundata*; tegmen, $\times 16$ (Evans, 1956).

Family ARCHEGOCIMICIDAE Handlirsch, 1906

[Archegocimicidae HANDLIRSCH, 1906b, p. 493] [=Eonabidae HANDLIRSCH, 1920, p. 207; Diatillidae HANDLIRSCH, 1920, p. 210]

Small Heteroptera of uncertain relationships. Body dorsoventrally flattened; pronotum coarsely warty. Fore wing with apex rounded; clavus narrow; costal margin

strongly sclerotized; veins SC, R, and M coalesced for about two-fifths length of wing.

Jur.

Archegocimex HANDLIRSCH, 1906b, p. 493 [**A. geinitzi*; OD] [=Eonabis HANDLIRSCH, 1920, p. 207 (type, *E. primitiva*); *Archegocoris* HANDLIRSCH, 1939, p. 114 (type, *A. liadis*)]. Radial complex of fore wing (R and RS) with 3 branches terminating on anterior margin of wing. POPOV & WOOTTON, 1977. *Jur.*, Europe (Germany).—FIG. 165,1. **A. geinitzi*; dorsal view, $\times 14$ (Popov & Wootton, 1977).

Anosmus HANDLIRSCH, 1939, p. 115 [**A. spilopteris*; OD]. Costal area of fore wing narrow; radial complex branched. POPOV & WOOTTON, 1977. *Jur.*, Europe (Germany).

Corynecoris BODE, 1953, p. 132 [**C. semigranulatus*; OD]. Little-known genus, based on poorly preserved specimen. [Family assignment doubtful.] POPOV & WOOTTON, 1977. *Jur.*, Europe (Germany).

Diatillus HANDLIRSCH, 1920, p. 210 [**D. debilis*; OD]. Little-known genus with archegocimicid venation. POPOV & WOOTTON, 1977. *Jur.*, Europe (Germany).

Ensphingocoris BODE, 1953, p. 139 [**E. praerotundatus*; OD]. Little-known genus, based on poorly preserved specimen lacking wings; body apparently that of a large archegocimicid. POPOV & WOOTTON, 1977. *Jur.*, Europe (Germany).

Entomecoris BODE, 1953, p. 134 [**A. minor*; OD]. Fore wing with common stem of SC, R, and M curving away from costal margin; wing differentiated into corium and membrane. POPOV & WOOTTON, 1977. *Jur.*, Europe (Germany).

Eurynotis BODE, 1953, p. 134 [**E. incisus*; OD].

Similar to *Somatocoris*, but radial complex with 2 branches. BECKER-MIGDISOVA, 1962b; POPOV & WOOTTON, 1977. *Jur.*, Europe (Germany).

Macropterocoris BODE, 1953, p. 138 [**M. obtusus*; OD]. Little-known genus; head and thorax resembling those of the Archegocimicidae. POPOV & WOOTTON, 1977. *Jur.*, Europe (Germany).

Progonocoris HANDLIRSCH, 1939, p. 115 [**P. pictus*; OD]. Similar to *Anosmus*, but radial complex of fore wing apparently unbranched; costal area long. BECKER-MIGDISOVA, 1962b; POPOV & WOOTTON, 1977. *Jur.*, Europe (Germany).

Somatocoris BODE, 1953, p. 141 [**S. conservatus*; OD]. Similar to *Archegocimex* but smaller; radial complex (R and RS) with 3 branches. BECKER-MIGDISOVA, 1962b; POPOV & WOOTTON, 1977. *Jur.*, Europe (Germany).

Family ENICOCEPHALIDAE Stål, 1858

[Enicocephalidae STÅL, 1858, p. 81]

Similar to the Reduviidae, but head constricted behind eyes; rostrum with 4 segments; fore wings entirely membranous, with longitudinal veins but few crossveins. *Mio.*—*Holo.*

Enicocephalus WESTWOOD, 1838, p. 22. *Holo.*
Disphaerocephalus COCKERELL, 1917g, p. 361 [**D. constrictus*; OD]. Little-known genus of small, elongate species, with long, thin legs and antennae; body with long, fine pubescence; hind legs long and narrow; tarsi 1-2-2; wings unknown. [Genus based on nymph and adult male.] ŠTYS, 1969. *Mio.*, Burma.

Paenicotechys ŠTYS, 1969, p. 353 [**Enicocephalus fossilis* COCKERELL, 1916a, p. 135; OD]. Similar to *Aenicopechys* (recent), but posterior margin of pronotum excised; eyes contiguous dorsally; middle tarsi with 2 segments. ŠTYS, 1969. *Mio.*, Burma.

Family ENICOCORIDAE Popov, 1980

[Enicocoridae POPOV, 1980a, p. 50]

Apparently related to the Enicocephalidae. Head short, rostrum thick, curved. Tegmen entirely membranous, clavus and corium not differentiated; radial-medial and cubital-anal sectors of veins widely separated at base; veins nearly parallel distally. Legs thin, cursorial. *Cret.*

Enicocoris POPOV, 1980a, p. 50 [**E. manlaicus*; OD]. Head prognathous; pronotum transverse; scutellum much narrower than pronotum; subcostal area of tegmen wide. *Cret.*, Mongolia. —FIG. 165,2 **E. manlaicus*; dorsal view, ×9 (Popov, 1980a).

Family DIPSOCORIDAE Dohrn, 1859

[Dipsocoridae DOHRN, 1859, p. 36]

Similar to the Saldidae, but third antennal segment not thickened at base. *Mio.*—*Holo.*

Dipsocoris HALIDAY, 1855, fig. 61. *Holo.*

Ceratocombus SIGNORET, 1852, p. 542. WYGODZINSKY, 1959. *Mio.*, Mexico (Chiapas)—*Holo.*

Family CUNEOCORIDAE Handlirsch, 1920

[Cuneocoridae HANDLIRSCH, 1920, p. 208]

Small insects. Fore wings reaching end of abdomen and overlapped distally; pronotum wider than long; scutellum triangular. Fore wing not clearly differentiated into corium and membrane; veins M and CU branched. *Jur.*

Cuneocoris HANDLIRSCH, 1920, p. 208 [**C. geinitzi*; OD]. M and CU each with 2 branches. POPOV & WOOTTON, 1977. *Jur.*, Europe (Germany). — FIG. 165,3. **C. geinitzi*; dorsal view, ×22 (Popov & Wootton, 1977).

Family GERRIDAE Leach, 1815

[Gerridae LEACH, 1815, p. 123]

Body slender; rostrum with 4 segments; fore wings without differentiation of corium, membrane, or clavus; posterior femora extending well beyond end of abdomen; claws ante-apical. Semiaquatic. ANDERSEN, 1982b. *Eoc.*—*Holo.*

Gerris FABRICIUS, 1794, p. 187. COCKERELL, 1909j; HANDLIRSCH, 1910b; THÉOBALD, 1937a. *Oligo.*, USA (Colorado), Canada (British Columbia), Europe (France)—*Holo.*

Metrobates UHLER, 1871, p. 108. SCUDDER, 1890. *Oligo.*, USA (Colorado)—*Holo.*

Telmatrechus SCUDDER, 1890, p. 351 [**Hygrotrechus stali* SCUDDER, 1879a, p. 183b; SD CARPENTER, herein]. Eyes not prominent; first antennal segment only a little longer than second; thorax relatively short; legs very long, with the tibiae equal in length to femora of same leg. *Eoc.*, USA (Wyoming); *Mio.*, Canada (British Columbia).

Family HYDROMETRIDAE Stephens, 1829

[Hydrometridae STEPHENS, 1829, p. 352]

Very slender species; head long and narrow but widened distally; antennae with 4 (rarely 5) segments; legs very long and slender, claws

apical; rostrum with 3 segments; tegmen with corium and membrane. *Eoc.*—*Holo.*

Hydrometra LATREILLE, 1796, p. 86. *Holo.*

Eocenometra ANDERSEN, 1982a, p. 91 [**E. danica*; OD]. Similar to *Bacillometra* (recent) and *Hydrometra* (recent), but first antennal segment much longer than second; thorax relatively short and robust. ANDERSEN, 1982b. *Eoc.*, Europe (Denmark).

Family VELIIDAE

Amyot & Serville, 1843

[Veliiidae AMYOT & SERVILLE, 1843, p. 418]

Similar to the Gerridae, but rostrum with 3 segments; posterior femora shorter, extending very little beyond end of abdomen at most. *Oligo.*—*Holo.*

Velia LATREILLE, 1804, p. 270. MEUNIER, 1914a. *Oligo.*, Europe (France)—*Holo.*

Palaeovelina SCUDDER, 1890, p. 349 [**P. spinosa*; OD]. Similar to *Microvelia* (recent). Head small, recessed to level of eyes in emarginate prothorax; hind legs very short, reaching only tip of abdomen; femora and tibiae of equal lengths; hind tibiae with long spines distally. *Oligo.*, USA (Colorado).

Stenovelina SCUDDER, 1890, p. 349 [**S. nigra*; OD]. Similar to *Palaeovelina*, but hind tibiae without long spines distally. *Oligo.*, USA (Colorado).

Family NOTONECTIDAE

Latreille, 1802

[Notonectidae LATREILLE, 1802a, p. 253]

Aquatic species, similar to the Naucoridae, but forelegs raptorial, and hind tarsi without claws. *Jur.*—*Holo.*

Notonecta LINNÉ, 1758, p. 439. PITON, 1942; LAUCK, 1960; POPOV, 1964; MARTINI, 1971. *Oligo.*, USA (Colorado), Europe (Germany); *Mio.*, Europe (France)—*Holo.*

Anisops SPINOLA, 1837, p. 58. DEICHMÜLLER, 1881; STYS & RIHA, 1975a. *Oligo.*, Europe (Czechoslovakia)—*Holo.*

Asionecta POPOV in BECKER-MIGDISOVA & POPOV, 1963, p. 78 [**A. curtipes*; OD]. Similar to *Notonecta* but with first segment of front and middle legs very short. *Jur.*, USSR (Kazakh). —FIG. 166,2. **A. curtipes*; ventral view, $\times 5$ (Popov in Becker-Migdisova & Popov, 1963).

Clematina POPOV, 1964, p. 66 [**Notonecta primaeva* HEYDEN, 1859a, p. 11; OD]. Little-known genus, apparently related to *Clypostemma*. POPOV, 1971; STYS, 1973. *Oligo.*, Europe (Germany).

Clypostemma POPOV, 1964, p. 64 [**C. xyphiale*; OD]. Species of moderate size and of uncertain

relationship within the family. Rostrum with 4 segments; tarsi of all legs with 2 segments. POPOV, 1971; STYS, 1973. *Cret.*, USSR (Asian RSFSR).

Enithares SPINOLA, 1837, p. 60. A nymph is only fossil record. [Generic assignment uncertain.] STYS & RIHA, 1975a. *Oligo./Mio.*, Europe (Czechoslovakia)—*Holo.*

Liadonecta POPOV, 1971, p. 172 [**L. tomiensis*; OD]. Little-known genus, based on nymph. Body elongate-oval, head transverse; hind tibiae and tarsi of uniform width. *Jur.*, USSR (Asian RSFSR). —FIG. 166,3. **L. tomiensis*; dorsoventral view, $\times 14$ (Popov, 1971).

Nepidium WESTWOOD, 1854, p. 396 [**N. stolonae*; OD]. Little-known genus, based on poorly preserved specimen. [Put in Naucoridae by HANDLIRSCH (1906b) and in Notonectidae by POPOV (1971).] *Jur.*, England.

Notonectites HANDLIRSCH, 1906b, p. 639 [**Notonecta elterleini* DEICHMÜLLER, 1886, p. 64; OD]. Little-known genus, apparently close to *Notonecta* and *Anisops* (recent). POPOV, 1964, 1971; STYS & RIHA, 1975a. *Jur.*, Europe (Germany).

Pelonecta POPOV, 1971, p. 170 [**P. solnhofeni*; OD]. Body elongate-oval, widest near base of abdomen; hind tibiae shorter than femora or tarsi; femora thickened, strongly developed. *Jur.*, Europe (Germany). —FIG. 166,1. **P. solnhofeni*; ventral view, $\times 2.2$ (Popov, 1971).

Soevenia STATZ, 1950b, p. 63 [**Notonecta beydeni* DEICHMÜLLER, 1881, p. 328; OD]. Similar to *Anisops*; clypeus fused to frons. Body structure little known. *Oligo.*, Europe (Germany, Czechoslovakia).

Family SCAPHOCORIDAE

Popov, 1968

[Scaphocoridae POPOV, 1968, p. 106]

Body oval; head hypognathous; pronotum large, covering scutellum; tegmen with membrane; hind legs relatively short; tarsi with a single segment and dense hairs. Probably related to the Naucoridae. *Jur.*

Scaphocoris POPOV, 1968, p. 106 [**S. notatus*; OD]. Head strongly transverse from above; scutellum very small, triangular; clavus with distinct anal veins; membrane present; hind tarsi shorter than tibiae. *Jur.*, USSR (Kazakh).

Family NAUCORIDAE Leach, 1815

[Naucoridae LEACH, 1815, p. 123] [=Aphlebororidae HANDLIRSCH, 1906b, p. 494; Apopnidae HANDLIRSCH, 1920, p. 209]

Antennae four-segmented, shorter than head; fore wing membrane without veins; forelegs raptorial; tarsi with more than one segment; hind tarsi with claws. *Jur.*—*Holo.*




FIG. 166. Notonectidae (p. 263).

- Naucoris** GEOFFROY, 1762, p. 473. HEER, 1853a; PITON & THÉOBALD, 1937; STATZ, 1950b. *Oligo.*, Europe (Germany); *Mio.*, Europe (France, Croatia)—*Holo.*
- Aidium** POPOV, 1968, p. 103 [**A. pleurale*; OD]. Anterior margin of pronotum concave; scutellum very large; tegmen, except for clavus, without venation; embolium absent; clavus narrow, hind legs relatively short. POPOV, 1971. *Jur.*, USSR (Kazakh).
- Angaronecta** POPOV, 1971, p. 146 [**A. longirostris*; OD]. Rostrum very long, reaching hind margin of prothorax; forelegs short; femora of all legs thickened; hind tarsi with single segment. *Jur.*, USSR (Asian RSFSR). — FIG. 167,3. **A. longirostris*; ventral view, $\times 3.7$ (Popov, 1971).
- Aphlebocoris** HANDLIRSCH, 1906b, p. 495 [**A. nana*; OD]. Fore wing not differentiated into corium and membrane; clavus narrow, nearly quadrilateral. POPOV & WOOTTON, 1977. *Jur.*, Europe (Germany). — FIG. 167,6. *A. punctata* HANDLIRSCH; fore wing, $\times 11$ (Popov & Wootton, 1977).
- Apopnus** HANDLIRSCH, 1920, p. 209 [**A. magniclavus*; OD]. Little-known genus. Fore wing differentiated into corium and membrane; costal margin convex; clavus broad and triangular. POPOV & WOOTTON, 1977. *Jur.*, Europe (Germany).
- Diplonychus** LAPORTE, 1832, p. 18. HEER, 1853a. *Mio.*, Europe (Croatia)—*Holo.*
- Heleonaucoris** POPOV, 1971, p. 149 [**H. maculipennis*; OD]. Clavus of moderate size; embolium narrow and developed only at base of tegmen; border between corium and membrane indistinct; corium spotted. *Jur.*, USSR (Kirghiz). — FIG. 167,1. **H. maculipennis*; tegmen, $\times 4.2$ (Popov, 1971).
- Liadonaucoris** POPOV, 1971, p. 144 [**L. robdendorfi*; OD]. Tegmen longer than abdomen; clavus longer than scutellum; vein R present on tegmen. *Jur.*, USSR (Kirghiz). — FIG. 167,2. **L. robdendorfi*; $\times 5.5$ (Popov, 1971).
- Nectodes** POPOV, 1968, p. 105 [**N. maculatus*; OD]. Little-known genus, based on tegmen. Clavus large and broad; embolium distinct, extending for half length of corium; membrane large. *Jur.*, USSR (Kazakh).
- Nectonaucoris** POPOV, 1968, p. 104 [**N. lariversi*; OD]. Anterior margin of pronotum straight; tegmen without veins; embolium absent; clavus narrow; hind legs relatively short. *Jur.*, USSR (Kazakh).
- Sphaerodemopsis** HANDLIRSCH, 1906b, p. 543 [**Sphaerodema jurassicum* OPPENHEIM, 1888, p. 235; OD]. Tegmen strongly sclerotized; clavus usually long and heavily sclerotized. POPOV, 1971. *Jur.*, Europe (Germany). — FIG. 167,5. **S. jurassicum*; dorsal view, $\times 2.5$ (Popov, 1971).




FIG. 167. Naucoridae and Mesotrepidae (p. 264–265).

Family MESOTREPHIDAE
Popov, 1971

[Mesotrepidae POPOV, 1971, p. 160]

Small species, related to the Scaphocoridae. Pronotum transverse, convex, elliptical, completely covering head from above. *Cret.*

Mesotrepes POPOV, 1971, p. 160 [**M. striata*; OD]. Scutellum very small; tegmen with only one vein, extending along most of costal margin.

Hind legs relatively short; tibiae and tarsi thin. *Cret.*, USSR (Kazakh). — FIG. 167, 4. **M. striata*; dorsoventral view, $\times 20$ (Popov, 1971).

Family BELOSTOMATIDAE
Leach, 1815

[Belostomatidae LEACH, 1815, p. 123]

Similar to the Nepidae, but antennae with 4 segments; posterior legs adapted for swimming, the tibiae flattened; aquatic. *Jur.-Hol.*




FIG. 168. Belostomatidae (p. 266–267).

Belostoma LATREILLE, 1807, p. 144. GERMAR, 1837. *Oligo.*, Europe (Germany)—*Holo.*

Belostomates SCHÖBERLIN, 1888, p. 61 [**Belostomum speciosa* HEER, 1865, p. 303; OD]. Little-known genus, with very broad front femora. HEER, 1865. *Mio.*, Europe (Germany).—FIG. 168,2. **B. speciosa*; dorsal view, $\times 0.6$ (Heer, 1865).

Lethocerus MAYR, 1853, p. 17. ŘIHA & KUKALOVÁ, 1967; POPOV, 1971. *Oligo.*, USSR (Asian RSFSR); *Mio.*, Europe (Czechoslovakia)—*Holo.*

Mesonepa HANDLIRSCH, 1906b, p. 637 [**Nepa primordialis* GERMAR, 1839, p. 206; SD POPOV, 1971, p. 116]. Similar to *Belostoma*, but fore

wing with much larger membranous area; fore tarsi with one segment. POPOV, 1971. *Jur.*, Europe (Germany).—FIG. 168,4. **M. primordialis*; dorsal view, $\times 2$ (Popov, 1971).

Scarabaeides GERMAR, 1839, p. 218 [**S. deperditus*; OD] [= *Mesobelostomum* HAASE, 1890a, p. 21, obj. J. Similar to *Lethocerus* (recent), but fore wing with M and R widely separated and remote from costal margin; membranous area of wing without venation. OPPENHEIM, 1888; POPOV, 1971. *Jur.*, Europe (Germany).—FIG. 168,3. **S. deperditus*; ventral view, $\times 1.2$ (Popov, 1971).

Stygonepa POPOV, 1971, p. 119 [**S. foersteri*; OD].

Related to *Belostoma*, but pronotum more transverse; fore wing with vestiges of R, M, and CU; hind tibiae and one-segmented tarsi forming broad lobes. *Jur.*, Europe (Germany).—FIG. 168,1. **S. foersteri*; dorsal view, $\times 2$ (Popov, 1971).

Family NEPIDAE Latreille, 1802

[Nepidae LATREILLE, 1802a, p. 252]

Antennae with 3 segments, shorter than head; membrane of fore wings reticulate; hind legs adapted for walking; tibiae not flattened; aquatic. *Oligo.*—*Holo.*

Nepa LINNÉ, 1758, p. 440. HEER, 1853a; HUNGERFORD, 1932. *Oligo.*, USA (Colorado); *Mio.*, Europe (Germany)—*Holo.*

Family SHURABELLIDAE

Popov, 1971

[Shurabellidae POPOV, 1971, p. 121]

Small species, related to the Corixidae. Pronotum transverse, sculptured; tegmen broad; hind legs relatively slender. *Jur.*

Shurabella BECKER-MIGDISOVA, 1949b, p. 28 [**S. lepyroniopsis*; OD] [= *Coleopteropsis* BECKER-MIGDISOVA, 1949b, p. 31 (type, *C. dolichoptera*)]. Pronotum about three times as wide as long; tegmen strongly sclerotized; vestiges of SC, R, and M present on tegmen. *Jur.*, USSR (Kirghiz).—FIG. 169,4. **S. lepyroniopsis*; dorsoventral view, $\times 10$ (Popov, 1971).

Family CORIXIDAE Leach, 1815

[Corixidae LEACH, 1815, p. 124]

Head not inserted into prothorax; antennae shorter than head; fore tarsi consisting of only one spatulate segment. *Jur.*—*Holo.*

Corixa GEOFFROY, 1762, p. 477. HEER, 1853a; SCUDDER, 1890; SCHLECHTENDAL, 1894. *Oligo.*, USA (Colorado), Europe (Germany); *Mio.*, Europe (Germany)—*Holo.*

Archaeorixa POPOV, 1968, p. 101 [**A. lata*; OD]. Pronotum transverse; corium of tegmen with distinct venation; SC coalesced with R, M, and CU for varying lengths, finally terminating on costal margin; embolium absent. POPOV, 1971. *Jur.*, USSR (Kazakh).

Baisocorixa POPOV, 1966, p. 99 [**B. jaczewskii*; OD]. Similar to *Corixa*, but veins R, M, and CU more strongly developed; head narrow; eyes small; eighth abdominal tergite well developed. *Jur./Cret.*, USSR (Asian RSFSR).

Diacorixa POPOV, 1971, p. 137 [**D. miocaenica*;

OD]. Similar to *Sigara* (recent) but with deep furrow along entire length of pronotum; vein CU weakly formed. *Mio.*, USSR (Kirghiz).

Diapherinus POPOV, 1966, p. 97 [**D. ornatipennis*; OD]. Little-known genus, based on tegmen. SC, R, M, and CU visible on corium; anal veins clear on clavus; embolium weakly developed. *Jur./Cret.*, USSR (Asian RSFSR).—FIG. 169,2. **D. ornatipennis*; tegmen, $\times 5$ (Popov, 1966).

Gazimuria POPOV, 1971, p. 130 [**G. scutellata*; OD]. Elongate species. Antennae with 4 segments; pronotum not more than three times wider than its length; tegmen with veins R, M, and CU; hind legs densely covered with hairs. *Jur.*, USSR (Asian RSFSR).—FIG. 169,5. **G. scutellata*; dorsoventral view, $\times 6$ (Popov, 1971).

Ijanecta POPOV, 1971, p. 132 [**I. angarica*; OD]. Pronotum well developed; scutellum small; fore margin of wing with wide embolium; all veins apparently absent. *Jur.*, USSR (Asian RSFSR).—FIG. 169,1. **I. angarica*; dorsal view, $\times 1$ (Popov, 1971).

Karataviella BECKER-MIGDISOVA, 1949b, p. 25 [**K. brachyptera*; OD]. Pronotum twice as wide as long; only vein 1A on clavus. *Jur.*, USSR (Kazakh).—FIG. 169,3. **K. brachyptera*; dorsal view, $\times 7$ (Popov, 1971).

Mesosigara POPOV, 1971, p. 129 [**M. kryshstofovichi*; OD]. Similar to *Baisocorixa*, but fore wing with R coalesced with SC for its entire length; M fused at base with CU. *Cret.*, USSR (Asian RSFSR).—FIG. 169,7. **M. kryshstofovichi*; lateral view, $\times 12$ (Popov, 1971).

Sigaretta POPOV, 1971, p. 136 [**Corixa florissantiella* COCKERELL, 1906e, p. 209; OD]. Pronotum large but covering only part of the scutellum; tegmen with well-developed embolium rim; anal vein present on clavus. *Oligo.*, USA (Colorado).—FIG. 169,6. **S. florissantiella* (COCKERELL); dorsal view, $\times 9$ (Popov, 1971).

Family ARADIDAE Brullé, 1835

[Aradidae BRULLÉ, 1835, p. 326]

Body strongly flattened; head porrect; antennae and rostrum with 4 segments; clavus narrowed apically; wing membrane with few or no veins; abdomen broader than wings; tarsi with 2 segments. *Oligo.*—*Holo.*

Aradus FABRICIUS, 1803, p. 116. GERMAR & BERENDT, 1856; USINGER, 1941; POPOV, 1978. *Oligo.*, Europe (Baltic); *Mio.*, Europe (Croatia)—*Holo.*

Calisius STÅL, 1858, p. 67. USINGER, 1941. *Oligo.*, Europe (Baltic)—*Holo.*

Mezira AMYOT & SERVILE, 1843, p. 305. USINGER, 1941. *Oligo.*, Europe (Baltic); *Mio.*, Europe (Croatia)—*Holo.*




FIG. 169. Shurabellidae and Corixidae (p. 267).

Family SALDIDAE
Amyot & Serville, 1843

[Saldidae AMYOT & SERVILLE, 1843, p. xlix]

Head shorter than thorax and scutellum; antennae four-segmented, longer than head; third antennal segment thickened at base; rostrum three-segmented; fore wings without reticulate cells, but with 4 or 5 long, closed cells; corium with an embolium; forelegs not raptorial. *Jur.*—*Holo.*

Salda FABRICIUS, 1803, p. 113. GERMAR & BERENDT, 1856. *Oligo.*, Europe (Baltic)—*Holo.*
Oligosaldina CARPENTER, herein [**O. rottensis* STATZ

& WAGNER, 1950, p. 101; OD]. Fore wing similar to *Chiloxanthus* (recent) but with cells of membrane nearly the same length. [The original generic name, *Oligosaldina*, was a *nomen nudum* (STATZ & WAGNER, 1950).] *Oligo.*, Europe (Germany).

Saldonia POPOV, 1973, p. 704 [**S. rasnitsyni*; OD]. Pronotum transverse; RS close to front margin of tegmen; membrane not present on tegmen; scutellum small, shorter than claval suture. *Jur.*, USSR (Asian RSFSR).

Family COREIDAE Leach, 1815

[Coreidae LEACH, 1815, p. 121]

Head much narrower and shorter than prothorax; antennae longer than head, with




FIG. 170. Coreidae, Alydidae, Lygaeidae, Pachymeridiidae, and Miridae (p. 269–275).

4 segments; fore wing not reticulate, its membrane with many longitudinal veins, often uniting; body stout, legs thick. *Jur.*—*Holo.*

Coreus FABRICIUS, 1794, p. 120. THÉOBALD, 1937a. *Oligo.*, Europe (France)—*Holo.*

Achrestocoris SCUDDER, 1890, p. 413 [**A. cinerarius*; OD]. Tegmen with large rhomboidal cell at apex of corium. *Oligo.*, USA (Colorado).

Anasa AMYOT & SERVILLE, 1843, p. 209. SCUDDER, 1890. *Oligo.*, USA (Colorado)—*Holo.*

Berytopsis HEER, 1853a, p. 54 [**B. femoralis*; OD]. Little-known coreid, apparently related to *Berytus* (recent). *Mio.*, Europe (Germany).

Corizus FALLÉN, 1814, p. 8. SCUDDER, 1890; COCKERELL, 1926a. *Oligo.*, USA (Colorado); *Tert.* (epoch unknown), Argentina—*Holo.*

Harmostites HEER, 1853a, p. 49 [**H. oeningensis*;

OD]. Little-known genus, based on poorly preserved specimen. *Mio.*, Europe (Germany).

Heeria SCUDDER, 1890, p. 430 [**H. gulosa*; SD HANDLIRSCH, 1907, p. 1049]. Similar to *Arenocoris* (recent) but with second and third antennal segments unequal. *Oligo.*, USA (Colorado).

Hypselonotus HAHN, 1833, p. 186. HEER, 1853a. *Mio.*, Europe (Germany)—*Holo.*

Jadera STÅL, 1860, p. 59. COCKERELL, 1909j. *Eoc.*, USA (Colorado)—*Holo.*

Karatavocoris BECKER-MIGDISOVA, 1962b, p. 222 [**K. asiatica*; OD]. Head much narrower than pronotum; first antennal segment short, not extending beyond apex of head; femora without spines. *Jur.*, USSR (Kazakh). — FIG. 170, 5. **K. asiatica*; ventral view, $\times 6$ (Becker-Migdisova, 1962b).

Leptoscelis LAPORTE, 1832, p. 31. HEYDEN, 1858.

- Palaeocoris** HEER, 1853a, p. 46 [**P. spectabilis*; OD]. Little-known genus, apparently related to *Acanthosoma* (recent). *Mio.*, Europe (Croatia).
- Phthinocoris** SCUDDER, 1890, p. 414 [**P. colligatus*; SD HANDLIRSCH, 1907, p. 1049]. Similar to *Achrestocoris*, but thorax much longer. *Oligo.*, USA (Colorado).
- Piezocoris** SCUDDER, 1890, p. 416 [**P. peritus* SCUDDER, 1890, p. 416; SD CARPENTER, herein]. Similar to *Phthinocoris*, but head large, one-half to two-thirds width of thorax. *Oligo.*, USA (Colorado).
- Spartocera** LAPORTE, 1832, p. 42. HEER, 1853a. *Mio.*, Europe (Croatia)—*Holo.*
- Syromastes** LATREILLE, 1829, p. 196. HEER, 1853a; STATZ & WAGNER, 1950. *Oligo.*, Europe (Germany); *Mio.*, Europe (Germany)—*Holo.*

Family ALYDIDAE Stål, 1872

[Alydidae STÅL, 1872, p. 53]

Similar to Coreidae, but head nearly as broad and as long as the prothorax, broader than anterior margin of pronotum; body and legs elongate. ŠTYS & RIHA, 1977. *Jur.*—*Holo.*

- Alydus** FABRICIUS, 1803, p. 248. *Holo.*
- Cydamus** STÅL, 1858, p. 33. [Generic assignment of fossil doubtful.] SCUDDER, 1890; ŠTYS & RIHA, 1975b, 1977. *Oligo.*, USA (Colorado)—*Holo.*
- Daclera** SIGNORET, 1862, p. 27. [Generic assignment of fossil doubtful.] THÉOBALD, 1937a; ŠTYS & RIHA, 1977. *Oligo.*, Europe (France)—*Holo.*
- Heeralydus** ŠTYS & RIHA, 1975b, p. 190 [**H. bucculatus*; OD]. Similar to *Alydus* (recent), but head relatively short and having long bucculae reaching proximally between insertion of antennae and anterior margins of eyes. ŠTYS & RIHA, 1977. *Oligo.*, Europe (Germany).
- Monstrococcus** POPOV, 1968, p. 109 [**M. quadrimaculatus*; OD]. Antennae thin and long, nearly as long as body; rostrum also very long; tegmen long, with only one distinct vein; legs long and thin, femora about as long as tibiae and much broader; tarsi with 3 segments. ŠTYS & RIHA, 1977. *Jur.*, USSR (Kazakh).—FIG. 170.4. **M. quadrimaculatus*; lateral view, $\times 3$ (Popov, 1968).
- Orthricorisa** SCUDDER, 1890, p. 429 [**O. longipes*; OD]. Little-known genus, similar to *Leptocoris* (recent). ŠTYS & RIHA, 1977. *Oligo.*, USA (Colorado).
- Protenor** STÅL, 1867, p. 543. [Generic assignment of fossil doubtful.] SCUDDER, 1890; ŠTYS & RIHA, 1977. *Oligo.*, USA (Colorado)—*Holo.*
- Sulcalydus** ŠTYS & RIHA, 1975b, p. 186 [**S. kalabisi*; OD]. Similar to *Alydus* (recent), but membrane of tegmen with more veins and apical part of corium longer. *Oligo.*, Europe (Czechoslovakia).

Family MESOPENTACORIDAE

Popov, 1968

[Mesopentacoridae Popov, 1968, p. 112]

Pronotum transverse, anterior corners projecting; anterior margins of tegmen thickened, forming a ridge along entire length of corium; venation vestigial. Tegmina and pronotum coarsely punctate. [Apparently related to the Urostylidae (recent).] *Jur.*

- Mesopentacoris** POPOV, 1968, p. 112 [**M. costalis*; OD]. Head narrower than pronotum; second antennal segment longest; corium with one vein; tibiae very slender. *Jur.*, USSR (Kazakh).

Family LYGAEIDAE

Schilling, 1829

[Lygaeidae SCHILLING, 1829, p. 85]

Head shorter than thorax and scutellum; antennae straight, not elbowed; four to five veins in membrane of fore wing, not forming ante-apical cells. SLATER, 1964. *Jur.*—*Holo.*

- Lygaeus** FABRICIUS, 1794, p. 133. HEER, 1853a; SCUDDER, 1890; PITON & THÉOBALD, 1935; THÉOBALD, 1937b; SLATER, 1964. *Oligo.*, USA (Colorado), Europe (France, Germany); *Mio.*, Europe (France, Germany, Croatia)—*Holo.*
- Aphanus** LAPORTE, 1832, p. 35. THÉOBALD, 1937a; SLATER, 1964. *Oligo.*, Europe (Germany, France)—*Holo.*
- Catopamera** SCUDDER, 1890, p. 387 [**C. augheyi*; SD SLATER, 1964, p. 1519]. Related to *Myodochina* (recent); head subtriangular, slightly broader than long; antennae slender, no longer than the head and thorax together. SLATER, 1964. *Oligo.*, USA (Colorado).
- Cephalocoris** HEER, 1853a, p. 61 [**C. pilosus*; OD]. Similar to *Cymus* (recent). SLATER, 1964. *Mio.*, Europe (Germany).
- Chilacis** FIEBER, 1864, p. 72. STATZ & WAGNER, 1950. *Oligo.*, Europe (Germany)—*Holo.*
- Cholula** DISTANT, 1882, p. 210. SLATER, 1964. *Eoc.*, USA (Wyoming)—*Holo.*
- Cophocoris** SCUDDER, 1890, p. 391 [**C. tenebricosus*; OD]. Little-known genus; probably close to *Catopamera*, but head rounded; antennae only half as long as body. SLATER, 1964. *Oligo.*, USA (Colorado).
- Coptochromus** SCUDDER, 1890, p. 405 [**C. manium*; OD]. Little-known genus; head fully as long as broad and as broad as apex of thorax. SLATER, 1964. *Oligo.*, USA (Colorado).
- Cryptochromus** SCUDDER, 1890, p. 409 [**C. letatus*; OD]. Related to *Coptochromus*; head large,

- much broader than long. SLATER, 1964. *Oligo.*, USA (Colorado).
- Ctereacoris** SCUDDER, 1890, p. 394 [**C. primigenus*; OD]. Little-known genus; probably related to *Catopamera* but with much shorter middle femora. SLATER, 1964. *Oligo.*, USA (Colorado).
- Diniella** BERGOTH, 1893, p. 202. SLATER, 1964. *Oligo.*, Europe (France, Germany)—*Holo.*
- Drymus** FIEBER, 1860, p. 178. STATZ & WAGNER, 1950. *Oligo.*, Europe (Germany)—*Holo.*
- Eucorites** SCUDDER, 1890, p. 392 [**E. serescens*; OD]. Little-known genus; probably related to *Catopamera* but with more rounded head; antennae longer than head and thorax. SLATER, 1964. *Oligo.*, USA (Colorado).
- Exitelus** SCUDDER, 1890, p. 408 [**E. exsanguis*; OD]. Similar to *Cryptochromus*, but head only a little broader than long. SLATER, 1964. *Oligo.*, USA (Colorado).
- Geocoris** FALLÉN, 1814, p. 10. SCUDDER, 1890. *Oligo.*, USA (Colorado)—*Holo.*
- Heterogaster** SCHILLING, 1829, p. 37. HEER, 1853a; SLATER, 1964. *Oligo.*, Europe (France); *Mio.*, Europe (Germany, Croatia)—*Holo.*
- Ischnodemus** FIEBER, 1837, p. 337. STATZ & WAGNER, 1950. *Oligo.*, Europe (Germany)—*Holo.*
- Ligyrocoris** STÅL, 1872, p. 51. SCUDDER, 1890. *Oligo.*, USA (Colorado)—*Holo.*
- Linnaea** SCUDDER, 1890, p. 396 [**L. carcerata*; SD SLATER, 1964, p. 1523]. Little-known lygaeid; head small; antennae as long as combined head and thorax; thorax very broad. *Oligo.*, USA (Colorado).
- Lithochromus** SCUDDER, 1890, p. 402 [**L. gardneri*; SD SLATER, 1964, p. 1524]. Little-known lygaeid; head as broad as long; antennae about half as long as body. *Oligo.*, USA (Colorado).
- Lithocoris** SCUDDER, 1890, p. 390 [**L. evulsus*; OD]. Little-known genus, apparently related to *Myodochina* (recent); head large, subrotund. SLATER, 1964. *Oligo.*, USA (Colorado).
- Lygaenocoris** POPOV, 1961, p. 1211 [**L. prynadai*; OD]. Eighth abdominal segment strongly developed, covering the ninth. *Jur.*, USSR (Kazakh). —FIG. 170,3. **L. prynadai*; whole insect, $\times 6.5$ (Popov, 1961).
- Lygaeosoma** SPINOLA, 1837, p. 254. SLATER, 1964. *Oligo.*, Europe (Germany)—*Holo.*
- Mesolygaeus** PING, 1928, p. 43 [**M. laiyangensis*; OD]. Similar to *Lygaeus*, but veins of fore wing membrane more prominent. SLATER, 1964. *Cret.*, China (Shantung).
- Miognates** SAILER & CARVALHO in A. R. PALMER, 1957, p. 256 [**M. subimpunctatus*; OD]. Similar to *Lethaeus* and other recent lethaeini but with a smoother integument than is characteristic of the related genera. *Mio.*, USA (California).
- Necrochromus** SCUDDER, 1890, p. 406 [**N. cockerelli*; SD SLATER, 1964, p. 1525]. Body regularly oval; head as broad as apex of thorax. *Oligo.*, USA (Colorado).
- Phrudopamera** SCUDDER, 1890, p. 388 [**P. wilsoni*; SD SLATER, 1964, p. 1521]. Similar to *Catopamera*, but antennae much longer than combined head and thorax. *Oligo.*, USA (Colorado).
- Pionosomus** FIEBER, 1860, p. 48. SLATER, 1964. *Oligo.*, Europe (Germany)—*Holo.*
- Praenotochilus** THÉOBALD, 1937a, p. 289 [**P. parallelus*; OD]. Similar to *Aphanus*, but body more cylindrical and first antennal segment much longer. *Oligo.*, Europe (France).
- Procoris** SCUDDER, 1890, p. 392 [**P. bechleri*; SD SLATER, 1964, p. 1521]. Little-known genus; probably similar to *Eucorites*, but posterior margin of thorax more truncate. *Oligo.*, USA (Colorado).
- Procrophius** SCUDDER, 1890, p. 382 [**P. communis*; SD SLATER, 1964, p. 1512]. Similar to *Crophius* (recent) but with shorter antennae. *Oligo.*, USA (Colorado).
- Procyomophyes** SAILER & CARVALHO in A. R. PALMER, 1957, p. 255 [**P. lithax*; OD]. Similar to *Cymophyes* (recent), but eyes well removed from anterior margin of pronotum. *Mio.*, USA (California).
- Procyms** USINGER, 1940, p. 79 [**P. cockerelli*; OD]. Similar to *Cymus* (recent), but body short, broad, and covered with cymine punctures. SLATER, 1964. *Oligo.*, USA (Colorado).
- Prolygaeus** SCUDDER, 1890, p. 405 [**P. inundatus*; OD]. Body very regularly oval. Antennae as long as head and thorax, the first segment not extending beyond frons, the last two segments longer than first two. *Oligo.*, USA (Colorado).
- Raglius** STÅL, 1872, p. 57. STATZ & WAGNER, 1950. *Oligo.*, Europe (Germany)—*Holo.*
- Rhyparochromus** HAHN, 1826, p. 17. [Most extinct species included here have uncertain generic positions; they were originally put in *Pachymerus* LEPELETIER & SERVILLE, 1825, which is now placed on the Official Index of Rejected and Invalid Names in Zoology (Op. 676, 1963, ICZN).] SLATER, 1964. *Eoc.*, USA (Wyoming); *Oligo.*, Europe (Germany, Baltic, France), USA (Colorado); *Mio.*, USA (Colorado), Europe (Germany, Croatia)—*Holo.*
- Scolopostethus** FIEBER, 1860, p. 188. STATZ & WAGNER, 1950. *Oligo.*, Europe (Germany)—*Holo.*
- Stenopamera** SCUDDER, 1890, p. 385 [**S. senebrosa*; SD SLATER, 1964, p. 1521]. Little-known genus, related to *Catopamera*. *Oligo.*, USA (Colorado).
- Tiromerus** SCUDDER, 1890, p. 401 [**T. torpefactus*; OD]. Little-known genus, similar to *Rhyparochromus*; second segment of antennae much longer than third or fourth. *Oligo.*, USA (Colorado).
- Trapezonotus** FIEBER, 1860, p. 50. SCUDDER, 1890; STATZ & WAGNER, 1950. *Oligo.*, Europe (Germany), USA (Colorado)—*Holo.*

Family BERYTIDAE Fieber, 1851

[Berytidae FIEBER, 1851, p. 9]

Body very slender; head conical, porrect; antennae and rostrum four-segmented; pronotum much longer than wide; scutellum armed; legs very slender; tarsi three-segmented. *Oligo.*—*Holo.*

Berytus FABRICIUS, 1803, p. 264. *Holo.*

Megalomerium FIEBER, 1859, p. 208. THÉOBALD, 1937a. *Oligo.*, Europe (France)—*Holo.*

Family PYRRHOCORIDAE
Fieber, 1860

[Pyrrhocoridae FIEBER, 1860, p. 43]

Body elongate-oval; antennae and beak four-segmented; ocelli absent; membrane with 2 large basal cells, giving rise to several (about 8) branching veins; tarsi three-segmented. *Oligo.*—*Holo.*

Pyrrhocoris FALLÉN, 1814, p. 9. STATZ & WAGNER, 1950. *Oligo.*, Europe (Germany)—*Holo.*

Dysdercus AMYOT & SERVILLE, 1843, p. 272. SCUDDER, 1890. *Oligo.*, USA (Colorado)—*Holo.*

Family CYDNIDAE Billberg, 1820

[Cydniidae BILLBERG, 1820, p. 7]

Similar to the Pentatomidae, but forelegs fossorial; tibiae strongly spinose, veins of fore wing membrane radiating from base. *Eoc.*—*Holo.*

Cydnus FABRICIUS, 1803, p. 184. HEER, 1853a; FÖRSTER, 1891; THÉOBALD, 1937a; STATZ, 1950a. *Oligo.*, Europe (Germany, France)—*Holo.*

Crocistethus FIEBER, 1860, p. 84. STATZ & WAGNER, 1950. *Oligo.*, Europe (Germany)—*Holo.*

Cyrtomenus AMYOT & SERVILLE, 1843, p. 90. SCUDDER, 1890. *Eoc.*, USA (Wyoming)—*Holo.*

Necroclydnus SCUDDER, 1890, p. 443 [**N. amyzonus* SCUDDER, 1890, p. 443; SD CARPENTER, herein]. Head only slightly sunk into prothorax; anterior-lateral angles of thorax rounded. *Eoc.*, USA (Wyoming); *Oligo.*, USA (Colorado).

Procydnus SCUDDER, 1890, p. 438 [**P. quietus* SCUDDER, 1890, p. 438; SD CARPENTER, herein].

Very similar to *Stenopelta*, but body less than twice as long as broad. *Eoc.*, USA (Wyoming); *Oligo.*, USA (Colorado).

Sehirus AMYOT & SERVILLE, 1843, p. 96. STATZ & WAGNER, 1950. *Oligo.*, Europe (Germany)—*Holo.*

Stenopelta SCUDDER, 1890, p. 437 [**Aethus punctulatus* SCUDDER, 1878b, p. 769; OD]. Scutellum triangular, as broad as long; head sunk deeply

into prothorax, the depth of the thoracic emargination being about half its width; body more than twice as long as broad. *Eoc.*, USA (Wyoming).

Teleocydnus HENRIKSEN, 1922b, p. 32 [**T. transitorius*; OD]. Similar to *Cydnus* (recent) but with a long, slender scutellum, reaching about to abdominal apex. *Eoc.*, Europe (Denmark).

Thlibomenus SCUDDER, 1890, p. 448 [**T. petreus*; OD]. Similar to *Necroclydnus*, but head even more prominent; anterior emargination of prothorax slight or absent. *Oligo.*, USA (Colorado).

Family SCUTELLERIDAE Leach, 1815

[Scutelleridae LEACH, 1815, p. 121]

Body oval, usually strongly convex; head triangular; 2 ocelli; rostrum four-segmented; scutellum very large, U-shaped; tarsi three-segmented. *Eoc.*—*Holo.*

Scutellera LAMARCK, 1801, p. 293. *Holo.*

Coptosoma LAPORTE, 1832, p. 73. PITON, 1940a. *Eoc.*, Europe (France)—*Holo.*

Poecilocoris DALLAS, 1848, p. 100. STATZ & WAGNER, 1950. *Oligo.*, Europe (Germany)—*Holo.*

Tectocoris HAHN, 1834, p. 33. HENRIKSEN, 1922b. *Eoc.*, Europe (Denmark)—*Holo.*

Family PACHYMERIDIIDAE
Handlirsch, 1906

[Pachymeridiidae HANDLIRSCH, 1906b, p. 495] [=Sisyrochoridae HANDLIRSCH, 1920, p. 210; Psychochoridae HANDLIRSCH, 1920, p. 298; Hypocimicidae HANDLIRSCH, 1939, p. 119]

Fore wing differentiated into corium and membrane; venation of membrane indistinct; clavus broad and nearly half as long as wing; vein SC separating from R + M near division of R and M. POPOV & WOOTTON, 1977. *Jur.*

Pachymeridium GEINITZ, 1880, p. 529 [**P. dubium*; OD]. Fore wing with SC, R, and M separating at a single point; R branched. POPOV & WOOTTON, 1977. *Jur.*, Europe (Germany). — FIG. 170,2. **P. dubium*; fore wing, X9 (Popov & Wootton, 1977).

Apsicoria HANDLIRSCH, 1939, p. 121 [**A. semideleta*; OD]. Similar to *Sisyrocoris*, but R branched; corium relatively smooth. POPOV & WOOTTON, 1977. *Jur.*, Europe (Germany).

Cathalus HANDLIRSCH, 1939, p. 121 [**C. alutaceus*; OD]. Little-known genus, based on wing fragment; apparently similar to *Sisyrocoris*, but R and M very close together in fore wing; corium less punctate. POPOV & WOOTTON, 1977. *Jur.*, Europe (Germany).

Hypocimex HANDLIRSCH, 1939, p. 119 [**H. membranaceus*; OD]. Little-known genus, based on

poorly preserved specimen; family position doubtful. POPOV & WOOTTON, 1977. *Jur.*, Europe (Germany).

Psychrocoris HANDLIRSCH, 1920, p. 208 [**P. cuneifera*; OD]. Little-known genus, apparently similar to *Sisyrocoris*. Fore wing slender; corium including more than three-fourths of wing surface. POPOV & WOOTTON, 1977. *Jur.*, Europe (Germany).

Sisyrocoris HANDLIRSCH, 1920, p. 210 [**S. rudis*; OD]. Little-known genus. Fore wing coarsely punctate; R unbranched. POPOV & WOOTTON, 1977. *Jur.*, Europe (Germany).

Family PENTATOMIDAE Leach, 1815

[Pentatomidae LEACH, 1815, p. 121]

Body oval; head triangular, porrect, much narrower than thorax; antennae five-segmented; rostrum four-segmented; ocelli present; scutellum extending beyond middle of abdomen, narrowed posteriorly to form triangle; membrane with numerous veins; tarsi two- or three-segmented. *Eoc.*—*Holo.*

Pentatoma OLIVIER, 1789, p. 25. HEER, 1853a; HEYDEN & HEYDEN, 1865; FÖRSTER, 1891; HANDLIRSCH, 1906b. *Eoc.*, Europe (Greenland); *Oligo.*, Europe (Germany, Baltic); *Mio.*, Europe (Croatia, Germany)—*Holo.*

Acanthosoma CURTIS, 1824, p. 28. HEER, 1853a; FÖRSTER, 1891; PITON & THÉOBALD, 1935. *Oligo.*, Europe (Germany); *Mio.*, Europe (Croatia, France)—*Holo.*

Arma HAHN, 1832, p. 91. FÖRSTER, 1891. *Oligo.*, Europe (Germany)—*Holo.*

Asopus BURMEISTER, 1834, p. 19. PITON, 1940b. *Eoc.*, Europe (France)—*Holo.*

Brachypelta AMYOT & SERVILLE, 1843, p. 89. NOVÁK, 1877. *Oligo.*, Europe (Germany, Czechoslovakia)—*Holo.*

Cacoschistus SCUDDER, 1890, p. 459 [**C. maceratus*; OD]. Similar to *Mataeoschistus* (recent) but with broader head and less prominent frontal area. *Oligo.*, USA (Colorado).

Carpocoris KOLENATI, 1846, p. 45. KUKALOVÁ & ŘIHA, 1957. *Mio.*, Europe (Czechoslovakia)—*Holo.*

Deryeuma PITON, 1940a, p. 159 [**D. primordialis*; OD]. Pronotum narrowed in front, notched in region of head; antennae five-segmented, the first segment very short, the second very long; tarsi three-segmented. *Eoc.*, Europe (France).

Dinidorites COCKERELL, 1921e, p. 34 [**D. marginiformis*; OD]. Body narrow; pronotum and scutellum with numerous, dark punctures. *Eoc.*, USA (Colorado).

Doryderes AMYOT & SERVILLE, 1843, p. 121. PITON, 1940a. *Eoc.*, Europe (France)—*Holo.*

Eurydema LAPORTE, 1832, p. 61. HEER, 1853a; PITON & THÉOBALD, 1935; THÉOBALD, 1937a. *Mio.*, Europe (Germany); *Mio./Plio.*, Europe (France)—*Holo.*

Eurygaster LAPORTE, 1832, p. 68. THÉOBALD, 1937a. *Oligo.*, Europe (France)—*Holo.*

Eysarcoris HAHN, 1834, p. 66. HEER, 1853a; NAORA, 1933b; THÉOBALD, 1937a. *Oligo.*, Europe (France); *Mio.*, Europe (Germany); *Tert. (epoch unknown)*, Japan—*Holo.*

Halys FABRICIUS, 1803, p. 180. HEER, 1853a. *Mio.*, Europe (Germany)—*Holo.*

Latahcoris COCKERELL, 1931b, p. 312 [**L. spectatus*; OD]. Head less than one-third width of pronotum; pronotum coarsely punctate, more than twice as long as wide; scutellum with straight sides. *Mio.*, USA (Washington).

Manevalia PITON, 1940a, p. 159 [**M. pachyliformis*; OD]. Little-known genus, apparently related to *Pachylis* (recent). *Oligo.*, Europe (France).

Mesohalys BEIER, 1952, p. 134 [**M. muezzenbergiana*; OD]. Pronotum and mesonotum very coarsely punctate; abdominal tergites finely punctate; front margin of pronotum notched. *Mio.*, Europe (Germany).

Neurocoris HEER, 1853a, p. 23 [**N. rotundatus* HEER, 1853a, p. 23; SD CARPENTER, herein]. Little-known genus; pronotum very broad; tegmen very short and broad. SCUDDER, 1885b. *Mio.*, Europe (Croatia).

Nezara AMYOT & SERVILLE, 1843, p. 143. THÉOBALD, 1937a. *Oligo.*, Europe (France); *Mio./Plio.*, Europe (France).

Pachycoris BURMEISTER, 1835, p. 391. HEER, 1853a. *Mio.*, Europe (Germany)—*Holo.*

Palomena MULSANT & REY, 1866, p. 277. MEUNIER, 1915a. *Mio.*, Europe (France)—*Holo.*

Pentatomites SCUDDER, 1890, p. 461 [**P. foliarum*; OD]. Similar to *Polioschistus* but with sides of thorax convex in front of lateral prominences. VERHOEFF, 1917; COCKERELL, 1927d. *Oligo.*, England, USA (Colorado); *Tert. (epoch unknown)*, USSR (Asian RSFSR).

Phloeocoris BURMEISTER, 1835, p. 371. HEER, 1853a. *Mio.*, Europe (Croatia)—*Holo.*

Poliocoris KIRKALDY, 1910, p. 130 [**P. amnesis*; OD]. Allied to *Teleoschistus*. Body oval; head longer than wide between eyes; scutellum extending halfway to apex of abdomen and rounded posteriorly. *Oligo.*, USA (Colorado).

Polioschistus SCUDDER, 1890, p. 460 [**P. ligatus* SCUDDER, 1890, p. 460; SD CARPENTER, herein]. General form as in *Euschistus* (recent); head in front of eyes subquadrate; thorax very short, about 4 times as broad as long. *Oligo.*, USA (Colorado).

Poteschistus SCUDDER, 1890, p. 458 [**P. obnubilus*; OD]. Little-known genus, with body regularly ovate. *Oligo.*, USA (Colorado).

Pycanum AMYOT & SERVILLE, 1843, p. 171. PITON, 1940a. *Eoc.*, Europe (France)—*Holo.*

Teleocoris KIRKALDY, 1910, p. 129 [**T. pothetias*; OD]. Head prominent, longer than its width between the eyes; pronotum more than 3 times as wide as base of head; scutellum regularly triangular, half the length of abdomen. *Oligo.*, USA (Colorado).

Teleschistus SCUDDER, 1890, p. 454 [**T. antiquus*; SD COCKERELL, 1909b, p. 74]. Head nearly half as broad as thorax and broader than long; apical border of prothorax emarginate; scutellum about as long as wide, reaching less than halfway to end of abdomen. COCKERELL, 1909b; HENRIKSEN, 1922b. *Eoc.*, Europe (Denmark); *Oligo.*, USA (Colorado).

Teryra FABRICIUS, 1803, p. 128. HEER, 1853a. *Mio.*, Europe (Germany)—*Holo.*

Thnetoschistus SCUDDER, 1890, p. 457 [**T. revulsus*; OD] [= *Mataeoschistus* SCUDDER, 1890, p. 459 (type, *M. limigenus* SCUDDER), obj.; the two species are based on counterparts of the same fossil]. Similar to *Euschistus* (recent) but more elongate. *Oligo.*, USA (Colorado).

Tiroschistus SCUDDER, 1890, p. 462 [**T. indurescens*; OD]. Head rounded, with very little extension in front of eyes; antennae 2 times as long as head and thorax together. *Oligo.*, USA (Colorado).

Family ANTHOCORIDAE

Amyot & Serville, 1843

[Anthocoridae AMYOT & SERVILLE, 1843, p. xxxvii]

Similar to the Saldidae, but fore wing membrane without long, closed cells; corium with an embolium. *Oligo.*—*Holo.*

Anthocoris FALLÉN, 1814, p. 9. *Holo.*

Temnostethus FIEBER, 1860, p. 263. STATZ & WAGNER, 1950. *Oligo.*, Europe (Germany)—*Holo.*

Family NABIDAE Costa, 1852

[Nabidae COSTA, 1852, p. 66]

Similar to the Reduviidae but more slender; rostrum with 4 segments; membrane of fore wings with distinctly branched veins or with a few longitudinal veins emitting radiating veins. *Jur.*—*Holo.*

Nabis LATREILLE, 1802, p. 248. HEER, 1853a, 1865; THÉOBALD, 1937a; JORDAN, 1952. *Oligo.*, Europe (Baltic, France); *Mio.*, Europe (Croatia)—*Holo.*
Karanabis BECKER-MIGDISOVA, 1962b, p. 219 [**K. kiritshenkoi*; OD]. Antennae with 4 segments; pronotum conical, strongly narrowed anteriorly; legs long. *Jur.*, USSR (Kazakh).

Family REDUVIIDAE Latreille, 1807

[Reduviidae LATREILLE, 1807, p. 126]

Head shorter than thorax and scutellum, not constricted behind eyes; antennae four-segmented; rostrum three-segmented; fore wings not reticulate; forelegs raptorial. *Oligo.*—*Holo.*

Reduvius FABRICIUS, 1775, p. 729. *Holo.*

Eothes SCUDDER, 1890, p. 355 [**E. elegans*; OD]. Related to *Opsicoetus* (recent), but body more slender and terminal antennal segments stout. *Oligo.*, USA (Colorado).

Evagoras BURMEISTER, 1843, p. 368. HEER, 1853a. *Mio.*, Europe (Germany)—*Holo.*

Harpactor LAPORTE, 1832, p. 8. HEER, 1853a. *Mio.*, Europe (Germany, Croatia)—*Holo.*

Limnaxis GERMAR, 1856, p. 19. HEER, 1853a. *Oligo.*, Europe (Baltic)—*Holo.*

Miocoris COCKERELL, 1927e, p. 591 [**M. fagi*; OD]. Anterior femora stout; first antennal segment not as long as head. *Oligo.*, USA (Colorado).

Pirates BURMEISTER, 1835, p. 222. HEER, 1853a. *Mio.*, Europe (Germany)—*Holo.*

Poliosphageus KIRKALDY, 1910, p. 130 [**P. psychrus*; OD]. Similar to *Repipta* (recent) but with first antennal segment scarcely longer than head; second segment much longer than first. *Oligo.*, USA (Colorado).

Propilocerus WASMANN, 1933, p. 1 [**P. dolosus*; OD]. Similar to *Ptilocerus* (recent) but with second antennal segment and the 2 terminal segments longer and thicker. *Oligo.*, Europe (Baltic).

Prostemma LAPORTE, 1832, p. 12. HEER, 1853a. *Mio.*, Europe (Germany)—*Holo.*

Rhinocoris HAHN, 1833, p. 20. STATZ & WAGNER, 1950. *Oligo.*, Europe (Germany)—*Holo.*

Stenopoda LAPORTE, 1832, p. 26. HEER, 1853a. *Mio.*, Europe (Germany)—*Holo.*

Tagalodes SCUDDER, 1890, p. 356 [**T. inermis*; OD]. Similar to *Taglis* (recent) but with shorter thorax and without spines on fore femora. *Oligo.*, USA (Colorado).

Family TINGIDAE Laporte, 1833

[Tingidae LAPORTE, 1833, p. 47]

Head shorter than thorax; antennae shorter than head, with 4 segments; fore wings lace-like, entirely reticulate. *Oligo.*—*Holo.*

Tingis FABRICIUS, 1803, p. 124. DRAKE & RUHOFF, 1960. *Oligo.*, USA (Colorado); *Mio.*, Europe (Croatia)—*Holo.*

Cantacader AMYOT & SERVILLE, 1843, p. 299. DRAKE, 1950; DRAKE & RUHOFF, 1960. *Oligo.*, Europe (Baltic)—*Holo.*

- Celantia** DISTANT, 1903, p. 137. COCKERELL, 1921f; DRAKE & RUHOFF, 1960. *Oligo.*, Europe (England)—*Holo.*
- Diclyla** STÅL, 1874, p. 57. SCUDDER, 1890; DRAKE & RUHOFF, 1960. *Oligo.*, USA (Colorado), Europe (Czechoslovakia); *Mio.*, Europe (Germany)—*Holo.*
- Eotingis** SCUDDER, 1890, p. 359 [**E. antennata*; OD]. Similar to *Tingis*; pronotum smooth; costal area of fore wing enlarged apically. DRAKE & RUHOFF, 1960. *Oligo.*, USA (Colorado).
- Phatnoma** FIEBER, 1844, p. 57. DRAKE, 1950; DRAKE & RUHOFF, 1960. *Oligo.*, Europe (Baltic)—*Holo.*

Family MIRIDAE Hahn, 1831

[Miridae HAHN, 1831, p. 234]

Head porrect; eyes large; ocelli absent; antennae and beak with 4 segments, beak not held in a groove; scutellum distinct; membrane of tegmen usually with 2 basal cells, veins otherwise absent from membrane; tarsi two-segmented. *Jur.*—*Holo.*

- Miris** FABRICIUS, 1794, p. 183. *Holo.*
- Aporema** SCUDDER, 1890, p. 369 [**A. praestricum*; OD]. Little-known genus, probably close to *Phytocoris*; scutellum large, equiangular, with straight sides. *Oligo.*, USA (Colorado).
- Calocoris** FIEBER, 1858, p. 305. STATZ & WAGNER, 1950. *Oligo.*, Europe (Germany)—*Holo.*
- Capsus** FABRICIUS, 1803, p. 241. [Generic assignment of species doubtful.] SCUDDER, 1890. *Oligo.*, USA (Colorado)—*Holo.*
- Carmelus** DISTANT, 1884, p. 297. [Generic assignment of species doubtful.] SCUDDER, 1890. *Oligo.*, USA (Colorado)—*Holo.*
- Closterocoris** UHLER, 1890, p. 76. [Generic assignment of species doubtful.] SCUDDER, 1890. *Oligo.*, USA (Colorado)—*Holo.*
- Fulvius** STÅL, 1862, p. 322 [= *Oligocoris* JORDAN, 1944a, p. 8 (type, *O. bidentata*)]. CARVALHO, 1954. *Oligo.*, Europe (Baltic)—*Holo.*
- Fuscus** DISTANT, 1884, p. 299. [Generic assignment of species doubtful.] SCUDDER, 1890. *Oligo.*, USA (Colorado)—*Holo.*
- Hadronema** UHLER, 1872, p. 412. SCUDDER, 1890. *Oligo.*, USA (Colorado)—*Holo.*
- Jordanofulvius** CARVALHO, 1954, p. 188, *nom. subt. pro Electrocoris* JORDAN, 1944b, p. 133, *non* USINGER, 1942 [**Electrocoris fuscus* JORDAN, 1944b, p. 133; OD]. Little-known genus, apparently belonging to recent tribe Cylapinae. *Oligo.*, Europe (Baltic).
- Lygus** HAHN, 1831, p. 28. STATZ & WAGNER, 1950. *Oligo.*, Europe (Germany)—*Holo.*
- Miomonalonion** SAILER & CARVALHO in PALMER, 1957, p. 257 [**M. conoidifrons*; OD]. Related to *Monalonion* (recent), but frons conately pro-

duced between antennae; first antennal segment very thick. PALMER, 1957. *Mio.*, USA (California).

- Miridoides** BECKER-MIGDISOVA, 1962b, p. 217 [**M. mesozoicus*; OD]. Antennae shorter than body; tegmen reaching to end of abdomen with front margin convex and only 2 veins in corium. *Jur.*, USSR (Kazakh).—FIG. 170, I. **M. mesozoicus*; ×10 (Becker-Migdisova, 1962b).
- Phytocoris** FALLÉN, 1814, p. 10. GERMAR & BERENDT, 1856; THÉOBALD, 1937a. *Oligo.*, Europe (Baltic, France)—*Holo.*
- Poecilocapsus** REUTER, 1875, p. 73. SCUDDER, 1890. *Oligo.*, USA (Colorado)—*Holo.*
- Scutellifer** POPOV, 1968, p. 108 [**S. karataunicus*; OD]. Antennae longer than body, its first segment longer than pronotum; scutellum very large; membrane of tegmen without spots; fore femora long, slightly flattened; hind legs very long. *Jur.*, USSR (Kazakh).

Family UNCERTAIN

The following genera, apparently belonging to the suborder Heteroptera, are too poorly known to permit assignment to families.

- Cacalydus** SCUDDER, 1890, p. 419 [**C. exstertatus*; SD ŠTYS & ŘIHA, 1977, p. 180]. Little-known genus; probably a coreid. *Oligo.*, USA (Colorado).
- Copidopus** HANDLIRSCH, 1906b, p. 635 [**C. jurasicus*; OD]. Little-known genus. Large species; antennae with 5 segments; hind legs with thickened femora. *Jur.*, Europe (Germany).
- Coreites** HEER, 1853a, p. 56 [**C. crassus* HEER, 1853a, p. 56; SD CARPENTER, herein]. Little-known heteropteron, possibly belonging to Coreidae. PITON & THÉOBALD, 1935. *Oligo.*, Europe (France); *Mio.*, Europe (Croatia).
- Cydnopsis** HEER, 1853a, p. 13 [**C. haidingeri* HEER, 1853a, p. 13; SD CARPENTER, herein]. Little-known genus; legs without spines. COCKERELL, 1909j; HANDSCHIN, 1937; PITON & RUDEL, 1936. *Eoc.*, USA (Colorado); *Oligo.*, Europe (France)—*Mio.*, Europe (Croatia, Germany).
- Deraiocoris** BODE, 1953, p. 128 [**D. insculptus*; OD]. Little-known heteropteron; head and thorax punctate. *Jur.*, Europe (Germany).
- Dichaspis** BODE, 1953, p. 137 [**D. laesa*; OD]. Little-known heteropteron, with small head; wings and venation virtually unknown. *Jur.*, Europe (Germany).
- Electrocoris** USINGER, 1942, p. 43 [**E. brunneus*; OD]. Cimicoid genus, with ocelli present; 4 free, longitudinal veins in membrane of tegmen; abdominal trichobothria absent. *Oligo.*, Europe (Baltic).




FIG. 171. Uncertain (p. 276–277).

Engerrophorus BODE, 1953, p. 144 [**E. nitidus*; OD]. Little-known heteropteron, with small head. *Jur.*, Europe (Germany).

Engynabis BODE, 1953, p. 130 [**E. tenuis*; OD]. Little-known genus, based on incomplete wings and body; probably related to Gerridae. POPOV & WOOTTON, 1977. *Jur.*, Europe (Germany).

Eogerridium BODE, 1953, p. 131 [**E. gracile*; OD]. Little-known genus, based on fragment of body; legs long and slender. *Jur.*, Europe (Germany).

Etirocoris SCUDDER, 1890, p. 425 [**E. infernalis*; OD]. Little-known genus; head elongate, slender, prolonged between antennae. ŠTYS & RIHA, 1977. *Oligo.*, USA (Colorado).

Euraspidium BODE, 1953, p. 137 [**E. granulosum*; OD]. Little-known heteropteron, with punctations on thorax. *Jur.*, Europe (Germany).

Hadrocoris HANDLIRSCH, 1939, p. 117 [**H. oculatellaris*; OD]. Little-known heteropteron, with a large scutellum and punctate head and thorax. [Type of family Hadrocoridae HANDLIRSCH, 1939, p. 116.] *Jur.*, Europe (Germany).

Heteronella EVANS, 1961, p. 22 [**H. marksei*; OD]. Little-known genus. Tegmen with suggestion of costal fracture; venation distinct in basal half of tegmen only. *Trias.*, Australia (Queensland).—FIG. 171.2. **H. marksei*; fore wing, $\times 14$ (Evans, 1961).

Ischnocoris BODE, 1953, p. 136 [**I. bitoratus*; OD]. Little-known genus, with broad head. *Jur.*, Europe (Germany).

Leptoserinetha THÉOBALD, 1937a, p. 362 [**L. navicularis*; OD]. Little-known genus, based on poorly preserved specimen. ŠTYS & RIHA, 1977. *Oligo.*, Europe (France).

Liasocoris WENDT, 1940, p. 19 [**L. hainmulleri*; OD]. Little-known genus, with prominent scutellum. *Jur.*, Europe (Germany).

Megalocoris BODE, 1953, p. 127 [**M. laticlavus*;

OD]. Little-known heteropteron; body large, oval in form; venation unknown. *Jur.*, Europe (Germany).

Ophthalmocoris BODE, 1953, p. 126 [**O. liasscus*; OD]. Little-known insect; fore wings apparently membranous. Ordinal assignment doubtful. *Jur.*, Europe (Germany).

Palaeonepidoideus MEUNIER, 1900, p. 13 [**P. carinata*; OD]. Little-known heteropteron, possibly belonging to the Nepidae. *Jur.*, Europe (Germany).

Pricecoris PINTO & ORNELLAS, 1974b, p. 296 [**P. beckeras*; OD]. Little-known genus, based on poorly preserved specimen; venation not preserved. [Type of family Pricecoridae PINTO & ORNELLAS.] *Cret.*, Brazil (Maranhão).

Probascanion HANDLIRSCH, 1939, p. 118 [**P. megacephalum*; OD]. Little-known heteropteron, with relatively large head; venation unknown. [Type of family Probascanionidae HANDLIRSCH, 1939.] *Jur.*, Europe (Germany).

Pronabis BODE, 1953, p. 129 [**P. utroquelaesus*; OD]. Little-known genus; fore wing without distinct membranous area. *Jur.*, Europe (Germany).

Protocoris HEER, 1852, p. 15 [**P. planus*; OD]. Little-known genus; fore wing with distinct membranous area. [Type of family Protocoridae HANDLIRSCH, 1906b, p. 495.] *Jur.*, Europe (Germany).

Rhepocoris SCUDDER, 1890, p. 426 [**R. praetectus*; SD ŠTYS & RIHA, 1977, p. 182] [= *Parodarmistus* SCUDDER, 1890, p. 421 (type, *P. collisus* SCUDDER; SD ŠTYS & RIHA, 1977)]. Little-known genus, possibly related to family Pyrrhocoridae. *Oligo.*, USA (Colorado).

Stiphroschema BODE, 1953, p. 143 [**S. longealatum*; OD]. Little-known heteropteron, with small head and broad thorax; fore wing apparently very thin. *Jur.*, Europe (Germany).

Strobilocoris BODE, 1953, p. 138 [**S. mediocordatus*; OD]. Little-known heteropteron; thorax quadrate, with coarse sculpturing; venation unknown. *Jur.*, Europe (Germany).

Tenor SCUDDER, 1890, p. 425 [**C. speluncaae*; OD]. Little-known genus, based on poorly preserved specimen. ŠTYS & RIHA, 1977. *Oligo.*, USA (Colorado).

Trachycoris BODE, 1953, p. 142 [**T. abbreviatus*; OD]. Little-known heteropteron; similar to *Strobilocoris* but with broader wings; venation unknown. *Jur.*, Europe (Germany).

Triassocoris TILLYARD, 1922b, p. 466 [**T. myersi*; OD]. Tegmen with corium present in central part of wing; membrane submarginal, separated from corium by impressed line; M and CU arising independently from stem R; radiating veins extending from M to distal margin; clavus sharply defined, short. [Type of family Triassocoridae

TILLYARD.] EVANS, 1956. *Trias.*, Australia (Queensland).—FIG. 171, 1. **T. myersi*; tegmen, $\times 8$ (Tillyard, 1922b).

Suborder UNCERTAIN

The following genera, apparently belonging to the Hemiptera, are too poorly known to permit assignment to suborders.

Family PARAKNIGHTIIDAE

Evans, 1950

[Paraknightiidae EVANS, 1950, p. 250]

Paranotal lobes well developed. Tegmen with costal fracture in basal third of wing. Female with well-developed ovipositor. *Perm.*

Paraknightia EVANS, 1943b, p. 185 [**P. magnifica*; OD]. Tegmen with costal margin thickened; R + M dividing about one-fifth wing length from base. [Originally placed in Homoptera but transferred to Heteroptera by EVANS (1950); moved to a different suborder, Peloridiina, by POPOV (1980b) along with the existing family Peloridiidae.] EVANS, 1950; BECKER-MIGDISOVA & POPOV, 1962. *Perm.*, Australia (New South Wales).—FIG. 172. **P. magnifica*; a, dorsal view, $\times 3$ (Becker-Migdisova & Popov, 1962); b, tegmen, $\times 4.6$ (Evans, 1950).

Family UNCERTAIN

Docimus SCUDDER, 1890, p. 314 [**D. psylloides*; OD]. Little-known genus, based on fragment. HANDLIRSCH, 1907. *Oligo.*, USA (Colorado).

Prosigara SCUDDER, 1890, p. 343 [**P. flabellum*; OD]. Little-known genus, based on poorly preserved specimen. HANDLIRSCH, 1907. *Oligo.*, USA (Colorado).





FIG. 172. Paraknightiidae (p. 277).