



Woody and arboreal habitats of the Green Salamander (*Aneides aeneus*) in the Blue Ridge Mountains

CHRISTOPHER R. WILSON

Conservation Biologist, North American Land Trust / NC Field Office, 196 Arnett Hollow Road, Vilas, NC 28692, 828.963.9291. cwilson@nalt.org

ABSTRACT -- The green salamander (*Aneides aeneus*) is primarily considered a rock crevice dwelling species. However, many early observations from Kentucky, Tennessee, Virginia, and West Virginia report *A. aeneus* taken from woody and arboreal habitats. There have been only four published records of *A. aeneus* using such habitats within the Blue Ridge Disjunct population of southwest North Carolina, northeast Georgia, and northwest South Carolina, and no records since 1952. Here I report two personal observations of *A. aeneus* using arboreal habitats in North Carolina. Additionally, I report nine observations, made by others, of *A. aeneus* using woody, arboreal, or otherwise non-rock-crevice habitats in North and South Carolina, including the first non-rock-crevice *A. aeneus* nesting record for the Blue Ridge. I also speculate that woody and arboreal habitats play a much larger role in the life-history of *A. aeneus* than generally thought, and that the rarity of *A. aeneus* is linked to the loss of American Chestnut and old-growth forests.

INTRODUCTION AND METHODS

The green salamander (*Aneides aeneus*) is distributed from central Alabama to southwestern Pennsylvania along the Appalachian Plateau (Petranka, 1998). A separate and smaller cluster of populations, the "Blue Ridge Disjunct", occurs in the mountains of southwest North Carolina, northeast Georgia, and northwest South Carolina (Bruce, 1968; Petranka, 1998). The green salamander is the only representative of the genus *Aneides*, or "Climbing Salamanders", in the eastern United States and is generally uncommon across its range due to specialized habitat requirements (Petranka, 1998).

Aneides aeneus is almost exclusively observed inhabiting rock crevices of outcrops, which are located within associations of the mixed-mesophytic forest (Gordon, 1952; Corser, 1991; Petranka, 1998). However, early observations from Kentucky, Tennessee, Virginia, and West Virginia report *A. aeneus* breeding and foraging within woody and arboreal habitats (Gordon, 1952). Pope (1928) and Barbour (1949) report *A. aeneus* being taken almost exclusively from beneath the exfoliating bark of standing or fallen, yet "solid", dead trees, mostly American Chestnut (*Castanea dentata*). Fowler (1947) reports *A. aeneus* being found under the bark of a fallen American Chestnut tree. Barbour (1949) reports one *A. aeneus* within the cavity of a standing dead American Chestnut and a brooding female beneath the bark of a fallen tree. Pope (1928) reports one individual within a rotted log and a brooding female within the small cavity of a prostrate limb of an Oak (*Quercus* spp.) tree. Welter and Barbour (1940) and Canterbury (1991) also report observations of *A. aeneus* beneath the loose bark of dead trees or within rotted stumps.

In contrast with these widespread observations of the use of arboreal habitats and woody debris by *A. aeneus* in the main range populations, there are only four published records of *A. aeneus* individuals observed in habitats other than rock crevices in the Blue Ridge Disjunct. Three of these observations were adjacent to, or near, rock outcrops. Bishop (1928) reported an individual *A. aeneus* observed beneath a piece of bark lying on the surface of a rock several feet above the ground in Tallulah Gorge, Rabun Co., GA. Brimley (1941) reported a single *A. aeneus* found within a woodpile on August 1, 1938 in Highlands, Macon Co., NC. Gordon (1952) reported that this woodpile observation was within 100 feet of a rock outcrop colony of *A. aeneus*. Gordon (1952) reports an immature *A. aeneus* individual crawling down a dead oak (*Quercus* spp.) sapling leaning against a rock outcrop along Clear Creek in Rabun Co., GA. Brimley (1927) and Swartz (1954) report an individual *A. aeneus* taken by J.O. Pepper on August 3, 1926 from the “surface of up-standing tree at about 3 ½ ft up, in damp woods” at Pinnacle Mountain, Pickens Co., South Carolina. While the four non-rock-crevice records are of individuals, *A. aeneus* nests are known only from rock crevices in the Blue Ridge Disjunct (Snyder, 1971).

Here I report two personal observations of *Aneides aeneus* using arboreal habitats in North Carolina. Additionally, I report nine observations, made by others, of *A. aeneus* using woody, arboreal, or otherwise non-rock-crevice habitats in North and South Carolina, including the first non-rock-crevice *A. aeneus* nesting record for the Blue Ridge Disjunct. Observational descriptions, provided by others, were collected by contacting professional biologists and amateur naturalists familiar with *A. aeneus* populations in the region. Unless otherwise cited, the following information is summarized from personal communications provided by the observers.

FIELD OBSERVATIONS OF ANEIDES NON-ROCK-CREVICE HABITAT USE IN THE BLUE RIDGE DISJUNCT

The following two observations were made at Biscuit Rock, Highlands, Macon Co, North Carolina, USA, at ca. 1,219 m elevation (35°03'N, 82°11'W). On 6 June 2003, at 1900hrs, during a light rain, with the aid of a high power flashlight, I observed an individual *A. aeneus* in a horizontal position at the bottom of a small, vertical, crevice-like tree cavity (2cm x 37cm), 42cm vertical distance above the ground, and located within a live Rosebay Rhododendron, *Rhododendron maximum*. The diameter of the trunk was 11 cm at the height of the crevice. On 7 June 2003, at 12:20 hrs, during a heavy rain, with the aid of a high power flashlight, I again observed another individual *A. aeneus* in a horizontal position at the bottom of a small tree cavity (2cm x 4cm), 72cm vertical distance above the ground, and located within a separate, live Rosebay Rhododendron, *R. maximum*, approximately 2m from the 6 June 2003 observation. The diameter of the trunk was 8cm at the height of the crevice. Because the trunk of the shrub was slanted, the actual trunk distance the salamander had to travel to reach the crevice from the ground was 97cm. In both observations, the salamanders were small, appeared to be less than one year of age, and retreated further into the cavity upon illumination. Both cavities were apparently created when a horizontal limb detached from the main trunk. Both shrubs were attached to a rock outcrop known to contain a rock-crevice dwelling population of *A. aeneus*.

The following three observations were located in Cedar Mountain, Transylvania Co., North Carolina, USA, at ca. 900m elevation (35°08'N, 82°41'W). The nearest suitable rock outcrop habitat, referred to as “Sherwood Forest Cave”, is >750m from the observations and is known to contain a population of *A. aeneus* (Wilson, 2001). On September 1974, upon splitting a downed American Chestnut log, *Castanea dentata*, Herbert Ball discovered an adult female *A. aeneus* tending a clutch of nine+ eggs (Millie Blaha, pers. com.). The brooding female was photographed by the late George and Millie Blaha (Figure 1) and the following descriptions are approximated, by the author, from their photographs. The log was about 25cm in diameter and contained a hollowed chamber about 6cm in diameter. The clutch contained at least nine eggs suspended from the sides of the chamber. At least four hatchlings were produced within two weeks of the initial discovery (Millie Blaha, pers. com.).

A color photograph was deposited in the N.C. State Museum of Natural Sciences by the author (ALB 10409, Figure 1). On 24 October 1993, Beth and Speed Rogers discovered two *A. aeneus* individuals under a clay terra cotta dish resting upon a rotting Black Oak (*Quercus velutina*) stump. The two salamanders were photographed and documented by Mille Blaha (Figure 2). The stump was approximately 40cm in diameter, 75cm high (measured in the field by the author, May 2000), and was cut seven years prior to the observation (Millie Blaha, pers. com.). The two salamanders were about 7.5cm total length (approximated, by author, from photographs). Upon disturbance, the pair crawled down the side of the stump and into a crevice at the base of the stump. A color photograph was deposited in the N.C. State Museum of Natural Sciences by the author (ALB 10410, Figure 2). During the spring of 2000, Speed Rogers, amateur naturalist, found one large *A. aeneus* climbing on his wooden deck, approximately 4.5m above the ground (Speed Rogers, pers. com.).

The following four observations were made at the Falling Creek Camp, Tuxedo, Henderson Co., North Carolina, USA, at ca 790m elevation (35°11'N, 82°28'W). The nearest suitable rock outcrop habitat, known as "Falling Creek Camp", is >500m from the observations and is known to contain a population of *A. aeneus* (Wilson, 2001). Beginning in June, 1991, and repeatedly throughout the summer, Joe Duckett, resident camp naturalist, observed an individual *A. aeneus* living within the entrance of a small tree cavity (2.5cm × 5cm), approximately 1.5m above the ground (approximated, by author, from photographs), and located within a live standing Scarlet Oak (*Quercus coccinea*). The salamander was feeding on small insects that were attracted to a black sap oozing out of the cavity. The salamander was no longer found after late summer. Identification was confirmed, and the habitat was photographed, by Allen Boynton, non-game wildlife biologist, North Carolina Wildlife Resources Commission, during the summer of 1991. In May, 1997, upon submerging a decorative driftwood log into a fish aquarium, Joe Duckett observed an *A. aeneus* escape from within an approximately 10 cm wide hollow within the log. The driftwood log had previously been left outside on a porch. During May, 1999, Joe Duckett again repeatedly observed an individual *A. aeneus* living in a cavity of a standing, yet dead, tree throughout the summer approximately 10m from the 1991 observation. The cavity entrance was 7.5 cm × 15 cm and located 1.8m above the ground (measured in the field by the author, May 2001). The salamander was no longer found after late summer. During the summer of 1999, Joe Duckett, along with a young camper, observed an individual *A. aeneus* under a log.

Caesar's Head State Park, Greenville Co., South Carolina, USA, at ca 920m elevation (35°06'N, 82°37'W). On several occasions in the fall of 1989, Mae Lee Hafer, park naturalist and Clemson University graduate student studying *A. aeneus*, observed *A. aeneus* individuals climbing on the wooden sides of the Park Naturalist House and on the screened, wooden porch. There are no rock outcrops "relatively close" to the site of observation (Mae Lee Hafer, pers. com.).

Wildcat Wayside, Greenville Co., South Carolina, USA, at ca 340m elevation (35°03'N, 82°30'W). On the night of 25 April 1981, while searching for the Ringneck Snake (*Diadophis punctatus*), Rudy Mancke, natural history curator at the South Carolina State Museum, and Ralph Philips observed a large *A. aeneus* under the exfoliating bark of a fallen oak (*Quercus* spp.) log, (Rudy Mancke, unpublished field notes)

DISCUSSION

The above observations demonstrate unambiguously that populations of *A. aeneus* in the Blue Ridge Disjunct 1) utilize non-rock habitats for breeding and foraging, 2) are not restricted to the immediate vicinity of rock outcrops, and 3) will use non-rock habitats throughout the spring and fall dispersal, and summer breeding phases of its annual life cycle. These observations suggest that the microclimates provided by tree cavities or under exfoliating slabs of bark are similar to those of rock crevices which are used for breeding, foraging, and refuge purposes, and that arboreal habitats historically comprised an important component of the ecological niche of *A. aeneus*.

Woody Habitats

Breeding *A. aeneus* females require cool, clean, and moist, but not wet, horizontal crevices or narrow chambers in which to suspend their eggs from an overhead substrate (Gordon, 1952). Such habitat provides a specific microclimate necessary for successful embryonic development and is typically observed in shaded rock crevices (Gordon, 1952). Woody breeding habitat apparently occurs within narrow cavities or under slabs of exfoliating bark. These microhabitat features occur during the earlier decay classes of a dead tree (Maser et al., 1988), when the tree is still “solid” as Pope (1928) described. Within the range of *A. aeneus*, woody and arboreal micro-habitat features such as tree hollows and exfoliating bark are most abundant within temperate hardwood old-growth forests (Hardt and Swank, 1997; Hale and Rusterholz, 1999).

Arboreal Foraging

Aneides aeneus is exceptionally well adapted to forage and dwell in arboreal habitats due to its anatomical and physiological attributes. Although these features are commonly interpreted as adaptations to a life within rock crevices (Gordon, 1952), they strongly suggest an arboreal life-style as well. In behavioral experiments, *A. aeneus* has an instinctual tendency to climb when confronted with vertical surfaces (Baltar, 1983). Adaptations for climbing include long-legs, expanded toe-pads, a flattened body, and a muscular tail (Gordon, 1952; Wake, 1963). *Aneides aeneus* is physiologically more tolerant to desiccation than any other salamander in its range, which allows it to withstand the drier conditions encountered when climbing on exposed surfaces such as rock cliffs (Gordon, 1952; Canterbury, 1991) or in trees. The unique lichen colored and patterned dorsum of *A. aeneus* provides excellent camouflage among the lichens and mosses that are commonly distributed over the surfaces of both rock outcrops, and tree trunks and limbs, within the range of *A. aeneus* (Karin Heiman, pers. com.). The very presence of such a bright and striking color pattern suggests an adaptation to predation from visual predators such as birds (Brandon and Huheey, 1975), which forage more frequently in trees than on rock. In behavioral experiments, *A. aeneus* is actually known to prefer bark substrates to rock (Mushinsky, 1976). Finally, other species of *Aneides* are highly arboreal in habit (Petranka, 1998).

Food is known to be a limiting factor in competition between salamanders (Jaeger, 1972). *Plethodon cinereus*, an otherwise ground dwelling salamander, was shown to realize higher foraging efficiency by climbing plant stems, where it acquires larger and more abundant invertebrate prey items (Jaeger, 1978). Due to its lack of tolerance to desiccation, *P. cinereus* is only briefly able to forage on plant stems during periods of high humidity on rainy nights (Jaeger, 1978). A number of other plethodontids in the eastern U.S. have been observed climbing on plant stems on rainy nights, presumably to forage (Hairston, 1949; pers. obs.; R. Wayne Van Devender, pers. com.). *Aneides aeneus* is known to use its climbing abilities and tolerance to desiccation to avoid competition with other plethodontids by climbing to higher and drier positions on rock outcrops (Baltar, 1983; Cliburn and Porter, 1987). Given the abundance and diversity of potential plethodontid competitor species on the forest floor, the relatively unexploited food resource in arboreal habitats, and the unique adaptations of *A. aeneus* to take advantage of arboreal habitats, it seems reasonable that arboreal foraging plays a larger role in the life history of *A. aeneus* than generally reported. Old-growth temperate hardwood forests contain a more complex vertical structure with higher densities of invertebrates (Schowalter, 1995), and cooler and moister microclimates relative to logged forests (Burry and Corn, 1988; deMaynadier and Hunter, 1995).

Such habitats, which historically dominated the range of the Blue Ridge Disjunct populations of *A. aeneus* (Braun, 1950), likely provide it increased arboreal foraging opportunities and decreased likelihood of desiccation.

Life History Implications

Breeding and foraging in woody and arboreal habitats may explain some apparent inconsistencies in the known life history traits of *A. aeneus*. For example, the highest seasonal densities of *A. aeneus* are found in rockcrevices during the spring “posthibernation aggregation and dispersal” and fall “prehibernation dispersal and aggregation” periods, with many fewer individuals observed during the summer “breeding” period (Gordon, 1952; Cupp, 1991; Hafer, 1992). Cupp (1991) suggested that this decrease in density is explained by emigration to outlying outcrops as far as 100m distant, while other herpetologists suggest the missing salamanders simply retreat further into the crevices, out of sight. *Aneides aeneus* certainly migrates between rock outcrops (Gordon, 1952; Canterbury, 1991), however many Blue Ridge colonies have no other apparently suitable rock outcrops in their vicinity, perhaps as far as 1km (pers. obs.). In addition, retreating deeper into crevices during the non-hibernating phases would seem to reduce foraging efficiency and cause greater intraspecific competition, particularly at colonial sites. Perhaps, instead, the lower densities of visible individuals at rock crevices in the summer are explained by extended foraging forays into arboreal habitats or the utilization of woody breeding habitats.

Another inconsistency that is potentially explained by arboreal foraging forays is the lower frequency of *A. aeneus* observations at rock crevices during periods of heavy or extended rain. *Aneides aeneus* is known to become more active during humid periods and is usually only observed outside of rock crevices, on the rock surface, during misting conditions, light rain, or moderate rain (Gordon, 1952; Cupp, 1991; Hafer, 1992). Yet, mysteriously, non-breeding individuals are observed in rock crevices far less often during periods of heavy rain or after long periods of rain, particularly in the summer (Hafer, 1992; pers. obs.). Hafer (1992) concluded that abundant rain makes conditions “too wet” for *A. aeneus* to come far enough out of the crevices to be observed. However, *A. aeneus* has been observed crossing roads on rainy nights (Williams and Gordon, 1961). While light or brief rains may create improved conditions for short forays near the rock crevice during the summer, perhaps heavy or extended periods of rain create ideal conditions for arboreal foraging explaining the brief decrease of observations at rock sites. Many plethodontid salamanders exhibit increased activity and foraging outside of their typical microhabitats during rainy periods (Bernardo, pers. com.; Hairston, 1949). In contrast, non-brooding *A. aeneus* likely seek transitory refuge in the humid conditions offered in rock or woody crevices during dry periods.

Rarity and Population Decline

The scarcity of woody and arboreal *A. aeneus* observations relative to rock observations is potentially due to a low population density across the landscape, the general inaccessibility of arboreal situations and micro-habitats such as tree cavities, and a general lack of suitable woody habitats (Petranka, 1998). Perhaps the rarity of *A. aeneus* is linked to the decline of mature American Chestnut trees and the loss of old-growth forests (Braun, 1950).

Many arboreal and woody habitat observations of *A. aeneus* are associated with American Chestnut, a tree which disappeared from the forest canopy by 1950 after the introduction of *Cryphonectria parasitica*, an Asiatic blight, around 1904 (Ronderos, 2000). The previous abundance of *C. dentata* and its rot resistant properties may have maintained the early decay classes necessary for *A. aeneus* breeding over a longer time frame than other types of woody debris, making habitat less spatially and temporally patchy.

The Clouded Salamander (*Aneides ferreus*), a close relative of *A. aeneus*, is found in western coastal forests of the United States and British Columbia and shares similar microhabitat preferences (Petranka, 1998). *Aneides ferreus* is found under loose bark, in rockface or talus crevices, in lightly decayed logs and stumps, and is arboreal (Petranka, 1998). Although this species is associated with old growth forests, it will thrive following clear-cut logging due to the subsequent abundance of woody debris in the early decay stages (Bury and Corn, 1988). However, *A. ferreus* populations decline as small diameter woody debris habitats quickly decay and disappear (Bury and Corn, 1988).

Similar observations of sudden and temporary local increases in *A. aeneus* populations were made by Barbour (1949, 1971) following clear-cuts in Kentucky. Over the long-term, *A. ferreus* may not persist in reforested areas that retain inadequate levels of coarse woody debris (Butts and McComb, 2000) and may take several hundred years to recover from logging (Corn and Bury, 1991).

A similar scenario may explain the rarity of *A. aeneus*. Old-growth forests likely provided abundant woody microhabitats and arboreal foraging opportunities for *A. aeneus*. Populations temporarily increased following a flush of woody habitats, in early decay stages, created by debris leftover from widespread logging and the mortality of American Chestnut during the first half of the last century. As woody habitats associated with old growth and American Chestnut decayed and disappeared across the landscape, *A. aeneus* populations decreased and became increasingly dependent on rock crevice habitats. The second- and third- growth forests that currently dominate the range of *A. aeneus* typically contain far less woody debris than the primary forest (Hardt and Swank, 1997; Hale and Rusterholz, 1999).

The loss of appropriate woody habitats and optimal arboreal foraging conditions due to the chestnut blight and logging may have restricted *A. aeneus* primarily to the crevices of rock outcrops (Gordon, 1952; Corser, 1991; Petrankska 1998; Corser, 2001). Populations within the Blue Ridge Disjunct comprise two genetically distinctive, and distantly related groups that harbor unique genetic diversity compared with populations throughout the main part of the range (Bernardo et al., manuscript in review). Moreover, Blue Ridge Disjunct populations have experienced severe declines (Corser, 2001) and are at a high risk of extinction according to the IUCN Redbook Criteria (Wilson, 2001), while populations within the Appalachian Plateau appear to be relatively stable (Snyder, 1983; Snyder, 1991). Rock outcrops containing appropriate rock-crevice habitat are small, few, patchily distributed, and isolated throughout the Blue Ridge Disjunct relative to the Appalachian Plateau (pers. obs.; Corser, 2001), perhaps making the Blue Ridge populations more susceptible to stochastic processes and contributing to the reported decline. If *A. aeneus* did historically utilize a broader spectrum of forest habitats, as the evidence reviewed herein suggests, then the habitat shift to using principally rock outcrops has an anthropogenic origin and the decline of the rock-crevice dwelling populations in the Blue Ridge Disjunct (Corser, 2001) is also attributable to human disturbance, via logging as well as by introduction of the exotic chestnut blight. Thus, conservation of Blue Ridge Disjunct populations is of significant concern.

The observations reported herein have important implications for conserving the remaining populations. While rock crevices currently represent the most critical habitat for *A. aeneus*, future conservation efforts should recognize the importance of woody and arboreal habitats in surrounding woodlands. Management should allow forests to mature and return to a natural "gap dynamic" (Runkle, 1985) method of regeneration and woody debris production over the next century. Such forests will provide abundant breeding and foraging habitat, as well as provide optimal corridors between existing rock outcrop colonies, facilitating metapopulation persistence (Hanski and Simberloff, 1997). In addition, supplementation of coarse woody debris in areas occupied by *A. aeneus* might aid recovery by enhancing the availability of both foraging and breeding sites, just as nest boxes are used to augment reproduction in bird populations.

ACKNOWLEDGEMENTS

I thank Millie Blaha, Speed Rogers, Allen Boynton, Joseph Duckett, Mae Lee Hafer, and Rudy Mancke for providing descriptions of their *A. aeneus* observations. William D. Hendricks and Gavin Wilson reviewed Figure 1 and commented on the identification and decay properties of the American Chestnut log. I thank R. Wayne Van Devender and Karin Heiman for their comments and assistance. I also thank Marla Wilson and the staff and board of North American Land Trust for their support and encouragement. Joseph Bernardo, Jim Petranka, and Jeff Corser provided suggestions that improved the manuscript.

LITERATURE CITED

BALTAR, A. C.

1983. Vertical stratification of some cliff dwelling plethodontid salamanders as related to competitive survival of the Mississippi population of *Aneides aeneus*. M. S. Thesis, University of Southern Mississippi. Mississippi.

BARBOUR, R. W.

1949. A study of the mammals, reptiles and amphibians of Big Black Mountain, Harlan County, Kentucky. PhD thesis. Cornell University, NY.

1971. Amphibians and Reptiles of Kentucky. University Press of Kentucky, Lexington, Kentucky.

BISHOP, S. C.

1928. Notes on some amphibians and reptiles from the southeastern states with description of a new salamander from North Carolina. *Journal of the Elisha Mitchell Scientific Society* 43:153-170.

BRANDON, R. A. AND J. E. HUHEEY.

1975. Diurnal activity, avian predation, and the question of warning coloration and cryptic coloration in salamanders. *Herpetologica* 31:252-255.

BRAUN, E. L.

1950. Deciduous forests of eastern North America. Blakiston, Philadelphia, Pennsylvania.

BRIMLEY, C. S.

1927. Some records of amphibians and reptiles from North Carolina. *Copeia* 162:10-12.

1941. The amphibians and reptiles of North Carolina. Installment No.15, *Carolina Tips* (Elon College, N. C.) 4:23.

BRUCE, R. C.

1968. The role of the Blue Ridge Embayment in the zoogeography of the green salamander, *Aneides aeneus*. *Herpetologica* 24:185-194.

BURY, R. B. AND P. S. CORN.

1988. Douglas-fir forests in the Oregon and Washington Cascades: Relation of the herpetofauna to stand age and moisture. Pages 11-22. in *Management of amphibians, reptiles and small mammals in North America* (R. C. Szaro, K. E. Severson and D. R. Patton, eds). USDA Forest Service General Technical Report RM-166.

BUTTS, S. R. AND W. C. MCCOMB

2000. Associations of forest-floor vertebrates with coarse woody debris in managed forests of western Oregon. *Journal of Wildlife Management* 64:95-104.

CANTERBURY, R. A.

1991. Ecology of the green salamander, *Aneides aeneus* (Cope and Packard), in West Virginia. M. S. Thesis, Marshall University, West Virginia.

CLIBURN, J. W., AND A. B. PORTER

1987. Vertical stratification of the salamanders *Aneides aeneus* and *Plethodon glutinosus* (Caudata, Plethodontidae). *Journal of the Alabama Academy of Science* 58:18-22.

CORN, P. S. AND R. B. BURY

1991. Terrestrial amphibian communities in the Oregon Coast Range. Pages 304-317 in *Wildlife and vegetation of un-managed Douglas-fir forests* (L. E. Ruggiero, K. B. Aubry, A. B. Carey, and M. H. Huff, tech. cords.). USDA Forest Service, Pacific Northwest Research Station, Olympia, Washington, General Technical Report PNW-GTR-285,

CORSER, J. D.

1991. The ecology and status of the endangered green salamander, *Aneides aeneus*, in the Blue Ridge Embayment of North Carolina. M.S. Thesis, Duke University, North Carolina.

2001. Decline of disjunct green salamander (*Aneides aeneus*) populations in the southern Appalachians. *Biological Conservation* 97:119-126.

CUPP, P. V., JR.

1991. Aspects of the life history and ecology of the green salamander, *Aneides aeneus*, in Kentucky. *Journal of the Tennessee Academy of Science* 66:171-174.

DEMAYNDAIER, P. G. AND M. I. HUNTER, JR

1995. The relationship between forest management and amphibian ecology: A review of North American literature. *Environmental Review* 3: 230-261.

FOWLER, J. A.

1947. Record for *Aneides aeneus* in Virginia. *Copeia* 1947:144.

GORDON R. E.

1952. A contribution to the ecology and life history of the plethodontid Salamander, *Aneides aeneus*. *American Midland Naturalist* 47:666-701.

HAFER, M. A.

1992. A survey of the green salamander in South Carolina. M. S. Thesis. Clemson University, South Carolina.

HAIRSTON, N. G. 1949. The local distribution and ecology of the plethodontid salamanders of the southern Appalachians. *Ecological Monographs* 19: 47-73.

HALE, C. M. AND K. A. RUSTERHOLZ.

1999. Comparison of structural and compositional characteristics in old-growth and mature, managed hardwood forests of Minnesota, U.S.A. *Canadian Journal of Forest Research* 29:1479-1489.

HANSKI, I AND D. SIMBERLOFF.

1997. The metapopulation approach, its history, conceptual domain, and application to conservation. Pages 5-26 in *Metapopulation Biology* (I. A. Hanski and M. E. Gilpin, eds.). Academic Press, San Diego, California.

HARDT, R. A. AND W. T. SWANK

1997. A comparison of structural and compositional characteristics of Southern Appalachian young second-growth, maturing second-growth, and old-growth stands. *Natural Areas Journal* 17:42-52.

JAEGER, R. G.

1972. Food as a limited resource in competition between two species of terrestrial salamanders. *Ecology* 53:535-546.

1978. Plant climbing by salamanders: periodic availability of plant dwelling prey. *Copeia* 1978:686-691.

LARSON, A., D. B. WAKE, L. R. MAXSON and R. HIGHTON.

1981. A molecular phylogenetic perspective on the origins of morphological novelties in the salamanders of the tribe Plethodontini (Amphibia, Plethodontidae). *Evolution* 35:405-422.

MASER, C., R. F. TARRAN, J. M. TRAPPE AND J. F. FRANKLIN (eds).

1988. *From the Forest to the Sea: a Story of Fallen Trees*. USDA Forest Service General Technical Report PNW-GTR-229, Pacific Northwest Research Station, Portland, OR. 153pp.

MUSHINSKI, H. R.

1976. Ontogenetic development of microhabitat preference in salamanders: The influence of early experience. *Copeia* 1976:755-758.

PETRANKA, J. W.

1998. *Salamanders of the United States and Canada*. Smithsonian Institution Press, Washington, D.C. 587pp.

POPE, C. H.

1928. Some plethodontid salamanders from North Carolina and Kentucky, with description of a new race of *Leurognathus*. *American Museum Novitates* 306:1-19.

RONDEROS, A

2000. Where giants once stood: the demise of the American Chestnut and efforts to bring it back. *Journal of Forestry* 98:10-11.

RUNKLE, J. R.

1985. Disturbance regimes in temperate forests. Pages 371-384 in *The Ecology of Natural Disturbance and Patch Dynamics* (S. T. A. Pickett and P. S. White, eds.). Academic Press, New York.

SCHOWALTER, T. D.

1995. Canopy invertebrate community response to disturbance and consequences of herbivory in temperate and tropical forests. *Selyana* 16:41-48.

SNYDER, D. H.

1971. The function of brooding behavior in the plethodontid salamander, *Aneides aeneus*: A field study. Ph. D. Dissertation, University of Notre Dame, Illinois.

1983. The apparent crash and possible extinction of the green salamander, *Aneides aeneus*, in the Carolinas. *Association of Southeastern Biologists Bulletin* 30:82.

1991. The green salamander *Aneides aeneus* in Tennessee and Kentucky, with comments on the Carolina's Blue Ridge populations. *Journal of the Tennessee Academy of Sciences* 66:165-169.

SCHWARTZ, A.

1954. The salamander *Aneides aeneus* in South Carolina. *Copeia* 154:296-298.

WAKE, D. B.

1963. Comparative osteology of the plethodontid salamander genus *Aneides*. *Journal of Morphology* 113:77-118.

WILLIAMS, K. L. AND R. E. GORDON.

1961. Natural dispersal of the Green Salamander *Aneides aeneus*. *Copeia* 1961:353.

WILSON, C. R.

2001. Status of the Green Salamander (*Aneides aeneus*) in the Blue Ridge Mountains. M.S. Thesis. Appalachian State University, North Carolina.

WELTER, W. A. AND R. W. BARBOUR.

1940. Additions to the herpetofauna of northeastern Kentucky. *Copeia* 1940:132.

© 1997 - 2003 All Rights Reserved, Contemporary Herpetology.