

VOLUME 12, NUMBER 2 JUNE 2005

IGUANA

CONSERVATION, NATURAL HISTORY, AND HUSBANDRY OF REPTILES

International Reptile Conservation Foundation

www.IRCF.org



GLENN MITCHELL

Anegada or Stout Iguana (*Cyclura pinguis*) (see article on p. 78).




JOHN BINNIS

A subadult Stout Iguana (*Cyclura pinguis*) awaiting release at the head-starting facility on Anegada (see article on p. 78).



RICHARD SANDAK

Timber Rattlesnakes (*Crotalus horridus*) reach the northernmost extent of their range on the bluff prairies in Wisconsin (see article on p. 90).



This cross-stitch was created by Anne Fraser of Calgary, Alberta from a photograph of Carley, a Blue Iguana (*Cyclura lewisi*) at the Captive Breeding Facility on Grand Cayman. The piece, which measures 6.8 x 4.2 inches, uses 6,100 stitches in 52 colors and took 180 hours to complete. It will be auctioned by the IRCF at the Daytona International Reptile Breeder's Expo on 20 August 2005. Profits from the auction will benefit conservation projects for West Indian Rock Iguanas in the genus *Cyclura*.



ROBERT POWELL

Red-bellied or Black Racers (*Alsophis rufiventris*) remain abundant on mongoose-free Saba and St. Eustatius, but have been extirpated on St. Christopher and Nevis (see article on p. 62).



ROBERT POWELL

Spiny-tailed Iguanas (*Ctenosaura similis*) are abundant in archaeological zones on the Yucatán Peninsula. This individual is on the wall around the Mayan city of Tulum (see article on p. 112).