

VOLUME 13, NUMBER 1


MARCH 2006

# IGUANA

CONSERVATION, NATURAL HISTORY, AND HUSBANDRY OF REPTILES

International Reptile Conservation Foundation

[www.IRCF.org](http://www.IRCF.org)



L. LEE GRISMER

Red-headed Kraits (*Bungarus flaviceps*) occur in lowland rainforests of southern Malaysia, one of the most biologically diverse and complex terrestrial ecosystems in the world (see related article on p. 2).



KENNETH L. KRYSKO

Brown Basilisks (*Basiliscus vittatus*) are well-established in southern Florida and are expanding their range (see article on p. 24).



ALEJANDRO SANCHEZ

The Puerto Rican Dwarf Gecko, *Sphaerodactylus klauberi*, is found only in mesic, montane habitats (see article on p. 16).



JOSEPH BURGESS

Cuban Iguanas (*Cyclura nubila*) are but one of several Cuban reptiles that use the U.S. Naval Base at Guantanamo Bay as a wildlife refuge (see article on p. 8).



ROBERT POWELL

Male anoles are fiercely territorial. Here two male *Anolis cristatellus* on Guana, British Virgin Island, battle over a territory centered on a small palm tree (see article on p. 22).



JOSEPH M. POLANCO

Adult male Emerald Tree Boa (*Corallus caninus*) with head elevated in response to stimuli indicating the presence of a possible meal. Long thought to be a difficult captive, the key to successful husbandry is careful preparation (see article on p. 37).



L. LEE GRISHIER

The Cat Gecko (*Aeluroscalabotes felinus*) is a slow moving, almost catatonic species with a range that extends from southern Thailand south through peninsular Malaysia and Singapore to Borneo (see article on p. 2).



JOSEPH BURGESS

“El Lagarto Azul,” the Blue Anole (*Anolis gorgonae*) of Isla Gorgona, Colombia (see travelogue on p. 47).