VOLUME 13, NUMBER 2 JUNE 2006

GUAINA

CONSERVATION, NATURAL HISTORY, AND HUSBANDRY OF REPTILES

International Reptile Conservation Foundation

www.IRCF.org





Gliding Lizards in the Genus *Draco* use wings for "flight" and also to facilitate species recognition and intra-species communication (see article on p. 86).



Exploitation of the Pacific Spiny-tailed (Black) Iguana, *Ctenosaura pectinata*, was a major topic of discussion during the VIII National Meeting on Iguanas in México (see article on p. 130).



Ctenosaura bakeri lives an unusual life in the mangrove forests on Isla de Utila, Honduras; both the mangrove forests and nesting beaches are threatened by development (see article on p. 108).



The conservation status of the Exuma Island Rock Iguana, *Cyclura cychlura figginsi*, was one of many discussed during the IUCN Iguana Specialist Group Meeting in November 2005 (see article on p. 116).



Chironius monticola is a terrestrial snake most often seen basking in warm, open areas of the Santa Lucia cloud forest reserve (see article on p. 94).



Nose-rub, clearly evident in this Emerald Treeboa (*Corallus caninus*), is indicative of being kept in an inappropriate enclosure. This may reflect neglect in other aspects of care (see article on p. 133).



IMPORTANT NOTICE: Please support the August 2006 National Reptile Breeder's Expo Auction to benefit the critically endangered Guatemalan Beaded Lizard, *Heloderma horridum charlesbogerti* (see p. 176). For details visit www.IRCF.org.