

QE
113
.A2
#52
p4

**TABULAR DESCRIPTION OF OUTCROPPING
ROCKS IN KANSAS**

By

**RAYMOND C. MOORE, JOHN C. FRYE, and
JOHN MARK JEWETT**

**UNIVERSITY OF KANSAS PUBLICATIONS
STATE GEOLOGICAL SURVEY OF KANSAS, BULLETIN 52, PART 4
LAWRENCE, KANSAS**

Generated at University of Kansas on 2023-09-20 18:29 GMT / <https://hdl.handle.net/2027/mdp.39015006833084>
Public Domain in the United States; Google-digitized / http://www.hathitrust.org/access_use#pd-us-google

STATE OF KANSAS

ANDREW F. SCHOEPPLE, *Governor*

STATE BOARD OF REGENTS

LESTER MCCOY, *Chairman*

JERRY E. DRISCOLL

FRED M. HARRIS

MRS. ELIZABETH HAUGHEY

WILLIS N. KELLY

DREW McLAUGHLIN

GROVER POOLE

LA VERNE B. SPAKE

OSCAR STAUFFER

MINERAL INDUSTRIES COUNCIL

JOHN ALLISON, *Chairman*

M. L. BREIDENTHAL

HOWARD CAREY

ANTHONY FOLGER

LESTER MCCOY

J. E. MISSIMER

BRIAN O'BRIAN, *Vice-Chairman*

J. A. SCHOWALTER

CHESTER SCOTT

K. A. SPENCER

W. L. STRYKER

B. O. WEAVER

STATE GEOLOGICAL SURVEY OF KANSAS

DEANE W. MALOTT, M.B.A., LL.D., Chancellor of the University of Kansas, and
ex officio Director of the Survey.

RAYMOND C. MOORE, Ph.D., Sc.D., Director and State Geologist†

JOHN C. FRYE, Ph.D., Asst. Director and Asst. State Geologist, in charge

EDITH HICKS LEWIS, Secretary

STRATIGRAPHY, PALEONTOLOGY, AND AREAL GEOLOGY:

Raymond C. Moore, Ph.D., Sc.D., Geol-
ogist†

John M. Jewett, Ph.D., Geologist

Walter H. Schoewe, Ph.D., Geologist

Arthur L. Bowshe, B.S., Geologist†

SUBSURFACE GEOLOGY:

M. L. Thompson, Ph.D., Geologist

Ada Swineford, M.S., Geologist

Philip Kaiser, A.B., Geologist

Eileen Martin, B.M., Stenographer

Ethelyn B. McDonald, M.A., Well

Sample Curator, Wichita Branch

Arden D. Brown, Assistant

Carrie Thurber, Assistant

MINERAL RESOURCES:

George E. Abernathy, E.M., Ph.D.,
Geologist

Robert M. Dreyer, Ph.D., Geologist†

Walter A. Ver Wiebe, Ph.D., Geologist

Norman Plummer, A.B., Ceramist

John I. Moore, M.S., Petroleum Engr.†

Frances L. Schloesser, Assistant Chemist

W. P. Ames, A.B., Laboratory Assistant

Ethel Owen, Assistant

PUBLICATIONS AND RECORDS:

Donald E. Dowers, Draftsman†

Robyn Ashby, Draftsman

Betty J. Hagerman, Bookkeeper

COOPERATIVE PROJECTS WITH UNITED STATES GEOLOGICAL SURVEY

GROUND-WATER RESOURCES:

Stanley W. Lohman, M.S., Geologist in
charge

John C. Frye, Ph.D., Geologist

Thad G. McLaughlin, Ph.D., Geologist

Bruce Latta, A.B., Geologist

Charles C. Williams, M.A., Geologist

V. C. Fishel, B.S., Physicist

James B. Cooper, A.B., Well Driller†

Oscar S. Fent, B.S., Well Driller

Charles K. Bayne, A.B., Instrumentman

Fern H. Ashby, Stenographer

Betty Ann Ball, Stenographer

W. W. Wilson, Well Observer†

MINERAL FUELS RESOURCES:

Wallace Lee, E.M., Geologist in charge

Constance Leatherock, B.S., Geologist

TOPOGRAPHIC SURVEYS:

C. L. Sadler, Division Engineer

Max J. Gleissner, Section Chief

J. P. Rydeen, Topographer

SPECIAL CONSULTANTS: Ray Q. Brewster, Ph.D., Chemistry; Claude W. Hibbard, Ph.D.,
Vertebrate Paleontology; Eugene A. Stephenson, Ph.D., Petroleum Engineering.

COOPERATING STATE AGENCIES: *State Board of Agriculture, Division of Water Resources*,
George S. Knapp, Chief Engineer; Robert Smrha, Assistant Chief Engineer; *State*
Board of Health, Division of Sanitation, Paul D. Haney, Chief Engineer and Di-
rector; Ogden S. Jones, Geologist.

† Absent on leave for military service.

STATE GEOLOGICAL SURVEY OF KANSAS, BULLETIN 52
1944 REPORTS OF STUDIES, PART 4, PAGES 137-212, FIGS. 1-9
OCTOBER 25, 1944

TABULAR DESCRIPTION OF OUTCROPPING ROCKS IN KANSAS

By
RAYMOND C. MOORE, JOHN C. FRYE, AND
JOHN MARK JEWETT

UNIVERSITY OF KANSAS PUBLICATIONS
STATE GEOLOGICAL SURVEY OF KANSAS, BULLETIN 52, PART 4
LAWRENCE, KANSAS

CONTENTS

	PAGE
ABSTRACT	141
INTRODUCTION	141
Main rock divisions and their distribution	142
Distribution and structure of Cenozoic rocks	143
Distribution and structure of Mesozoic rocks	143
Distribution and structure of Paleozoic rocks	143
Igneous rocks	144
CENOZOIC ROCKS	145
Quaternary system	145
Recent series	145
Pleistocene series	146
Tertiary system	148
Pliocene series	148
MESOZOIC ROCKS	151
Cretaceous system	151
Gulfian series	151
Comanchean series	154
(?) Triassic system	154
PALEOZOIC ROCKS	156
Permian system	156
Guadalupian series	156
Leonardian series	157
Wolfcampian series	160
Pennsylvanian system	170
Virgilian series	170
Missourian series	184
Desmoinesian series	194
Mississippian system	200
Meramecian series	200
Osagian series	200
IGNEOUS ROCKS	200
BIBLIOGRAPHY OF OUTCROPPING ROCKS	202
INDEX OF STRATIGRAPHIC NAMES	209

ILLUSTRATIONS

FIGURE	PAGE
1. Map showing distribution of main divisions of outcropping rocks of Kansas	140
2. Generalized sections of Cenozoic rocks in Kansas	150
3. Generalized section of Cretaceous rocks in Kansas	155
4. Generalized section of upper Permian rocks in Kansas	161
5. Generalized section of middle Permian rocks in Kansas	162
6. Generalized section of lower Permian rocks in Kansas	171
7. Generalized section of upper Pennsylvanian rocks in Kansas	185
8. Generalized section of middle Pennsylvanian rocks in Kansas	198
9. Generalized section of lower Pennsylvanian rocks in Kansas	201




FIG. 1. Map showing distribution of main divisions of outcropping rocks of Kansas.

TABULAR DESCRIPTION OF OUTCROPPING ROCKS IN KANSAS

By RAYMOND C. MOORE, JOHN C. FRYE, and JOHN MARK JEWETT

ABSTRACT

This paper summarizes knowledge of the outcropping rocks of Kansas, indicating in tabular form their present classification and nomenclature. Range in thickness and average thickness of each described unit, based on observation of outcrops but not on subsurface data, are indicated. It is the purpose of the contribution to make available a compact outline of the stratigraphy of the Kansas region, as represented by exposed formations. References to important publications that offer detailed treatment of various rock divisions in Kansas are arranged by main parts of the geologic column.

INTRODUCTION

At the suggestion of geologists in Wichita who were charged with responsibility of arrangements for the annual meeting of the American Association of Petroleum Geologists in Wichita, in March 1935, the State Geologist prepared a tabular description of the rocks of Kansas. This chart was printed on large sheets and was distributed to geologists attending the convention. Additional copies were made available to most geologists and oil companies in Kansas through the agency of the Kansas Geological Society. Eventually the supply was exhausted, and copies were made by photostat or other means of reproduction. The usefulness of this tabulation, as indicated by the demand for it, has been called to our attention on numerous occasions, and the desirability of a revision or new summary on similar lines has been emphasized.

The present contribution differs from the earlier publication in treating only the rocks that are exposed at the surface in different parts of Kansas and in being arranged to go on successive pages rather than on a single large sheet. The omission of descriptions of subsurface rock units—that is, those which are not represented by outcrops anywhere in Kansas—may be con-

sidered to be a serious deficiency, especially from the standpoint of petroleum geologists who are concerned with exploration for oil and gas in Ordovician or other pre-Mississippian rocks of the state. Reason for this omission is the complexity of features belonging to these subsurface rocks, as now known in various parts of Kansas, and the present unreadiness of the authors to deal with these unexposed rocks satisfactorily. It may be possible to summarize subsurface stratigraphy of the Kansas region in a useful manner at a later time. The differentiation of outcropping rocks from those that are not exposed in the state is a convenient, if somewhat arbitrary, basis for limiting the scope of descriptions. The arrangement of material on pages, rather than in chart form, has the advantage of permitting flexible variation in space allotted to items, and the unwieldiness of a large chart is avoided. Effort has been made to preserve tabular features and conciseness, which are attributes of a stratigraphic chart.

MAIN ROCK DIVISIONS AND THEIR DISTRIBUTION

Outcropping rocks of Kansas are of Cenozoic, Mesozoic, and Paleozoic age. They are assigned to seven (or, according to alternative classification, six) geologic systems that embrace a time span of approximately 300 million years. They record the successive invasions of shallow seas that covered much or all of the state during many thousands of years. Some rock layers are of nonmarine origin, having been deposited by streams, air currents, or the still water of lakes. Such formations lack fossilized marine organisms, like those which are so abundant in many sea-laid strata of Kansas; they may contain leaves of land plants or traces of fresh-water life, or air-breathing animals. Also, there are numerous records of more or less widespread and prolonged erosion, when varying quantities of previously formed rocks were removed. These times of denudation indicate an emergent condition of the earth's surface in Kansas. They are defined by the local or regional absence of strata that are recognized elsewhere, and commonly there are irregularities along the contact of the sedimentary units that are separated by a hiatus in deposition. The hiatus is termed a disconformity if adjoining strata lie parallel, and it may be called an unconformity if locally or regionally the rocks on opposite sides of the break show some divergence in structure or if there is evidence that interruption of deposition was very widespread and prolonged. Strictly speaking, however,

the term unconformity embraces disconformity and gentle to strong angular discordance of beds separated by an erosional break.

The distribution of outcrops belonging to all main divisions and representing many subdivisions of the rock column of Kansas is adequately shown on the half-million scale geologic map of the state, on which 1 inch equals approximately 8 miles (Moore and Landes, 1937). A generalized map showing rock outcrops is given in figure 1.

Distribution and structure of Cenozoic rocks.—Deposits of Cenozoic age occurring in Kansas constitute a discontinuous thin veneer that broadly conforms to the east-sloping land surface. Excepting the widespread Tertiary deposits of western Kansas, the Cenozoic sedimentary units are so patchy and irregular that they may hardly be said to exhibit regional structure.

Distribution and structure of Mesozoic rocks.—The outcrops of Mesozoic strata in Kansas are confined to the western two thirds of the state. They extend much farther eastward along the north border of Kansas (Marshall county) than in the south (Barber county); in central Kansas exposures are found eastward as far as Dickinson and Marion counties. Mesozoic rocks underlie thousands of square miles in western Kansas where Cenozoic formations cover the surface. In general, the Mesozoic strata are gently inclined toward the northwestern corner of Kansas, dipping westward, northwestward, or northward.

Distribution and structure of Paleozoic rocks.—The eastern and south-central parts of Kansas contain extensive outcrops of Upper Paleozoic rocks, chiefly Permian and Pennsylvanian. This is a region of roughly parallel belts of plains that are developed at the outcrop of weak rocks and on the upper surface of some gently inclined hard formations. The plains are separated by hilly country that terminates eastward in a more or less prominent escarpment. The general trend of these escarpments is south-southwest across eastern Kansas. The Flint Hills, which are formed by Lower Permian strata, are an example.

At the outcrop, Paleozoic strata in Kansas show a very gentle west or west-northwest dip that ranges mostly from 15 to 35 feet to the mile. Locally, there are dips in other directions or the layers may be quite horizontal; nowhere are the strata steeply inclined. A noteworthy belt of distinct although gentle easterly

dips may be traced across the state from the vicinity of Arkansas City in the south to the neighborhood of Seneca in the north; this defines the east limb of the anticlinal structure that overlies the buried Nemaha Ridge. Drilling along this anticline in south-central Kansas has led to the discovery of important oil and gas fields. Faults having small displacement have been observed in a few places, but nowhere in Kansas is faulting a very noticeable feature of the rock structure.

Igneous rocks.—Although outcropping rocks of Kansas are almost exclusively of sedimentary nature, rocks of igneous origin occur in at least two areas. It is not possible to determine the age of these rocks with any exactness, and consequently they cannot be fitted satisfactorily into a tabular description. In Woodson county boulders of granite-like rock are found in association with partly metamorphosed Pennsylvanian sedimentary rocks. The igneous rock is evidently a dike-like intrusion that invaded the strata some time after Late Pennsylvanian time. In Riley county there are at least three small outcrops, each less than an acre in extent, that furnish record of intrusion of dark-colored igneous rock into Lower Permian strata. The known exposures of this igneous rock are several miles distant from one another. Based on very insecure considerations, it is supposed that the igneous rocks of Kansas just noted may be of Cretaceous age.

CENOZOIC ROCKS.—Formations belonging to the Cenozoic era, last of first rank geologic time divisions, are widespread in Kansas. They comprise Quaternary sediments, which include glacial deposits of northeastern Kansas and nonglacial deposits of Pleistocene and Recent time in all parts of the state, and Tertiary deposits, which cover most of western Kansas.

QUATERNARY SYSTEM.—Deposits of Quaternary age occur widely over Kansas. These are all nonmarine in origin and include glacial, fluvial, and eolian deposits. Glacial sediments occur only in northeastern Kansas, whereas stream-laid deposits occur generally in the western half of the state and along the valleys in the eastern half. The maximum aggregate thickness of these deposits is nearly 700 feet, but inasmuch as the maximum thickness of several units is never found in the same area the greatest thickness of Quaternary deposits known at any locality is 275 feet.

RECENT SERIES (including some undifferentiated Pleistocene deposits).—Sedimentary deposits formed since the time of disappearance of the last glacial ice sheet from North America are classed as belonging to the Recent series. They are chiefly stream-laid deposits in continental areas such as Kansas, but on borders of continents they include marine sediments. Inasmuch as the last ice sheet did not reach Kansas and conditions that are typical of Recent time began in this region approximately in mid-Pleistocene time, the Pleistocene-Recent boundary in Kansas is not clearly recognizable at some places.

☐ **Alluvium.**—Gravel, sand, silt, and loam; gray to tan; underlying the valley flat of most stream valleys throughout the state. In western Kansas and along the through valleys of central and eastern Kansas the alluvium consists of sediments derived from the Rocky Mountain region mixed with local materials. In northeastern Kansas the alluvium is largely derived from glacial deposits; in southeastern Kansas it consists predominantly of chert gravel and sand and other material derived from the Paleozoic rocks of the region. Alluvium is generally thicker under major valleys and ranges in thickness from a few feet to more than 80 feet. Between Hutchinson and Wichita the alluvium of the Arkansas river valley attains a known thickness of 275 feet

☐ **Dune sand.**—Eolian sand, well-sorted, predominantly quartz; gray to tan; mantling the surface of much of central and southwestern Kansas, especially in the Great Bend area, south of Arkansas river. The dune sand in many areas represents several cycles of dune development. The younger cycles are Recent in age and consist of loose sand, whereas the older cycles are Pleistocene in age and in some places are red-brown and are partly indurated. Maximum thickness, about 50 feet

☐ **Loess.**—Eolian silt, well-sorted; gray, tan, buff, and red-brown. Widespread in western and central Kansas. Surficial loess is still in the process of deposition in some areas of the state, and in some places a thin mantle of Recent material overlies more widespread deposits of Pleistocene loess. Includes the upper part of the Kingsdown silt and the Sanborn formation. Maximum known thickness, about 25 feet

~~~~~? Unconformity ?~~~~~

CENOZOIC

QUATERNARY

Recent and Pleistocene

**PLEISTOCENE SERIES.**—The distinctive sedimentary deposits formed in Pleistocene time, which comprises the older and longer part of the Quaternary system, are glacial materials. Some glacial deposits, both those formed directly by ice work and those resulting from action of melt-water derived from ice sheets,

### EASTERN KANSAS

□ **Loess.**—Eolian silt, massive, tan. Caps the uplands of northeastern Kansas, especially along the Missouri river valley. Maximum thickness, about ..... 35 feet

□ **Terrace deposits.**—Sand, gravel, boulders, and silt; forming terraces along major stream valleys. In northeastern Kansas the terrace deposits are derived principally from glacial sediments and contain ice-carried pebbles and boulders of igneous rocks and quartzite. In south-eastern Kansas these deposits consist of silt, clay, sand, and chert pebbles derived mainly from near-by Paleozoic rocks and high-level Tertiary deposits. Maximum thickness, about ..... 80 feet

■ **Kansan till.**—Glacial till in northeastern Kansas, coarse-textured, brown and tan; containing boulders of rocks foreign to Kansas, especially pink quartzite. Thickness, a featheredge to ..... 75 feet

■ **Aftonian interglacial deposits.**—Sand and silt in Atchison and adjacent counties; thin-bedded and thinly cross-bedded; tan and gray. Maximum thickness, in eastern Atchison county, ..... 70 feet

■ **Nebraskan till.**—Glacial till in northeastern Kansas, fine-textured, blue-gray, contains few pebbles or boulders. Thickness, a featheredge to a maximum, near the Missouri valley, of ..... 80 feet

### CENTRAL KANSAS

□ **Loess.**—Eolian silt, well-sorted; tan, brown, and gray; locally, volcanic ash at base. Widespread in McPherson and Republic counties and mantles the uplands at scattered localities. Maximum thickness, in McPherson county, ..... 110 feet

■ **Gerlane formation.**—Clay, silt, sand, and gravel in south-central Kansas. Maximum thickness, in Barber county, 104 feet; thickness commonly ..... 60 feet

□ **Terrace deposits.**—Sand, gravel, silt, clay, and volcanic ash; tan and gray. Mantles terraces of different ages which occur at heights of 10 to 100 feet above the flood plains of valleys of central Kansas, and underlies the floors of several abandoned high-level valleys. Some local terrace deposits have been called "Salt Creek gravel beds," and some deposits called "Abilene conglomerate" may be terrace deposits. Snails locally abundant. Representative mammals from terraces of three ages include: (1) *Cynomys*; (2) *Castoroides*, *Neofiber*, *Blarina*; (3) *Sorex*, *Microtus*, *Pitymys*. Thickness, thin veneer to ..... 80 feet

■ **McPherson formation.**—Sand, gravel, and silt; gray and tan. Contains bones of mammals: *Megalonyx*, *Paramylodon*, *Smilodon*, *Mastodon*, *Elephas*, *Parelephas*, *Equus*, and *Camelops*. Thickness, a featheredge to ..... 150 feet

■ **Belleville formation.**—Sand, gravel, silt, and clay, in Republic county. Contains *Equus* and *Stegomastodon*. Maximum thickness more than ..... 150 feet

~~~~~ Major unconformity ~~~~~

~~~~~ Unconformity ~~~~~

CENOZOIC

QUATERNARY

Pleistocene



[PLEISTOCENE SERIES, Continued] occur in Kansas, but they belong to the early and middle part of the Pleistocene epoch. Outside the glaciated area in early Pleistocene time and throughout the state during the latter part of the Pleistocene epoch, conditions resembled those of Recent time.

### SOUTHWESTERN KANSAS

■ **Kingsdown silt.**—Silt and fine sand, containing nodules and bands of caliche; tan to gray. Thin-bedded fine sand at base grading upward into loess (Recent in upper part). Mantles uplands and covers floor of the Meade basin. Mammal remains include *Paramylodon* and *Equus* in lower part and *Cynomys* in upper part. Thickness, typically 10 to 45 feet, but attaining a maximum of more than 90 feet in southern Ford county; commonly ..... 40 feet

□ **Terrace deposits.**—Gravel, sand, silt, and clay; mantling terrace surfaces at several levels along the Cimarron, Arkansas, and adjacent valleys. The deposits consist largely of pebbles and grains of Rocky Mountain rocks and abraded caliche and "mortar bed" pebbles. Contain bones of *Parelephas* and *Equus*. Thickness ..... 5 to 65 feet

■ **Meade formation.**—Gravel, sand, silt, clay, volcanic ash, and caliche; gray, tan, red, pink, and buff; selenite crystals prominent in redbeds. Gravel and sand contain pebbles derived from the Rocky Mountain region and from near-by Pliocene and older rocks. Channel deposits rest unconformably on the lower, more widespread, part. Well exposed in the Meade basin. Snails locally abundant. Five distinct vertebrate zones have been recognized. The most typical mammals from these zones, in descending order, are: (1) *Ambystoma*, *Sorex*, *Microtus*; (2) *Paramylodon*, *Equus*, *Camelops*; (3) *Sigmodon*, *Onychomys*, *Perognathus*; (4) *Sorex*, *Microtus*, *Pitymys*; (5) *Equus*, *Camelops*. Thickness, commonly ..... 10 to 150 feet

### NORTHWESTERN KANSAS

■ **Sanborn formation.**—Consists mainly of loess; sand and, locally, gravel occur commonly at the base; generally well-sorted, tan to gray-buff. Mantles the uplands and highest terrace levels. Snails are abundant locally. From Jewell county westward to Decatur county a prominent soil zone that has yielded remains of *Citellus* and *Cynomys* occurs at the base of the gray-tan loess. Below the soil zone are heterogeneous deposits of clay, silt, sand, and gravel, usually red to red-brown in color. These deposits have yielded remains of the fossil vertebrates *Citellus* and *Cynomys*. Thickness ..... 5 to 100 feet

~~~~~Local unconformity~~~~~

~~~~~Unconformity~~~~~

**TERTIARY SYSTEM.**—Deposits of Tertiary age occur widely in central and western Kansas, and locally in eastern Kansas. These deposits are all continental in origin and were deposited mostly by streams and in lakes. The source area of most of these deposits was to the west, principally the area of igneous rocks in the Rocky


**PLIOCENE SERIES.**—The Tertiary rocks of Kansas, so far as is known, are all included within the Pliocene series.

**EASTERN KANSAS**

□ **Chert gravels.**—Gravel and sand; consist predominantly of poorly sorted chert gravels, brown to gray-tan, derived principally from the Flint Hills region of east-central Kansas. Underlie the surface of abandoned high-level valleys and high terraces. In northeastern Kansas these chert gravel deposits underlie the Nebraskan and Kansan tills. Pliocene or older in age. Thickness, thin veneer to ..... 30 feet

**CENTRAL KANSAS**

■ **Emma Creek formation.**—Sand, gravel, sandy silt, and clay; brown, buff, and gray. Consists largely of pebbles and grains derived from adjacent Cretaceous rocks. Widespread in McPherson and adjacent counties. These deposits were probably deposited by streams flowing toward the south and southwest at a time prior to the establishment of east drainage across the Flint Hills region. A prominent disconformity occurs within the formation and the lower (older) part is probably equivalent in part to the Ogallala formation of western Kansas. The age of the beds above this disconformity is in doubt. They may prove in part to be of Pleistocene age. Mammal remains from the beds below the disconformity include *Amebelodon*, *Tetralophodon*, *Neohipparion*, *Pliohippus*, *Nannippus*, and *Teleoceras*. Thickness, less than 10 to 180 feet; commonly ..... 40 feet



~~~~~ Major unconformity ~~~~~ ~~~~~ Major unconformity ~~~~~

[**TERTIARY SYSTEM, Continued**] Mountain region and the area of sedimentary rocks in eastern Colorado and western Kansas. The thickness of these deposits ranges from a thin veneer to about 350 feet, but the maximum thickness has nowhere been observed at the surface.

SOUTHWESTERN KANSAS

■ **Ogallala formation.**—Gravel, sand, silt, clay, and caliche; pink, gray, and tan. Widespread over western Kansas. Thickness, maximum, 350 feet, commonly 125 feet

● **Rexroad member.**—Sand, silt, clay, caliche, and locally peat; gray, tan, and pink. Occurs in the Meade basin and vicinity. Contains mammals: *Parahodomy*, *Sigmodon*, *Ogmodontomys*, *Plesipus*, and *Nannippus*. The exact age of these beds within the standard time scale is in doubt. Snails locally abundant. Thickness (maximum thickness known only from well samples), less than 50 to 250 feet

○ **Unnamed middle Pliocene member.**—Gravel, sand, sandy silt, and caliche; massive to cross-bedded; pink, tan, and gray. Contains fossil mammals *Amebelodon*, *Plihippus*, and *Aphelops*, and fruits of the herb *Biorbia fossilia*. Thickness, 10 to 150 feet, commonly 100 feet

In Meade and Seward counties an angular unconformity at base.

~~~~~ Local unconformity ~~~~~

■ **Laverne formation.**—Shale, sandstone, and limestone; buff, blue-gray, and gray. Silty chalky limestone contains ostracodes and diatoms. Snails locally abundant in shales. Mammal remains include *Nannippus*, *Calippus*, and *Eucastor*. Occurs in Meade and Seward counties, locally truncated by overlying Ogallala. Lower Pliocene and possibly upper Miocene in age. Exposed thickness 60+ feet; from well logs believed to attain a maximum thickness of more than 60 feet

~~~~~ Major unconformity ~~~~~

NORTHWESTERN KANSAS

■ **Ogallala formation.**—Gravel, sand, silt, and caliche; massive and cross-bedded; contains pebbles of igneous rocks (derived from the west), limestone, and sandstone; pink, gray, and tan. Gravel and sand locally cemented to form "mortar beds." Widespread under the High Plains of western Kansas. Locally capped by hard limestone or caliche bed containing algal-like structures. Contains Edson Quarry fauna and *Biorbia fossilia* fruits. Middle Pliocene in age. Green and red-brown bentonitic clay beds locally occur in the base of the Ogallala, and in Wallace county this zone has been called the *Woodhouse clay*. Locally this formation unconformably overlies undifferentiated lower Pliocene deposits (Norton, Rawlins, Phillips, and Trego counties) that have yielded bones of the fossil beaver, *Eucastor*. Thickness, 10 to 200 feet, commonly 150 feet

~~~~~ Major unconformity ~~~~~




FIG. 2. Generalized sections of Cenozoic rocks in Kansas.

MESOZOIC ROCKS.—Deposits of Mesozoic age comprising part of the geologic section of Kansas belong mostly to the Cretaceous system. These rocks cover many thousand square miles in the western part of the state. Older Mesozoic rocks, probably representing both the Jurassic and Triassic systems, are identified in well borings but are absent or, in the case of Triassic, somewhat doubtfully recognized very locally in surface exposures.

CRETACEOUS SYSTEM.—Rocks of Cretaceous age crop out at the surface or underlie much of central and western Kansas. These rocks are mostly marine, but deposits of supposed continental origin are found in the Cheyenne sandstone and Dakota formation. Clayey and calcareous shale is the dominant constituent, but fine-grained platy and chalky limestone comprises most of the Greenhorn and Niobrara, and sandstones occur in the Cheyenne, Kiowa, Dakota, Graneros, and Carlile. In general, these rocks dip very gently westward, but locally dips may occur in any direction. Small outcrops of dark-colored igneous rocks that penetrate Permian strata in Riley county are possibly Cretaceous in age. The thickness of the Cretaceous system in Kansas is about 2,750 feet.

GULFIAN SERIES.—The upper part of the Cretaceous system is defined as the Gulfian series. At the beginning of Gulfian time, continental and littoral deposits accumulated in Kansas, but these conditions rapidly gave way to marine conditions and the shales and limestones constituting the Colorado and Montana groups were deposited. Thickness about 2,500 feet.

Montana group.—Dark-colored marine shales that crop out and underlie the surface of northwestern Kansas. Thickness, 1,000 to 1,400 feet; average 1,235 feet.

■ **Pierre shale.**—Shale, thin-bedded, black to dark gray, a few beds lighter in color, weathers to coffee-brown and gray; marine; contains concretions, selenite crystals, thin beds of bentonite, and locally chalky beds. Occurs in northwestern Kansas. Thickness, 1,000 to 1,400 feet; average 1,235 feet

● **Beecher Island shale member.**—Shale, gray. Irregular concretionary limestone near top, limonite concretions throughout, thin beds of bentonite and limestone concretions in lower part. Characterized by the marine clams *Tardinacara (Pseudoptera) fibrosa* and *Inoceramus sagensis*, and the cephalopods *Baculites grandis* and *Discoscaphites abyssinus*. Thickness 100 feet

○ **Unnamed shale member.**—Shale, black to gray. Thickness, 500 to 600 feet, average 550 feet

● **Salt Grass shale member.**—Shale, clayey, gray, containing numerous thin bentonite beds, limestone concretions, and concretionary limonite zones distributed throughout. Contains *Baculites pseudovatus* and *Acanthoscaphites nodosus*. Thickness 60 feet

● **Lake Creek shale member.**—Shale, thin-bedded, flaky, dark gray and black; limestone concretions, zones of concretionary limonite, and locally gypsum are present. Contains *Baculites compressus*, *Serpula wallacensis*, and *Acanthoscaphites nodosus*. Thickness 200 feet

MESOZOIC
CRETACEOUS
Gulfian
Montana

Montana

- **Weskan shale member.**—Shale, clayey, gray. Bentonite beds more abundant in lower part, large limestone concretions and some limonite. Contains *Serpula wallacensis*, *Acanthoscaphites nodosus*, *Anomia subtrigonalis*, *Ostrea*, and *Crassatella evansi*. Thickness170 feet
- **Sharon Springs shale member.**—Shale, flaky, black, somewhat bituminous, large septarian and ordinary limestone concretions abundant in upper part; a few beds of light-gray shale. Thickness, 155 feet

Colorado group.—Shales, calcareous and noncalcareous, that crop out and underlie the surface of north-central, northwestern, and west-central Kansas. Limestone beds occur interbedded with the calcareous shales. These rocks are all of marine origin. Thickness, about 1,050 feet.

- **Niobrara chalk.**—Calcareous shale and chalk, soft, interbedded, light gray. Crops out in a belt trending generally northeast-southwest and extending from north-central to western Kansas. Thickness, 500 to 750 feet in Logan and Wallace counties; average 605 feet

- **Smoky Hill chalk member.**—Shale, chalky, interbedded, gray, weathers white, yellow, and orange; contains limonitic concretions. Forms badland topography. Characterized by the marine clams *Inoceramus (Haploscapha) grandis* and *Ostrea congesta*. Thickness, 450 feet, to 700 feet in Logan county; average 550 feet

- **Fort Hays limestone member.**—Chalk or chalky limestone, gray to cream-colored, massively bedded; thin beds of light to dark-gray chalky clay shale separate the massive chalky limestone beds. Extensively used for building stone. Contains *Inoceramus deformis*. Thickness, 50 feet in Phillips county, to 65 feet near the Colorado line; average 55 feet

- **Carlile shale.**—Shale, chalky in lower part and containing thin chalk beds near the base, black fissile shale and large septarian concretions in upper part, fine-grained sandstone at top; marine. Occurs in northwestern and western Kansas. Thickness about 300 feet

- **Codell sandstone member.**—Sandstone, fine-grained, and siltstone, somewhat calcareous, gray and tan; sharply defined at top but grading downward into dark-gray shale. Thickness, 3 feet in Ness county, 22 feet in northern Ellis county, to 25 feet in Hamilton county; average 15 feet

- **Blue Hill shale member.**—Shale, clayey, gray-black to dark gray, noncalcareous; abundant ordinary and septarian concretions and selenite crystals. Thickness, 75 feet in Hamilton county to 200 feet in Russell county; average 160 feet

- **Fairport chalky shale member.**—Shale, calcareous, and thin chalk beds, blue-gray to gray, weathers to light orange-tan; chalky limestones more abundant near base. Contains *Ostrea congesta*. Thickness, 85 feet in Russell county to 147 feet in Hamilton county; average 125 feet

- **Greenhorn limestone.**—Chalky limestone and calcareous shale, interbedded, thin-bedded, light gray to dark gray, weathers yellow-gray to light gray; marine. Occurs in northwestern and western Kansas. Thickness, 85

MESOZOIC

CRETACEOUS

Gulfan

Colorado

feet in northern Barton county, 95 feet in Ellis county, to 132 feet in Hamilton county; average 100 feet

● *Pfeifer shale member*.—Chalky shale and chalky limestone in alternating layers, "Fencepost limestone bed" at top, blue-gray, weathers to light-tan. In Hamilton county the Pfeifer shale member and the underlying Jetmore chalk member are thicker than farther east. They cannot be distinguished and have been together designated the *Bridge Creek limestone member*, having a total thickness of 74 feet. Thickness of Pfeifer shale is typically 19 to 21 feet in Ellis and Russell counties; average 20 feet

● *Jetmore chalk member*.—Chalky shale and chalky limestone, interbedded, gray, weathers to light gray. Contains *Inoceramus labiatus*. Thickness, 20 feet (Russell and Ellis counties) to 25 feet; average 22 feet

● *Hartland shale member*.—Chalky shale, a few thin beds of chalky limestone and bentonite, gray. Thickness, 23 feet in Kearny county to 35 feet in Ellis county; average 30 feet

● *Lincoln limestone member*.—Chalky shale and chalky limestone, interbedded, light gray; beds of dark-gray hard crystalline limestone at base and top; shale contains thin beds of bentonitic clay. Member weathers to yellow-gray or yellow-tan. Thickness, 20 feet (Ellis county) to 35 feet; average 28 feet

■ *Graneros shale*.—Shale and clay shale, fissile, noncalcareous, blue-black, weathers to dark gray and coffee-brown; largely or entirely marine but mostly unfossiliferous; locally contains sandstone beds and beds of "clay ironstone"; selenite crystals abundant. Thickens southward across western Kansas. Thickness, 30 to 35 feet in Russell county to 65 feet near the Colorado line; average 45 feet

The rocks which comprise the Dakota formation, Kiowa shale, and Cheyenne sandstone formerly were classed as the Dakota group. The use of this grouping has been discontinued.

■ *Dakota formation*.—Clay, shale, siltstone, and sandstone, interbedded and lenticular; contains carbonaceous material, lignite, concretions of hematite and limonite, and locally quartzitic sandstone; white, gray, red, brown, and tan. Occurs in north-central and western Kansas. The *Cockrum sandstone* of southwestern Kansas is equivalent in age to part of the Dakota formation. Contains stratigraphic units formerly called "Rocktown channel sandstone," "Ellsworth formation," "Solomon formation," "Reeder sandstone," "Marquette sandstone," "Spring Creek clay," and others. Contains plant fossils and land vertebrates. Thickness, 100 to 300 feet; average 215 feet

● *Janssen clay member*.—Clay, silt, and fissile shale, with lenticular sandstone, lignite, and lignitic clay common; gray to dark gray, contains concretionary hematite and limonite and beds of "ironstone." Central and north-central Kansas. Thickness, 30 to 80 feet; average 55 feet

● *Terra Cotta clay member*.—Clay, shale, sandstone, and quartzitic sandstone, interbedded, red, gray, brown, and tan. Central and north-

Colorado

Gulfian

MESOZOIC

CRETACEOUS

Gulfian

central Kansas. Sandstones are lenticular and weather brown, coarse-grained to fine-grained, abundant hematite and limonite concretions throughout. Quartzitic sandstone near top. Thickness, 70 to 220 feet; average 160 feet

COMANCHEAN SERIES.—The lower part of the Cretaceous system is defined as the Comanchean series. At the beginning of Comanchean time the Kansas area was being eroded and only upper Comanchean deposits occur in the state. The oldest Comanchean deposits in Kansas represent continental and littoral deposits laid down as the sea advanced northward. These nonmarine conditions rapidly gave way to marine conditions, and marine shales overlie the nonmarine and littoral deposits. These rocks crop out in a belt extending diagonally north to south across central Kansas, and at a few places in southwestern Kansas. Thickness, featheredge to 400 feet; average about 250 feet.

■ **Kiowa shale.**—Shale, fissile, light gray, dark gray, and black, contains thin limestone beds throughout, with the *Champion shell bed* at the base in the type area. Locally, lenticular sandstones occur at any position within the shale; selenite crystals common. Abundant marine molluscan fauna in the thin limestones, shale, and sandstone. Probably contains stratigraphic units formerly called "Greenleaf sandstone," "Mentor beds," and others, and is in part equivalent to the "Belvidere formation," "Medicine bed," "Elk River beds," and others. Thickens across central and western Kansas toward the south and southwest. Typical thickness, 60 to 150 feet; average .. 100 feet

■ **Cheyenne sandstone.**—Sandstone, white, buff, gray, tan, red, yellow, and purple, with white, light gray, and buff dominant; and shale, light gray to dark gray. Sandstone, very fine-grained to coarse-grained, dominantly quartz, cross-bedded; shale, sandy and silty, cross-bedded, dark gray, contains plant remains. Cobble zone at base, locally conglomeratic; locally a zone of redeposited red Permian sand at base. Contains stratigraphic units formerly called "Stokes sandstone," "Lanphier beds," and "Natural Corral sandstone." Thickens westward in the subsurface. Maximum thickness exposed, more than 75 feet; maximum thickness in the subsurface, 300 feet; average 150 feet

A major unconformity marks the base of the Cretaceous rocks. The Cretaceous rocks overlap northward on the pre-Cretaceous erosion surface, successively younger beds lying above the contact northward across the state, ranging from the Cheyenne sandstone in Comanche and Kiowa counties to the Dakota formation in Washington county. Rocks underlying the unconformable contact range in age from the Herington limestone in Washington county to beds believed to be of Jurassic age in northwestern Kansas.

Major unconformity

(?) **TRIASSIC SYSTEM.**—Continental deposits of early Mesozoic age, identified as Triassic on the basis of vertebrate and plant remains, are widespread in western Texas and New Mexico. Mainly on lithologic grounds, certain outcrops in the panhandle of Oklahoma and southwestern Kansas are believed to be equivalent to the undoubted Triassic of areas farther south. Thickness in Kansas about 20 feet.




FIG. 3. Generalized section of Cretaceous rocks in Kansas.

| | | |
|----------|------------|--------------|
| MESOZOIC | | (?) |
| | | TRIASSIC (?) |
| | Dockum (?) | |

(?) **Dockum group.**—Inasmuch as the Triassic rocks above mentioned, belonging to the late part of the period, are called Dockum group in Texas, this name is tentatively employed for the supposed Triassic of southwestern Kansas. Thickness, 20 feet.

□ **Unnamed formation.**—Red siltstone, buff and white sandstone, and a small amount of gypsum, exposed at two places in the Cimarron valley in Morton county, are differentiated from other rocks of the region. They are believed to correspond to deposits in Texas county, Oklahoma, that are identified as Triassic. The maximum exposed thickness of these rocks in Kansas is about 40 feet but locally beneath the surface their thickness seemingly is greater; average 20 feet

Regional unconformities and overlaps bring Cenozoic and Mesozoic rocks into contact with various Paleozoic formations.

Unconformity

PALEOZOIC ROCKS.—Sedimentary, igneous, and metamorphic rocks of all sorts formed during the great span of geologic time called the Paleozoic era have no classificatory designation that is generally used to refer to them collectively. Somewhat loosely, therefore, we term them Paleozoic rocks. The outcropping rocks throughout most of eastern and central Kansas are all sedimentary strata of Late Paleozoic age. They include shale, sandstone, limestone, and conglomerate having widely varied characteristics, as well as coal beds, dolomite, and some other rock types. Paleozoic rocks underlie all of western Kansas. Locally exposed igneous rocks in Woodson county may be Paleozoic but they are probably much younger. The total thickness of exposed Paleozoic rocks in Kansas, derived by compiling average measurements of recognized subdivisions, may be given as approximately 6,000 feet.

PERMIAN SYSTEM.—Rocks of Permian age crop out in eastern Kansas in a belt extending from Washington, Marshall, Nemaha, and Brown counties, on the northern boundary, to Meade, Clark, Comanche, Barber, Harper, Sumner, and Cowley counties, on the Kansas-Oklahoma line. The total outcrop thickness is about 3,000 feet.

GUADALUPIAN SERIES.—This division of the Permian system, which is defined from fossiliferous marine rocks in western Texas and southeastern New Mexico, crops out in southern Kansas. In this region, it comprises unfossiliferous deposits that seem to have been laid partly on land by sluggish streams and action of winds and partly in shallow basins occupied by strongly saline waters. Bedding is mostly irregular. Reddish color, which signifies abundance of ferric oxide, prevails. Thickness of outcropping rocks belonging to this series in Kansas is about 290 feet.

Quartermaster group.—Permian beds above the Day Creek dolomite have been classified as belonging to the Quartermaster group. They seem to be equivalent to the lower part of the Quartermaster formation of western Oklahoma and the panhandle region of Texas. The maximum outcrop thickness of these rocks in Kansas is about 45 feet.

Quartermaster

■ **Taloga formation.**—Redbeds of silty and sandy shale and sandstone, called "Big Basin" in some reports, belong to the Taloga formation. The lower 25 feet is chiefly silty shale. Outcrops are in western Clark and eastern Meade counties. The strata are seemingly equivalent, all or in part, to the Quartermaster formation in Oklahoma. Thickness, maximum about 45 feet; average 20 feet

Beds below the Taloga formation and above the Dog Creek shale are not assigned to a group.

■ **Day Creek dolomite.**—Fine-grained dense dolomite cropping out in western Clark county. It is seemingly absent between the northern part of T. 33 S., R. 24 W. and a point in Oklahoma near the center of T. 25 N., R. 25 W. (Indian meridian). Thickness in Kansas is about 2 feet

■ **Whitehorse sandstone.**—Redbeds of sandstone, siltstone, and shale, and a minor amount of dolomite. Cross bedding is common and the formation is characterized by "sand balls," which are water-worn small "sand crystals" consisting of sand pseudomorphs after calcite, and by crystals of barite, anhydrite, calcite, and gypsum. The outcrops are in southeastern Meade, Clark, and southwestern Kiowa counties. Thickness is about 270 feet

O *Upper shale member.*—Redbeds of shale and a minor amount of sandstone, with a zone of dolomitic beds in the basal part and a zone of gray-green sandy shale in the upper part, mostly brick-red or maroon. Thickness is about 38 feet

O *Even-bedded member.*—Sandstone and shaly siltstone, mostly even-bedded but locally cross-bedded in the upper part; containing "sand balls" and "sand crystals"; maroon. Thickness is about, 100 feet

● **Relay Creek? dolomite and sandstone member.**—Two beds of dolomite separated by red and white sandstone of variable texture. The dolomite beds range in thickness from a few inches to about 1 foot. Locally the dolomite beds are altered to anhydrite or gypsum. Individual layers in the sandstone are identified with difficulty; locally the rock is cross-bedded. Thickness is about 22 feet

● **Marlow sandstone member.**—Redbeds of sandstone, fine-grained, locally shaly or silty, cross-bedded. "Sand balls" locally prominent. Thickness about 110 feet

Conformity

LEONARDIAN SERIES.—This major division of the Permian, named from strata in the Glass Mountains region of western Texas, is judged to be represented in Kansas by about 1,900 feet of rocks that are chiefly unfossiliferous clastics (sandstone, shale) and evaporites (anhydrite, gypsum, salt). Red shale, siltstone, and sandstone predominate in the upper part. Gray shale is the most common rock type in the lower part, although red and other bright colors are present.

Leonardian

■ **Nippewalla group.**—The upper part of strata assigned to the Leonardian series, named from Nippewalla river in Kingman county, is most widely exposed in south-central Kansas west of Arkansas river. This group consists mostly of redbeds that form a plain. In Barber county and adjacent areas the topography

of the Nippewalla outcrop area locally has considerable relief and is "badland" in type. Gypsum beds make prominent escarpments. The total thickness of the group is approximately 930 feet.

PALEOZOIC
PERMIAN
Leonardian
Nippewalla

■ **Dog Creek shale.**—Maroon shale, sandstone, thin layers of dolomite, dolomitic sandstone, and gypsum. The top generally is marked by about 3 feet of maroon shale, but locally a gypsum bed about 1 foot thick and having red stripes occurs at the top. The most persistent part is a bed of white and red sandstone, about 6 feet thick, that locally is capped by dolomitic sandstone, which occurs next below the upper maroon shale portion. The outcrops are in southern Kiowa, eastern Comanche, and western Barber counties. In Kiowa county the formation is overlapped by Cretaceous sandstone (Cheyenne). The reported thickness ranges from 14 to 53 feet; average about 35 feet

■ **Blaine formation.**—Gypsum beds separated by dolomite and red shale. The formation is divided into four members. Outcrops are in Clark, Comanche, Barber, and Pratt counties. The thickness is about 50 feet

● **Haskew gypsum member.**—One foot or less of gypsum underlain by about 5 feet of red shale. The gypsum bed has been removed by solution at many places. The thickness of the gypsum and underlying shale beds is about 6 feet

● **Shimer gypsum member.**—A thick bed of gypsum overlying a bed of dolomite that ranges in thickness from about 6 inches to 1½ feet. Excessive solution and erosion of the gypsum bed have greatly reduced its thickness in many places. Measured sections believed to show the original thickness range from 14 to 24 feet; average 19 feet

● **Nescatunga gypsum member.**—A bed of gypsum with overlying and underlying red shale. The gypsum bed varies in thickness within short distances and is known to range from about 2 to 8 feet. Locally as much as 8 feet of red shale separates the gypsum from the next higher and lower gypsum beds. Average 5 feet

● **Medicine Lodge gypsum.**—The thickest bed of gypsum in Kansas. Ordinarily there is a bed of dolomite at the base which ranges from 6 inches to 1 foot in thickness. Exposures are in Barber, Clark, and Comanche counties. This gypsum bed forms a conspicuous rim rock at the top of steep slopes of the Flowerpot shale. The maximum thickness is 30 feet or more; average 20 feet

■ **Flowerpot shale.**—Shale, red, soft, gypsiferous. A thin lenticular bed of dolomite has been observed in the middle part, and the formation is cut by intersecting veins of satin spar gypsum. Outcrops are in Barber, southeastern Kiowa, and Comanche counties. Commonly stands in steep slopes, eroded into innumerable gullies, and strewn with clear, white, pink, and red satin spar and clear crystals of selenite. Measured thicknesses range from 170 to 190 feet, average 180 feet

■ **Cedar Hills sandstone.**—Sandstone and siltstone, chiefly red, containing beds of white sandstone in the upper and lower parts; the upper one contains "snow balls" of white gypsum. Shaly siltstone separates the more resistant and more massive sandstones. The upper part below the "snow ball"-bearing sandstone commonly is eroded into a badland plateau. The more

massive sandstones are generally weathered into rounded hills and the shaly portions are weathered into canyon-dissected slopes. Outcrops are in Barber and Harper counties. Thickness is about 180 feet

■ **Salt Plain formation.**—Chiefly red, flaky, silty shale and some siltstone and sandstone. There are two prominent sandstone beds, the upper, about 25 feet thick, occurring about 42 feet below the top of the formation and the lower (*Crisfield*), about 29 feet thick, occurring about 115 feet below the top of the formation. The sandstones are partly cross-bedded. Outcrops are in southeastern Barber, Harper, and southern Kingman counties where the formation has weathered into a nearly featureless plain. In Kingman county the formation is overlapped by Cenozoic deposits. The thickness is reported by George H. Norton to be 265 feet

■ **Harper sandstone.**—Chiefly red, divided into two members. The outcrops are in Harper, Kingman, Reno, and Rice counties. Northward the formation is overlapped by Cenozoic rocks. Thickness is about 220 feet

● **Kingman sandstone member.**—Red sandstone and a few beds of red shale and white sandstone. A prominent bed of white sandstone, 3 feet thick, occurs at the base. Thickness is about 80 feet

● **Chikaskia member.**—Sandstone and shale, mostly red but some gray; fissile, fine-grained, ripple-marked and locally cross-bedded sandstone in the lower part. Some white sandstone and dolomite lenses and concretions occur in the upper part. Along the outcrop in Kansas the formation thins northward, and the thickness ranges from about 100 to 160 feet; average 140 feet

Sumner group.—This division comprises about 1,000 feet of strata at the outcrop, chiefly shale. Thick beds of salt occur in the subsurface. Gray shale predominates but there are also beds of red and green shale and deposits of dolomite, limestone, gypsum, and anhydrite.

■ **Stone Corral dolomite.**—Dolomite, anhydrite, and gypsum. The anhydrite and gypsum portions are lost in exposed sections and the remaining dolomite ledge is cellular or contains numerous calcite-filled vugs. The color is chiefly gray but locally there are red and pink streaks. At some exposures the formation is chiefly red shale, bounded above and below by thin dolomite beds. Ripple marks on dolomite slabs are more or less characteristic. The outcrop belt is interrupted in eastern Rice, western McPherson, and other counties to the south by eastward overlap of Cenozoic deposits. This is one of the most readily recognized "key beds" in the Kansas redbed section. It is reported to produce well-marked reflections in seismograph surveys. Maximum measured thickness at the outcrop is about 6 feet

■ **Ninnescah shale.**—Predominantly shale, mostly red, but containing some gray shale, impure limestone, and calcareous sandstone. The *Runnymede sandstone*, 7 to 8 feet thick, forms the upper part of the formation. G. H. Norton has found that seven other distinctive beds of sandy or calcareous material can be traced for long distances. *Estheria* shells are rather common. Some of the beds show ripple marks. Rosette-shaped calcareous concretions occur in the middle part. Weathering and erosion in this part have produced the "Red Jaw" country in Reno county. The formation thins north-

Nippewalla

Leonardian

Sumner

PALEOZOIC
PERMIAN

ward. In the subsurface near the Nebraska line it is about 50 feet thick. The maximum outcrop thickness is about 450 feet; average thickness at outcrop, about 300 feet

■ **Wellington formation.**—Chiefly shale, with a few hundred feet of salt in the middle part in the subsurface. Outcrops contain several more or less lenticular beds of gypsum and impure limestone. The *Milan limestone member*, consisting of 1 foot of greenish-gray shaly limestone that on the outcrop is characterized by bright-green copper carbonate, occurs at the top and thus marks the top of the Wellington formation in a comparatively large area. Elsewhere change in color from gray to red may be regarded as the upper boundary, which may be at different horizons in different places. The upper 300 feet of the Wellington is largely gray shale with thin interbedded deposits of calcareous material. The *Hutchinson salt member* occurs in the middle part but is not exposed. The fossil insect-bearing *Carlton limestone member* occurs a short distance below the Hutchinson salt. Bright red and green shale is conspicuous in the lower part, which contains more or less discontinuous beds of impure limestone and gypsum. A bed of impure dolomitic limestone, not definitely known to be of widespread occurrence, has been named the *Hollenberg limestone*. In general, outcrops of the Wellington formation are poor and discontinuous. The total thickness of the formation is about 700 feet

Conformity

WOLFCAMPIAN SERIES.—This division, formerly called the Big Blue series, contains the older Permian rocks of Kansas. Shales and limestones predominate, subdivisions being remarkably persistent. The thicker shales are bright colored and the limestones are mostly light gray bleached-bone color. The outcrop thickness in Kansas is about 835 feet.

Chase group.—The topmost group of Wolfcampian beds is made up of about 370 feet of prominent escarpment-making limestones and shales. The shale units are characterized by bright reds and greens. Flint-bearing limestone forms an important part of this division, which largely comprises the rocks in which the Flint Hills are developed from Washington, Marshall, and Nemaha counties on the northern boundary of Kansas to Cowley county on the Kansas-Oklahoma line.

■ **Nolans limestone.**—This formation consists of upper and lower limestone members separated by shale. The thickness ranges from about 22 to 40 feet; average 34 feet

● **Herington limestone member.**—Limestone and dolomite, yellowish-tan, soft and dense, more dolomitic in southern Kansas than in the northern and central parts of the state. Outcrops are characterized by siliceous and calcareous geodes and concretions and cauliflower-like masses of drusy flint weathered from the matrix. Fossil mollusks are locally abundant. Thickness ranges from about 7 to 10 feet in the northern part and about 30 feet in the southern part of Kansas; average 20 feet

Leonardian
Sumner

PALEOZOIC
PERMIAN

Wolfcampian
Chase




FIG. 4. Generalized section of upper Permian rocks in Kansas.




FIG. 5. Generalized section of middle Permian rocks in Kansas.

PALEOZOIC
PERMIAN
Wolfcampian
Chase

● **Paddock shale member.**—Shale, gray, in northern Kansas containing stringers and vein fillings of calcite; in southern Kansas buff, calcareous, and containing limestone in the lower part. Fossil pelecypods are locally abundant in northern outcrops. The thickness is about 9 feet in Cowley county and ranges between 11 and 13 feet farther north; average 11 feet

● **Krider limestone member.**—Commonly two beds of limestone separated by a bed of shale, each about 1 foot thick. In southern Kansas the separating shale is somewhat thicker. In most exposures the color is yellowish-brown. Characteristic thickness is about 3 feet

■ **Odell shale.**—Shale, gray, yellow, and red chiefly, but with some blue shale in the southern part of Kansas. Along most of the outcrop, red is predominant in the upper and middle parts; gray or yellow is characteristic of the lower part. Thicknesses range from about 20 to 40 feet; average 30 feet

■ **Winfield limestone.**—A thick upper limestone underlain by about 10 feet of fossiliferous gray shale, which is underlain in turn by a thin bed of flinty limestone. The members are not definitely identified in southern Kansas. The combined thickness is about 28 feet

● **Cresswell limestone member.**—This member consists of limestone and locally shale in the upper and middle part, and of massive fossiliferous limestone in the lower part. The upper part, which has been called "Luta limestone," is thinner bedded and more shaly. The shaly middle part commonly contains calcareous concretions and geodes. Echinoid spines and other fossils are plentiful in the lower massive limestone. Throughout a considerable distance the lower massive ledge is almost constantly 3 feet thick. Total thickness of the member is commonly about 17 feet

● **Grant shale member.**—Shale, gray, calcareous and fossiliferous. This is a distinct unit except in southern Kansas, where it probably has been included in the lower part of the Cresswell limestone. Thickness is about 10 feet

● **Stovall limestone member.**—Limestone, gray, dense, with an abundance of steel-gray flint. Fossils are rare. The thickness is commonly about 1 foot

■ **Doyle shale.**—Two shale members and a separating limestone member. The thickness is about 80 feet

● **Gage shale member.**—This part of the Doyle consists mostly of clayey shale, but calcareous shale and a minor amount of limestone occur in the upper part. The lower and middle parts are chiefly varicolored noncalcareous shale consisting of red, green, purple, and chocolate-colored zones interbedded with gray and yellow layers. Locally, a limestone lentil occurs not far from the top and is separated from the overlying Winfield limestone by the more calcareous fossiliferous part of the member. The characteristic thickness is approximately 48 feet

● **Towanda limestone member.**—Limestone, light bluish-gray to yellow, slabby and platy. Fossils are generally absent or rare. About 15 feet has been measured in Geary county but the common thickness ranges from about 5 to 9 feet; average 7 feet

PALEOZOIC

PERMIAN

Wolfcampian

Chase

● *Holmesville shale member.*—Shale, gray, yellow, red, green, unfossiliferous. The thickness ranges from about 20 to 30 feet; average 25 feet

■ *Barneston limestone.*—This formation comprises two limestone members separated by a thin shale member. The upper limestone makes an extensive dip slope and crops out as a steep escarpment that extends from north to south across Kansas. The west portion of the Flint Hills is capped chiefly by the Barneston limestone. The thickness of the formation ranges from about 80 to 90 feet; average 84 feet

● *Fort Riley limestone member.*—Limestone, light gray and tan, massive and thin-bedded, and a minor amount of gray shale. This limestone is naturally divided into four parts, the thicknesses of which vary from place to place. In the basal part there are thin, more or less shaly beds that are overlain by a massive "rim rock," which is a conspicuous outcrop maker. Thin shaly beds and locally clay shale deposits occur in the middle part. The upper strata are rather massive but less so than the "rim rock." The youngest fusulines known in Kansas occur in the Fort Riley limestone. The thickness ranges from about 30 to 45 feet; average 40 feet

● *Oketo shale member.*—Shale, gray, calcareous. Generally absent in southern Kansas and locally absent elsewhere. The thickness, according to measured sections, ranges from a featheredge to 17 feet, but where present this shale parting between flinty limestone below and nonflinty limestone above is generally less than 6 feet; average 5 feet

● *Florence limestone member.*—Limestone, an abundance of flint, and a minor amount of shale. The limestone is commonly lighter in color than the included nodules and layers of steel-gray flint. Shale partings are rather common locally. Fossils, including fusulines, bryozoans, corals, pelecypods, and brachiopods, are somewhat sparse. In most places along the outcrop the thickness is between 35 and 45 feet; average 40 feet

■ *Matfield shale.*—Two units of varicolored shale separated by a limestone member. The thickness ranges from about 60 to 90 feet; average 78 feet

● *Blue Springs shale member.*—Shale, chiefly red and gray, and a relatively minor amount of limestone except in southern Kansas where several limestone beds occur in the upper part of the member. Farther north the member is less calcareous and limestone is absent. Fossils occur in thin limestones and in gray shale beds in the southern part of the outcrop area. The thickness is about 40 feet

● *Kinney limestone member.*—Limestone and shale. Locally there is a single bed of gray, massive or thin-bedded limestone but elsewhere two or more limestone beds are separated by shale. The limestone is sparsely fossiliferous and the associated shale is commonly abundantly fossiliferous. The thickness ranges from a few feet to about 20 feet; average 8 feet

● *Wymore shale member.*—Shale, chiefly gray and yellow but including red, green, and purple bands. Locally most of the unit is composed of bright-colored clay shale. Limestone beds and fossilifer-

ous shale beds are included in the lower part in the southern part of the state. The thickness is about 30 feet

■ **Wreford limestone.**—Two limestones and a shale member. The limestone members are characterized by an abundance of flint in some of the beds. The thickness ranges between about 30 and 40 feet; average 35 feet

● **Schroyer limestone member.**—Limestone, light gray to nearly white, mostly flint-bearing, but commonly containing a nonflinty bed (about 3 feet thick) in the upper part. The thickness is about 18 feet

● **Havensville shale member.**—Shale, gray and calcareous, and thin limestone beds. This unit cannot be easily identified in parts of southern Kansas where the Wreford limestone comprises a continuous limestone section. Thickness in northern Kansas is about 10 feet; average 7 feet

● **Threemile limestone member.**—Limestone, light gray to nearly white, flinty in part, but containing more massive and more resistant nonflinty beds. The thickness is about 10 feet

Council Grove group.—This division of the Permian rocks comprises about 320 feet of limestone and shale. Throughout most of the outcrop area little or no flint occurs in these rocks. In general, they include less massive and thinner limestone units than occur in the overlying group. The group includes strata that, according to older classification, were divided (in descending order) into "Garrison shale," Cottonwood limestone, Eskridge shale, Neva limestone, "Elmdale shale," and Americus limestone.

■ **Speiser shale.**—Shale and limestone, formerly classed in the upper Garrison shale, make up the Speiser shale. The upper part consists of gray fossiliferous shale underlain by a remarkably persistent limestone bed, which is commonly less than 1 foot thick and occurs about 3 feet below the Threemile limestone. The remainder of the formation consists of beds of varicolored shale, red shale being predominant. The thickness of the formation is about 18 feet in northern Kansas and about 35 feet in the southern part of the state where a lenticular bed of sandstone occurs in the middle part; average 25 feet

■ **Funston limestone.**—Light-gray to bluish-gray limestone and gray to yellow shale are contained in the Funston limestone. The limestone occurs in rather massive, sparsely fossiliferous layers, and commonly the beds are separated by shale. In southern Kansas, bluish to nearly black shale occurs in the lower part, and locally the lower limestone beds contain flint. The Funston limestone, Blue Rapids shale, and Crouse limestone were formerly classified as comprising the Bigelow formation. The thickness of the Funston ranges from about 5 to 11 feet; average 8 feet

■ **Blue Rapids shale.**—Shale, gray, containing local limestones. The thickness ranges from about 16 feet in southern Kansas to 25 feet in northern Kansas; average 20 feet

■ **Crouse limestone.**—Limestone and shale. Generally the formation comprises an upper and lower limestone separated by a few feet of shale. The upper part generally displays platy structure and weathers brown. In gen-

Chase

Wolfcampian

Council Grove

PALEOZOIC
PERMIAN

PALEOZOIC
— PERMIAN
— Wolfcampian
— Council Grove

eral the limestone beds are light gray and locally are flinty. The thickness ranges from about 10 to 13 feet; average 12 feet

■ **Easy Creek shale.**—Shale, red and gray, partly calcareous and containing local limestone beds. The upper part is light-colored and calcareous; the lower part is largely red shale. In Marshall county a bed of gypsum occurs near the base. The thickness is approximately 15 feet

■ **Bader limestone.**—Two limestones and a shale member. The thickness range is between approximately 18 and 25 feet; typically about 23 feet

● *Middleburg limestone member.*—Limestone and shale, gray, sparsely fossiliferous. Generally there are two limestone beds separated by shale. Productids and *Composita* are somewhat plentiful in the lower limestone. The thickness ranges from about 4 feet near the Nebraska line to about 8 feet in southern Kansas; average 6 feet

● *Hooser shale member.*—Shale and impure limestones. The shale is gray and fossiliferous, pelecypods predominating. The thickness along the outcrop line across the state is remarkably constant, about 10 feet

● *Eiss limestone member.*—Limestone and shale. Two beds of limestone separated by shale are remarkably persistent across Kansas from Marshall and Nemaha counties to Cowley county. The upper limestone bed, 2 to 3 feet thick, is siliceous and locally contains some flint. The middle part (2 to 7½ feet locally) consists of gray fossiliferous shale in which ramose bryozoans, chonetid brachiopods, and the brachiopod *Meekella* are plentiful. The lower limestone, 1½ to 2½ feet thick, is shaly, thin-bedded, and fossiliferous. The massive upper part of the Eiss limestone is a prominent bench maker in the eastern part of the Flint Hills. The characteristic thickness of the member is about 7 feet

■ **Stearns shale.**—This shale contains a minor amount of impure limestone. The color is mostly gray to olive but red shale occurs in the middle and lower parts. In southern Kansas the upper part contains fossil pelecypods and chonetid brachiopods. Thickness is 8 to 10 feet in southern Kansas and about 17 to 20 feet in the northern part of the state; average 14 feet

■ **Beattie limestone.**—Two limestones and a shale member. The thickness ranges from about 15 to 20 feet; average 18 feet

● *Morrill limestone member.*—This unit consists of brown to gray impure cellular limestone that in the northern part of the outcrop area contains a thin shale parting. In southern Kansas the Morrill thickens, mainly as a result of algal accumulations in the upper part. The thickness ranges from about 2 to about 8 feet; average 5 feet

● *Florena shale member.*—A highly fossiliferous gray shale, in southern Kansas containing thin limestone beds, is called Florena. It is the lowermost part of strata formerly classed under the name Garrison formation. The brachiopod *Chonetes granulifer* is abundant in this shale throughout its line of outcrop. In southern Kansas the variety of fossils is greater than in the north; the fauna contains numerous species of pelecypods and brachiopods and well-preserved specimens of a small trilobite are locally common. The thickness ranges from about 5 to 10 feet; average 7 feet

PALEOZOIC
PERMIAN
Wolfcampian
Council Grove

● **Cottonwood limestone member.**—Limestone, massive, light buff, weathering nearly white, containing abundant slender fusulines in the upper part. Except in the southern part of the state, this is the most prominent limestone of the Council Grove group. In southern Kansas it is thin-bedded and shaly. Throughout most of its line of outcrop it is marked by a fringe of shrubs along the grass-covered Flint Hills slopes. Except for thinning toward the south, the thickness of the Cottonwood limestone is remarkably constant, amounting to about 6 feet

■ **Esckridge shale.**—Shale and a minor amount of impure limestone. The upper and middle parts are more calcareous and fossiliferous. Pelecypods and ostracodes are fairly abundant in thin limestone beds. The lower part is varicolored, consisting of red, chocolate-colored, and green clay shale interbedded with gray shale. The thickness is constantly about 37 feet

■ **Grenola limestone.**—As defined by the Kansas Geological Survey, this formation comprises two limestones and a shale member. The Nebraska Geological Survey includes in the Grenola a few feet of strata that are here classified as belonging in the upper part of the Roca shale. The thickness of the Grenola formation is about 38 feet

● **Neva limestone member.**—This member consists mainly of light-gray massive to flaggy limestone, but includes some gray fossiliferous shale. In general, weathered outcrops of the limestone are brown and pitted, and they display marly facies and breccias of limestone fragments in a calcareous matrix. Fossils include fusulines, corals, and brachiopods. The shale partings are rather non-persistent. The common thickness is about 20 feet

● **Salem Point shale member.**—Shale, gray, calcareous, without fossils or very sparingly fossiliferous. The thickness is about 8 feet

● **Burr limestone member.**—Limestone, light gray, and shale, gray to olive and locally nearly black in the lower part. Ostracodes are plentiful in upper layers and mollusks are rather abundant in the limestone beds. The thickness ranges from about 8 to 14 feet; average 10 feet

■ **Roca shale.**—Shale and limestone; chiefly gray and olive clay shale but containing a large amount of red and locally some green shale. Subdivisions called *Legion shale* and *Sallyards limestone* at the top of the Roca are included in the Grenola formation by the Nebraska Geological Survey. The Sallyards limestone is a thin rather persistent bed but is not present in all exposures. The thickness of the Roca shale is about 20 feet

■ **Red Eagle limestone.**—Two limestones and a shale member. In southern Kansas, the Red Eagle limestone is a single ledge about 20 feet thick, but the two limestone members that are differentiated in central and northern Kansas can be identified also in southern Kansas. The thickness across the state is rather constant, ranging from about 18 to 20 feet; average 19 feet

● **Howe limestone member.**—Limestone, gray to brown, rather impure. Thickness in northern Kansas is commonly between 2 and 4 feet; average 3 feet

PALEOZOIC

PERMIAN

Wolfcampian

Council Grove

● **Bennett shale member.**—Shale and locally some impure limestone. The shale is characteristically dark gray to black but locally the upper part is light gray or green. At some exposures, white fossil brachiopods are abundant in the black shale, forming coquinas composed largely of specimens of *Crurithyris* sp. The Bennett shale is absent in southern Kansas where the two bounding limestones are in contact. The thickness ranges from a featheredge to about 14 feet; average approximately 7 feet

● **Glenrock limestone member.**—Limestone, gray and brown, impure in the northern part of the outcrop area, purer limestone in the southern part. In southern Kansas this member is believed to comprise the greater part of the conspicuous cliff-making Red Eagle limestone. In northern Kansas it is variable in lithology, changing from dense gray limestone to a mass of loosely cemented fusulines or to thin-bedded, slabby limestone. The thickness in the northern part of the state ranges from 1 to 2 feet but in southern Kansas it is about 17 feet; average 9 feet

■ **Johnson shale.**—Shale, gray, locally in northern Kansas containing thin beds of argillaceous limestone. Dark carbonaceous material occurs in the upper part of the shale in the north. The lower and middle parts are commonly somewhat sandy. The thickness ranges from about 16 feet in the northern part of the state to about 25 feet in the southern part; average 20 feet

■ **Foraker limestone.**—Two limestones and a shale member. The thickness is about 50 feet

● **Long Creek limestone member.**—Alternating beds of yellow limestone and shale, or thin-bedded limestone, locally containing an abundance of fusulines in the northern part of the outcrop area. In southern Kansas light-gray limestone, more or less massive in the upper and lower parts, sparsely fossiliferous. Thickness is 5 to 8 feet; average 7 feet

● **Hughes Creek shale member.**—In northern Kansas this part of the Foraker comprises light-gray to nearly black shale and thin limestone beds containing abundant fusulines and, in the lower part, a brachiopod fauna. In southern Kansas the member is a nearly continuous limestone section, massive in the lower part and containing much light-blue flint. Fusulines are plentiful in the flint and limestone. Thickness is about 40 feet

● **Americus limestone member.**—Limestone, commonly in two beds separated by a thin shale, the upper bed locally containing flint nodules. The limestone is commonly bluish-gray and the shale is dark gray. Fusulines are abundant; pelecypods and brachiopods are plentiful. The thickness is about 4 feet throughout most of the line of outcrop but is somewhat less in the southern part of Kansas; average 3 feet

Admire group.—This division, formerly classed as Upper Pennsylvanian, comprises the lowermost 125 to 225 feet of the rocks in Kansas that now are con-

sidered assignable to the Permian system. The strata are chiefly clastic deposits, but they contain thin beds of limestone and some coal. Shale is the predominant rock type.

■ **Hamlin shale.**—Two shales and a limestone member. Thickness about 50 feet

● **Oaks shale member.**—Shale, mostly gray. Crystals of pink celestite occur as vein fillings in Brown county. The thickness ranges from about 10 feet in southern Kansas to perhaps as much as 20 feet in the northern part of the state; average 12 feet

● **Houchen Creek limestone member.**—Limestone and shale, gray and porous, an algal deposit in part. This limestone zone seems to be lenticular, but is recognized north of Kansas river and has been identified in Chase and Elk counties. Measured thicknesses range from 1 foot in northern Kansas to 4 feet in the southern part of the state; average 2 feet

● **Stine shale member.**—Shale, sandstone, and thin limestone beds. In northern Kansas the upper part is sandy shale, the middle part contains limestone beds, and the lower part contains some red shale below gray beds. In southern Kansas it is chiefly gray shale containing some limestone in the middle part. The thickness ranges from about 30 to 40 feet; average 36 feet

■ **Five Point limestone.**—Limestone, light to brownish-gray, slabby to nodular, containing fusulines, bryozoans, brachiopods, and algal remains. Thickness is 1 to 5 feet; average 3 feet

■ **West Branch shale.**—Shale, gray, green, and red, upper part sandy in southern Kansas. Limestone lenses occur in the West Branch shale in northern Kansas. Thickness ranges from about 10 feet in southern Kansas to about 30 feet in the northern part of the state; average 20 feet

■ **Falls City limestone.**—Two thin limestone beds separated by about 8 feet of fossiliferous shale in northern Kansas; in central and southern Kansas a single limestone bed, or locally absent. The fossils include algal remains, fusulines, bryozoans, mollusks, and brachiopods. Locally pelecypods are abundant in the upper part. This limestone is classed as the upper member of the Chicago Mound formation by the Nebraska Geological Survey. The thickness ranges from about 2½ to 10 feet; average 7 feet

■ **Hawxby shale.**—Shale deposits containing thin, lenticular, mollusk-bearing limestones and, more locally, coal beds; sandy in the middle and lower parts. Thickness ranges from about 12 feet in northern Kansas to about 40 feet in the southern part of the state; average 30 feet

■ **Aspinwall limestone.**—Limestone or two thin limestones separated by shale. Unfossiliferous limestone, 1 to 3 feet thick, in the northern part of the outcrop area, but in southern Kansas, where it comprises two limestone beds (3 feet) and as much as 5 feet of shale, the lower limestone bears thick-shelled pelecypods and the upper limestone contains pelecypods and brachiopods. It is classed as the lower member of the Chicago Mound formation by the Nebraska Geological Survey. Average thickness 5 feet

PALEOZOIC

PERMIAN

Wolfcampian

Admire

- PERMIAN —
Wolfcampian —
Admire —
- **Towle shale.**—Clayey and sandy shale, siltstone, and sandstone. The upper part is chiefly gray slightly sandy clay shale, but in northern Kansas it contains some red and green clay. The *Indian Cave sandstone* occurs in the lower part and is only locally present, filling channels as deep as 120 feet. The thickness of the Towle formation ranges from about 15 to 135 feet; average 30 feet
- **Indian Cave sandstone member.**—Sandstone, siltstone, and locally conglomerate in the lower part. Local channel fillings that grade upward into the basal part of the unnamed upper member of the Towle shale. Thickness ranges from a featheredge to about 120 feet

This seemingly important, although obscure, unconformity is marked mainly by large channel sand fillings. It is traceable from Nebraska into Oklahoma. Erosion channels below the Indian Cave sandstone bring deposits classified as lowermost Wolfcampian into contact with rocks ranging downward from the Brownville limestone, or possibly slightly younger beds, to the Dover formation or lower. This evidence of widespread erosion, accompanied by rather noteworthy change in the lithologic characters of limestones, and paleontologic change, which is accentuated by the disappearance of old species and the appearance of some new genera and species, warrant the placement of the Wolfcampian-Virgilian boundary at this position.

Unconformity

PENNSYLVANIAN SYSTEM.—This division of Late Paleozoic rocks is widely represented throughout the world, being distinguished by importance of its coal deposits and by characters of its large assemblage of marine invertebrates and varied land plants. It is one of the most important among outcropping strata of Kansas, both on account of the economic value of its contained materials and the scientific value of its rock succession as a standard of reference in studies of equivalent-aged deposits in other parts of the continent. The Pennsylvanian rocks comprise the upper part of a large division that has long been known as the Carboniferous system. The term Carboniferous is almost universally used by geologists outside of North America and has been employed by the State Geological Survey of Kansas in designating outcrops shown on the large geological map of the State (1937) and described in numerous reports. Pennsylvanian rocks have been classified in these publications as a subsystem. Here, we recognize the validity of considerations that have led a great majority of American geologists to treat the Pennsylvanian as an independent geologic system. The aggregate thickness of exposed formations belonging to this system in Kansas is about 3,100 feet. Outcrops occur throughout the eastern one fourth of the State.

VIRGILIAN SERIES.—The Virgilian series comprises the youngest Pennsylvanian rocks of the midcontinent region. They are separated from rocks above and below by unconformities. In Kansas this series is divided into three groups on the basis of general differences in lithology and the nature of cyclic deposits. The thickness is commonly about 1,200 feet.

Wabaunsee group.—The uppermost major division of Virgilian strata includes beds below the unconformity that separates Permian and Pennsylvanian rocks




FIG. 6. Generalized section of lower Permian rocks in Kansas.

and above the top of the Topeka limestone. Limestones in this group are uniformly thin but persistent. Sandstone is plentiful, but shale is the chief rock. The thickness of the Wabaunsee group is about 500 feet, excepting in places where the overlying unconformity cuts out the upper beds.

■ **Brownville limestone.**—Bluish-gray limestone that weathers yellowish or brown, occurring in one bed or in two beds separated by a thin layer of shale. Varying texture from impure sandy appearing rock to fairly pure dense limestone. Characteristic fossils include large fusulines, brachiopods *Meekopora* and *Chonetes granulifer*, crinoid fragments, and bryozoans. The thickness ranges from about 2 to 8 feet; average 5 feet

■ **Pony Creek shale.**—Bluish-gray shale chiefly, some red clay shale or sandy shale locally, and sandstone in the middle part in some places. A thin coal bed occurs in the upper middle part in southern Kansas. Brachiopods and bryozoans more or less common in upper layers. The Pony Creek shale is classified as the upper part of the Woods Siding formation in Nebraska. Thickness ranges from about 5 to 20 feet; average 14 feet

■ **Caneyville limestone.**—Three limestones and separating shales in the southern part of the outcrop area; two limestones and a shale elsewhere. In northern Kansas, where no discernible boundary between Caneyville and Pony Creek is found, strata from the base of the Nebraska City limestone to the base of the Brownville limestone are designated Caneyville-Pony Creek, or they may be called the Woods Siding formation, which is defined by the Nebraska Geological Survey to have limits that correspond exactly to Caneyville-Pony Creek. Thickness about 21 feet

● **Grayhorse limestone member.**—Gray, medium- to coarse-grained limestone, fragmental or coquinoid, ferruginous, broken surfaces showing curved cleavage surfaces of iron- or magnesium-bearing carbonate crystals. Large specimens of *Myalina* are rather common. This member is well developed in southern Kansas and northern Oklahoma but is not known to extend north of Greenwood county. The thickness ranges from about 0.5 to 6 feet; average 1 foot

O **Shale** ranging in thickness from about 5 to 15 feet lies between the Grayhorse limestone and the limestone described below; average 10 feet

O **Limestone**, bluish-gray, slightly arenaceous, massive, fusuline-bearing, occurs in the middle part of the Caneyville in southern Kansas. Thickness is commonly about 1 foot

O **Shale**, gray, sandy, 5 to 10 feet thick, occurs in southern Kansas between the fusuline-bearing limestone of the Caneyville limestone and the Nebraska City limestone. Average thickness 7 feet

● **Nebraska City limestone member.**—Bluish to greenish-gray, impure, soft limestone that weathers light yellowish-brown; contains bryozoans, brachiopods, and sparse mollusks. This member, which is well developed in northern Kansas, is traced to Oklahoma. It is classed as the lower unit of the Woods Siding formation by the Nebraska Geological Survey. Thickness ranges from about 1 foot to 5 feet; average 2 feet

PALEOZOIC
PENNSYLVANIAN

Virgilian

Wabaunsee

■ **French Creek shale.**—Bluish-gray to yellowish-brown clayey and sandy shale commonly containing light-brown or tan sandstone and one or more coal beds in the upper part. The average thickness is about 30 feet

■ **Jim Creek limestone.**—A persistent bluish-gray, fine-grained, hard limestone, containing in many places slender fusulines and, more locally, a large variety of other marine fossils. It is traced across Kansas. The thickness ranges from a few inches to nearly 2 feet; average 1 foot

■ **Friedrich shale.**—Bluish-gray shale that weathers yellowish and brown. Sandstone is more or less common and in Greenwood county a thin coal bed occurs in the upper part. Pelecypods and sparse bryozoans and brachiopods are found near the top. Average thickness is about 15 feet

■ **Grandhaven limestone.**—Commonly two limestones separated by a few feet of shale. The upper limestone is light-gray and weathers almost white, algal deposits forming a prominent part of the rock. The shale is mostly bluish-gray and locally calcareous. The lower limestone is gray to bluish-gray and weathers brown. In exposures along Cottonwood river, it is cross-bedded and carries abundant brachiopods and bryozoans and some mollusks. Large fusulines are present at some outcrops and algae ("cryptozoans") are common. The formation is traced from Shawnee county southward to Oklahoma and may be recognized north of Kansas river. The thickness averages about 10 feet

■ **Dry shale.**—Bluish-gray, generally clayey, but locally somewhat sandy shale containing a thin coal bed near the top in southern Kansas. Marine fossils are abundant locally above the coal bed. The thickness ranges from about 5 to 20 feet; average 15 feet

■ **Dover limestone.**—Light-gray and blue limestone, most of which weathers almost white. Algae ("cryptozoans") and fusulines are characteristic. Southward from Greenwood county, a fine-grained blue limestone carrying a mixed fauna of mollusks and brachiopods occurs a few feet below the more persistent fusuline- and algal-bearing part. In northern Kansas the thickness is 2 to 4 feet but in southern counties the formation thickens to about 15 or 20 feet; average 10 feet

■ **Langdon shale.**—Bluish-gray clayey to sandy shale, containing the persistent thin *Nyman coal* and locally platy to massive sandstone in the upper part. Marine invertebrates are abundant at places in northern Kansas below the upper sandy facies. Langdon is a term recently proposed by the Nebraska Geological Survey. It here replaces the previously used name "Table Creek shale," because study of the Table Creek type locality has shown that this name is unsuitable for beds between the Dover and Maple Hill limestones. Thickness ranges from about 5 to 50 feet; average 30 feet

■ **Maple Hill limestone.**—Gray, medium hard, more or less sandy limestone, weathering reddish-brown; generally bearing crinoid remains, bryozoans, brachiopods, and sparse mollusks. Small slender fusulines are common in the vicinity of Emporia. The Maple Hill limestone is not definitely recognized south of Emporia but is believed to be represented in southern Kansas by thinner bedded limestone. Thickness about 1 to 5 feet; average 3 feet

PALEOZOIC

PENNSYLVANIAN

Virgilian

Wabausee

■ **Wamego shale.**—Bluish-gray clay shale and yellowish-brown micaceous sandy shale; nearly black shale locally in the upper part. Between Kansas and Cottonwood rivers a more or less persistent coal bed occurs in the upper-middle part. A mixed marine fauna is found in beds above the coal. The formation is not differentiated from the Willard shale south of Lyon county. The term Wamego, recently introduced by the Nebraska Geological Survey, has not been used previously by the Kansas Geological Survey; it replaces "Pierson Point shale," which is judged to be an inappropriate name. Thickness of the Wamego ranges from about 6 to 25 feet; average 17 feet

■ **Tarkio limestone.**—This formation, consisting of gray limestone that weathers deep yellowish-brown, is one of the most easily recognized units in the Wabausee group. It is characterized by the brown color of weathered outcrops and the abundance of large fusulines (*Triticites* sp.). Locally algal remains occur in the upper part. The formation is present from a point in northern Lyon county northward across the state, and seemingly occurs at some places in southern Kansas. The thickness ranges from a feather-edge to about 10 feet; average 6 feet

■ **Willard shale.**—Dark bluish-gray and brown shale, and sandstone. Sandstone is prominent in many places in northern Kansas but thin or absent farther south. Inasmuch as the Maple Hill and Tarkio limestones are not recognized in southern Kansas, the shale and sandstone section extending from the base of the Willard to the top of the Langdon shale is designated as the Willard-Langdon shale; its thickness ranges from about 5 to 15 feet. The thickness of the Willard in northern and central Kansas ranges from about 30 to 66 feet, the maximum being near Kansas river; average about 40 feet

■ **Elmont limestone.**—Dark-blue massive limestone, weathering bluish-gray and displaying closely spaced vertical joints; locally conglomeratic in the basal part. Fusulines are plentiful in parts of the limestone along most of the outcrop area; brachiopods and mollusks are present locally in the lower part. The upper part commonly contains algal remains and mollusks. Locally near Topeka, the upper algal-molluscan facies consists of 8 feet of coquinoïd cross-bedded limestone. The Elmont limestone was classed formerly as the upper part of the "Emporia limestone." In northern Oklahoma the Elmont is included in the upper part of the Stonebreaker limestone. The thickness along the Kansas outcrop line ranges from about 1 to 15 feet; average about 5 feet

■ **Harveyville shale.**—Bluish and greenish-brown clayey shale, locally sandy shale and thin platy sandstone. In places a coal bed occurs above the sandstone and marine invertebrate fossils are found above the coal horizon. The formation is continuous across Kansas, having a thickness range from less than 1 to 25 feet; average about 10 feet

■ **Reading limestone.**—The most persistent part consists of dark-blue, dense, hard limestone showing vertical jointing, and weathering light bluish-gray mottled with light brown or yellow. The formation comprises one to three beds of limestone and blue-gray shale beds between the limestone layers. Fusulines are characteristic of the persistent dense blue limestone. Locally an upper algal phase is present as a single distinct limestone above

the more persistent part, and a brachiopod- and pelecypod-bearing limestone may be present below the persistent bed. The Reading constitutes the lower part of the "Emporia limestone" as previously recognized. Thickness of the formation is about 1.5 to 15 feet; average 6 feet

■ **Auburn shale.**—A somewhat complex unit composed chiefly of gray shale, containing minor amounts of sandstone and limestone. Persistent platy sandstone occurs in the upper part in southern Kansas, and locally limestone is prominent in the upper middle part. Near Kansas river, this limestone is a cross-bedded 5-foot layer consisting partly of conglomerate and partly of a coquinoid mass of molluscan and algal remains. Dark shale containing numerous ostracodes and pelecypods occurs just below the cross-bedded limestone. A persistent impure limestone about 0.5 foot thick is found near the middle of the Auburn shale, and at a little lower position there is red shale and coal locally. The lower Auburn generally consists of yellowish-gray sandy shale. The thickness of the formation in Kansas ranges from about 20 to 70 feet; average 50 feet

■ **Wakarusa limestone.**—Dark bluish hard limestone that becomes light brown when weathered. Locally, an algal-molluscan or fine-grained sandy limestone occurs in the upper part and a mollusk- or brachiopod-bearing limestone may be present below the more persistent part. Shale separates the limestones where more than one is present. Fusulines, massive bryozoans, large brachiopods (*Dictyoclostus*), algal remains, and other fossils are characteristic of the persistent ledge. The Wakarusa is traced across Kansas into northern Oklahoma where the fusuline-bearing phase is represented by the so-called "Cryptozoan limestone." The thickness of the formation ranges from about 2 to 18 feet; average 8 feet

■ **Soldier Creek shale.**—Bluish-gray clayey to sandy shale; locally a thin coal bed in the upper part. Marine invertebrates occur in the upper part in places. Thickness ranges from about 12 to 18 feet; average 15 feet

■ **Burlingame limestone.**—Limestone, brown, fine-grained, hard, thick-bedded, appearing mottled and brecciated in some exposures. Shale commonly separates the limestone beds. Fusulines are fairly common in the more persistent part, and algal remains are conspicuous near the top. Sponge-like algal deposits, composed of *Spongiastroma* are numerous in a few exposures in northern Kansas. The thickness of the formation ranges from about 4 to 16 feet; average 9 feet

■ **Silver Lake shale.**—Gray and yellow clay shale, variably associated with platy impure limestone, sandy shale, or shaly sandstone. One or more coal beds occur in the upper part. Marine invertebrates and land plant fossils may be present. This shale comprises the topmost strata of the formerly recognized "Scranton shale" which includes all beds between the Burlingame and Howard limestones. The average thickness of the Silver Lake shale is about 25 feet

■ **Rulo limestone.**—Bluish-gray limestone that locally is mottled with light brown when weathered. Locally a molluscan phase occurs in the lower part and a thin algal zone in the upper part. Brachiopods and bryozoans are more or less common in the more persistent part. The thickness averages about 2 feet

PALEOZOIC

PERMIAN

Virgilian

Wabauense

PALEOZOIC

PENNSYLVANIAN

Virgilian

Wabausee

■ **Cedar Vale shale.**—Bluish to yellowish-brown clayey and sandy shale and sandstone with a persistent coal bed (*Elmo*) near the top. Marine invertebrate fossils occur above the coal. The average thickness is about 25 feet

■ **Happy Hollow limestone.**—Generally a single, very persistent massive bed of pinkish-brown limestone characterized by large fusulines. In some places there is an upper algal bed in contact with the more persistent part or separated from it by a thin shale bed, and locally a dark bluish-gray limestone occurs 1 or 2 feet below the fusuline-bearing bed. Thickness ranges from about 1 to 8 feet; average 4 feet

■ **White Cloud shale.**—Bluish-gray to yellowish-brown clayey and sandy shale and sandstone that locally fills channels and contains conglomerate at the base. Sparse marine and land plant fossils are present. The thickness ranges from about 30 to 80 feet; average 50 feet

■ **Howard limestone.**—Three limestone members and two shale members, of which the middle limestone is the most persistent. The thickness of the formation ranges from about 8 to 30 feet; average 13 feet

● **Utopia limestone member.**—Dark bluish-gray dense limestone that weathers light brownish-gray, resembling a coquina in many outcrops. Fossils include algal remains, bryozoans, brachiopods, and mollusks. The thickness ranges commonly from less than 1 foot to about 4 feet. Consists of two or more limestones and an ostracode-bearing shale in northern Kansas. Average thickness 2 feet

● **Winzeler shale member.**—Bluish-gray to yellowish-gray clay shale. Marine fossils occur in the lower part. Thickness is about 3 to 8 feet; average 4 feet

● **Church limestone member.**—Commonly one massive bed of blue to bluish-gray limestone that weathers to a deep rich brown. Crinoid remains, productid brachiopods, and "cryptozoan" algae are more or less characteristic. A thin zone at the top contains abundant bryozoans, and fusulines occur sparingly in the upper part in southern Kansas. The thickness ranges from about 1.5 to 6 feet; average 2 feet

● **Aarde shale member.**—Bluish-gray to yellowish-gray clayey and sandy shale containing a very persistent coal bed (*Nodaway*), ranging from about 1 inch to 2 feet in thickness, and persistent black fissile shale. A hard dense vertical-jointed limestone, 3 to 10 inches thick, occurs next below the black fissile shale in southern Kansas. Thickness ranges from about 3 to 7 feet; average 4 feet

● **Bachelor Creek limestone member.**—Commonly sparsely fossiliferous, bluish-gray, impure limestone which is present from a point in Greenwood county southward. Also well developed, perhaps locally, in southern Osage county. Thickness is commonly about 1 foot.

■ **Severy shale.**—Yellowish-brown and bluish-gray clay shale and a minor amount of platy to massive sandstone in the upper part. Sparse marine fossils occur in the uppermost part and in shales below some thin limestone beds which occur in southern Kansas. A disconformity in the lower part of the Severy is indicated by various outcrops in northern Kansas. The thickness is commonly about 75 feet

Shawnee group.—The Shawnee group comprises four limestone formations and three shale formations. Thick limestones and a distinctive type of cyclic sedimentation are characteristics that distinguish these rocks from those of neighboring groups. The average thickness is about 325 feet.

■ **Topeka limestone.**—Limestones and shales having a wide range of lithology. The two upper limestone members seem to be absent between Kansas river and Greenwood county, but are recognized in the north and tentatively identified in the south. In Oklahoma the upper part of the Pawhuska formation is correlated with the Topeka limestone. The thickness of the Topeka ranges from a nearly constant measurement of 33 feet in northern Kansas to 55 feet near the Oklahoma boundary. The average thickness is 35 feet

● **Coal Creek limestone member.**—Light bluish-gray limestone and nodular shale or dark-blue more massive limestone that weathers light bluish-gray or brown. The member is not positively identified south of Kansas river, but the "red limestone" in southern Kansas is probably its equivalent. Fusulines are locally abundant, and at many places numerous brachiopods, bryozoans, and other well-preserved invertebrate fossils are found. The thickness of the Coal Creek limestone ranges from 2 to 5 feet; average 3 feet

● **Holt shale member.**—Bluish-gray shale in the upper part and black shale in the lower part. This member is not recognized with certainty south of Topeka because the limestone beds above and below it disappear, at least for many miles, and because the distinctive black slaty beds do not persist southward. Shale beneath the uppermost Topeka limestone in southern Kansas probably represents the Holt member, however. Corneous brachiopods and conodonts occur in the black shale. The thickness ranges from 1.5 to 3 feet; average 2 feet

● **Du Bois limestone member.**—One or more dark-blue or greenish-blue fine-grained, somewhat earthy limestone beds with prominent vertical jointing. Mollusks and brachiopods are common. The thickness commonly ranges from about 0.5 to 2 feet, but the member is not definitely recognized south of Topeka; average 1 foot

● **Turner Creek shale member.**—Bluish or greenish-gray clayey and calcareous shale, bearing few invertebrate fossils. The thickness ranges from 1 to 5 feet in northern Kansas; shale in southern Kansas that is correlated with the Turner Creek measures 12 to 15 feet; average 5 feet

● **Sheldon limestone member.**—A massive light-gray to nearly white, very fine-grained, dense limestone that commonly contains numerous small algal growths of the type known as *Osagia*. The rock weathers light yellowish-gray and makes smooth rounded surfaces at the outcrop. The member is a persistent layer that is identified northward from Kansas river but it is absent to the south as far as Greenwood county; in southern Kansas there is an algal limestone at the Sheldon horizon. Thickness, 0.7 to 2 feet; average 1 foot

● **Jones Point shale member.**—Clayey, calcareous, and silty gray shale, locally containing nodular or platy limestone beds. The mem-

PALEOZOIC
PENNSYLVANIAN
Virgilian
Shawnee

PALEOZOIC

PENNSYLVANIAN

Virgilian

Shawnee

ber contains fairly numerous brachiopods and mollusks at some outcrops but is poor in fossils at many exposures. The thickness in northern Kansas ranges from 1 to 10 feet, but beds in Elk and Chautauqua counties that are correlated with the Jones Point shale are 10 to 15 feet. The average thickness is about 6 feet

● *Curzon limestone member.*—This is a persistent prominent subdivision of the Topeka formation that is readily and positively identifiable throughout northern Kansas. It consists of two or more beds of massive bluish-gray, brown-weathering limestone that is mostly hard and resistant, forming a well-marked escarpment. Nodules of chert are common. Fusulines occur sparingly to very abundantly in the lower and middle parts of the member, together with some brachiopods and other invertebrates; bryozoans and echinoid remains are common in upper layers. This limestone, which erroneously has been called Hartford in some earlier Kansas reports, is undoubtedly the same as that termed the Curzon limestone in publications of the Nebraska Geological Survey and, in spite of some objections to the origin of this term, it is employed here for rocks in Kansas. The thickness of the Curzon ranges from 5 to 12 feet; average about 8 feet

● *Iowa Point shale member.*—Yellowish-gray to bluish-gray, clayey to calcareous shale that locally contains sandstone layers and a thin coal bed. This shale is typically developed and thickest in northern Kansas near the Nebraska boundary; in sections near Topeka it is only 2 feet thick, and farther south it disappears, as indicated by fusuline-bearing lower layers of the Curzon resting directly on the algal upper part of the Hartford limestone. Thickness, a featheredge to 14 feet; average about 4 feet

● *Hartford limestone member.*—Massive, light bluish-gray limestone that weathers brownish. This member commonly bears numerous fusulines except in the upper part, which contains numerous *Osagia* and is of algal origin. The lower beds are characterized by presence of the chambered sponge, *Amblysiphonella*, specimens of which can be found at almost every outcrop from Nebraska to Oklahoma. The upper algal limestone is highly variable in thickness, ranging from almost nothing to 12 feet. This member, named many years ago from outcrops near Hartford in Neosho county, is the same as rocks recently called Wolf River limestone in Nebraska reports. The thickness of the Hartford limestone ranges from 3 to 13 feet; average 5 feet

■ *Calhoun shale.*—Clayey and sandy shale, a minor amount of limestone, and one or more coal beds. In northern Kansas a thin coal bed and much sandstone, a part of which fills channels, occur near the top of the formation. In southern Kansas the shale progressively diminishes in thickness until near the Oklahoma boundary it is locally absent. In northern Kansas, near the Nebraska line, the Calhoun is only 10 feet thick. Maximum thickness of this shale, about 60 feet, is developed near Kansas river; average 30 feet

■ *Deer Creek limestone.*—This persistent, escarpment-making formation comprises three limestone members and two shale members. The outcrop belt crosses Kansas from Doniphan county in the northeast to Chautauqua

county in the south. Lithologic characters are fairly constant. The thickness ranges from about 20 to a local abnormal maximum in Osage county of 80 feet; average 40 feet

● *Ervine Creek limestone member.*—This member is mainly composed of light-gray to nearly white or bluish-gray fine-grained limestone characterized by thin, wavy bedding that locally contains chert nodules. It bears a large assemblage of invertebrate fossils, including fusulines, corals, echinoid and crinoid fragments, bryozoans, brachiopods, and mollusks. The thickness, which ranges from 5 to 32 feet, averages about 14 feet. Above the wavy-bedded limestone is a less persistent limestone, generally massive, that is variable in lithology. Locally it is oölitic; elsewhere it is coquinoïd, nodular, very fine-grained or sandy; the alga called *Osagia* is common. This limestone, which ranges from a featheredge to about 6 feet, is generally separated from the underlying limestone by 1 to 2 feet of yellowish clayey to sandy shale but the shale is absent in many places. Average thickness of the Ervine Creek limestone member is 18 feet

● *Larsh-Burroak shale member.*—Gray or yellow clayey shale in the upper part and black, more or less fissile shale in the lower part. Fossils are rare in the upper part but conodonts occur in the black portion. Thickness ranges from 2 to 8 feet; average 3.5 feet

● *Rock Bluff limestone member.*—This is a distinctive, massive, vertical-jointed bed of dark-blue limestone that develops a bluish-gray or creamy surface film on weathering. Fusulines are common; a few brachiopods and other marine invertebrates are present. The thickness ranges from about 1 to 2.5 feet; average 1.5 feet

● *Oskaloosa shale member.*—Bluish or yellowish-gray shale in northern Kansas and sandy micaceous shale containing a red zone and some nodular limestone in the southern part of the state. The thickness ranges from an observed minimum of 3 feet to a maximum of 50 feet locally in Osage county; average about 12 feet

● *Ozawkie limestone member.*—Brownish-gray, brown-weathering, massive limestone. Fossils are somewhat sparse, but fusulines and other marine fossils are abundant in some outcrops. The thickness ranges from 1 to 22 feet (maximum in Osage county); average 5 feet

■ *Tecumseh shale.*—Chiefly clayey and sandy shale, locally having a more or less discontinuous limestone (*Ost*) in the upper part. This limestone is not persistent enough to call for subdivision of the formation as in Nebraska. Sandstone is locally present not far below the Deer Creek limestone. The thickness of the Tecumseh ranges from an observed minimum of 12 feet in southern Kansas to a maximum of 65 feet near Kansas river; average 35 feet

■ *Lecompton limestone.*—Four limestone members and three shale members, all of which can be traced across Kansas, make up this formation. The thickness ranges from about 30 to about 50 feet; average 34 feet

● *Avoca limestone member.*—Dark bluish-gray somewhat earthy limestone occurring in one or two beds. Large fusulines are the most common fossils. In southern Kansas a "cryptozoan"-bearing limestone is a characteristic element of the member. Algal-molluscan

PALEOZOIC
PENNSYLVANIAN
Virgilian
Shawnee

limestone beds occur at the top of the Avoca in some outcrops. Locally the thickness ranges to 15 feet but the average is about 4 feet

● *King Hill shale member*.—Bluish-green to reddish-gray shale, calcareous in northern Kansas. Brachiopods occur in the upper part. The thickness range is from 4 to 24 feet. Average 8 feet

● *Beil limestone member*.—Alternating beds of hard bluish-gray limestone, weathering yellowish, and calcareous shale, or light-gray, nearly white, wavy-bedded limestone in southern Kansas. Algal limestone resembling an oölite comprises the upper part in northern Kansas. Corals and fusulines are abundant and characteristic. The thickness ranges from 4 to 15 feet; average 7 feet

● *Queen Hill shale member*.—Bluish-gray or yellow shale in the upper part; hard black fissile shale, bearing conodonts, in the lower part. The thickness ranges from about 3 to 6 feet; average 4 feet

● *Big Springs limestone member*.—Dark bluish-gray dense limestone, commonly occurring as a single bed with prominent vertical jointing. Weathering produces a yellowish-brown film. Fusulines are extremely abundant. The thickness ranges from 1 to 5 feet; average 2 feet

● *Doniphan shale member*.—Chiefly bluish and yellowish-brown clay shale and in the southern part of the state some red shale. Sandy shale, an underclay-like deposit, and coaly zones occur locally in Shawnee county. The thickness ranges from about 5 to 10 feet; average 7 feet

● *Spring Branch limestone member*.—Gray, somewhat sandy limestone that weathers deep brown and occurs in massive slightly uneven beds. Fusulines are abundant in most outcrops. In northern Kansas the thickness is commonly 5 feet; in southern Kansas this member is a very sandy impure limestone 2 to 3 feet thick; average 4 feet

■ *Kanwaka shale*.—This formation is divisible in northern Kansas into two shale members and a limestone member. It consists chiefly of gray, yellow, and brown clayey to sandy shale containing much sandstone, one or more thin coal beds, and in southern Kansas red shale. The thickness ranges from about 40 feet in northern Kansas to 150 feet in Chautauqua county; average 80 feet

● *Stull shale member*.—Yellowish-brown sandy shale, sandstone, and locally a thin coal bed. Sandstone, partly of channel-filling type, occurs in the upper part. Sandstone deposits thicken southward and in southern Kansas are represented by the *Elgin sandstone*. Land plant fossils are more or less abundant in the sandy facies. The thickness of the member ranges from 22 feet to at least 45 feet; average 35 feet

● *Clay Creek limestone member*.—Dark-blue to bluish-gray dense or granular limestone showing distinct vertical jointing. Persistent northward from Osage county, and probably represented in southern Kansas by calcareous fossiliferous sandstone and shale. Fusulines and other marine invertebrates are present. Thickness ranges from a featheredge to 5 feet or slightly more; average 2 feet

● *Jackson Park shale member*.—Bluish-gray and yellowish-brown sandy shale and sandstone. Land plant fossils are more or less abundant. This and higher subdivisions of the Kanwaka here named are not differentiated in southern Kansas. The observed thickness ranges from 16 to 52 feet; average 43 feet

■ *Oread limestone*.—The basal formation of the Shawnee group comprises four limestone members and three shale members, the outcrop of which is marked by a prominent escarpment across the state from Doniphan county in the north to Chautauqua county in the south. The average thickness in northern and central Kansas is 52 feet; in southern Kansas the thickness increases to about 100 feet, owing to expansion of the interval between the lower two limestone members; average 70 feet

● *Kereford limestone member*.—This member is variable in lithology, commonly more or less oölitic or fine-grained and flaggy. In northern Kansas, upper slabby cross-bedded layers, containing algal and locally land plant remains, overlie more massive rock that contains fusulines and other marine invertebrates. At some outcrops the member consists of a single bed of dark bluish-gray dense limestone; elsewhere, especially in Osage county, it is made up of dense bluish flaggy limestone partly interbedded with shale. The Kereford is not recognized in southern Kansas. Thickness ranges from a feather-edge to more than 40 feet; average about 6 feet

● *Heumader shale member*.—Commonly dark-gray shale, locally bluish or greenish, clayey to silty, and in southern Kansas sandy. Well-preserved mollusks and other fossils occur commonly. Thickness in northern Kansas ranges from 2 to 5 feet; average 4 feet

● *Plattsmouth limestone member*.—The Plattsmouth is the main limestone member of the Oread formation. Commonly it is light bluish-gray, light-weathering, dense limestone that occurs in wavy beds. Flint is more or less abundant in the northern part of the outcrop. Fossils include fusulines, corals, bryozoans, brachiopods, sparse mollusks, and other invertebrates. Algal remains occur locally at the top. The thickness ranges from 2 feet in southern Greenwood county to 23 feet in Atchison county; in southern Kansas the member is commonly 15 to 20 feet thick; average 16 feet

● *Heebner shale member*.—In many outcrops this member consists of four distinct shale units which are, in descending order: calcareous clay shale, dark bluish-gray shale, black platy shale, and a thin bed of gray or yellow clay shale. The black platy shale is almost everywhere the thickest part, but is commonly less than 3 feet. Small brachiopods occur in the upper part, conodonts are found in the black shale, and locally there are numerous gastropods in the thin film of clay at the base of the member. The thickness of the Heebner shale across Kansas is very constant, ranging from 5 to 7 feet; average 6 feet

● *Leavenworth limestone member*.—A dark bluish-gray, dense, massive limestone occurring in a single bed having prominent vertical joints. Fusulines and brachiopods are the most common fossils. Almost everywhere across Kansas the thickness is more than 1 and less than 2 feet; average 1.5 feet

PALEOZOIC
PENNSYLVANIAN
Virgilian
Shawnee

Shawnee

● *Snyderville shale member*.—Gray and bluish-gray clay and in part red shale. In northern Kansas it is largely a structureless under-clay-like deposit; in southern Kansas it is partly red and sandy and locally contains sandstone and impure nodular limestone. Marine fossils, especially *Chonetes*, occur in the uppermost part, and in southern Kansas there are other fossiliferous zones. The thickness in the northern part of the state averages about 12 feet, but it increases to about 75 feet in southern Kansas; average 30 feet

● *Toronto limestone member*.—Brownish-gray massive limestone that weathers deep brown. Locally flinty or algal limestone occurs in the upper part. The Toronto is typically developed from Woodson county northward but is thinner and sandy or locally absent in southern Kansas. The member is absent also in part of southern Douglas county where a limestone conglomerate occupies its approximate stratigraphic position. Fusulines, corals, and small brachiopods are the most common fossils. The thickness of the Toronto in northern Kansas ranges from 8 to 12 feet; in southern Kansas it is generally less than 4 feet; average 6 feet

Douglas group.—The Douglas group underlies the Shawnee group conformably, except possibly in southern Douglas county. It comprises chiefly clastic rocks, shale, and sandstone. Limestone, coal, and conglomerates are quantitatively of minor importance. The thickness is approximately 250 feet.

■ **Lawrence shale.**—Blue-gray and yellowish shale, tan-colored sandstone, and a minor amount of coal, limestone, and conglomerate comprise the Lawrence shale. The formation extends from the base of the Toronto limestone to the disconformity below the Ireland sandstone or, in places where the disconformity is undeterminable, to the top of the Haskell limestone. Except for members known as the Amazonia limestone and Ireland sandstone, which are somewhat impersistent, the Lawrence shale is not subdivided. The thickness ranges from about 40 to 175 feet; average 100 feet

● *Amazonia limestone member*.—This is typically a light-gray, dense, hard limestone occurring about 25 or 30 feet below the top of the Lawrence shale. Locally it is partly coquinoïd or brecciated in appearance. Outcrops are found in Doniphan, Atchison, Franklin, Coffey, and Woodson counties and possibly farther south. In Franklin county and vicinity, the Amazonia occurs a few feet below the *Williamsburg coal*, which is mined at several places. Sponges are locally common in the member (Atchison county) and fusulines occur in limestone that may be Amazonia in southern Kansas. Fossils are not abundant generally. Thickness ranges from a featheredge to about 13 feet. In the southern outcrop area it averages about 1.5 feet

● *Ireland sandstone member*.—Light-buff or tan thin-bedded and massive sandstone, in part cross-bedded. Where thickest, the sandstone lies above a disconformity, which is identified in many places but seemingly is not continuous. In some places, especially where the Ireland cuts out the Haskell limestone and still lower beds, a limestone conglomerate is found at the base of the sandstone. The upper

PALEOZOIC

PENNSYLVANIAN

Virgilian

Douglas

boundary is somewhat indefinite because much sandy shale and thin-bedded sandstone occur in the upper part of the Lawrence shale. Local coal beds are associated with these sandy sediments. In Woodson county the top of the Ireland sandstone is only a few feet below the Amazonia limestone. The thickness of the Ireland sandstone ranges from a featheredge to about 100 feet.

Disconformity

■ **Stranger formation.**—Light-tan to yellowish-gray sandstone and shale and a minor amount of limestone, coal, and conglomerate. Thickness ranges from 40 to 220 feet; average 155 feet

● **Robbins shale member.**—Gray and yellowish-gray marine shale, locally cut out and overlapped by the Ireland sandstone in Franklin county and the southern part of Douglas county. This member is essentially restricted to areas southward from Woodson county. Phosphatic concretions occur in a zone at the base in the northern part of the outcrop area. Thickness ranges from a featheredge to 110 feet; average about 70 feet

● **Haskell limestone member.**—Bluish-gray fine-grained limestone, locally having oölitic layers at the top and base. Pelecypods and gastropods occur in the oölitic facies. The main part bears algal remains, fusulines, and brachiopods. In parts of Douglas and Franklin counties, the Haskell limestone is cut out and overlapped by the Ireland sandstone. The thickness, which ranges from a featheredge to about 10 feet, averages 3 feet

● **Vinland shale member.**—Gray, clayey, calcareous, and sandy shale and locally some sandstone underlying the Haskell is named the Vinland shale. Locally in Woodson county, a dark-green layer occurs in the middle part. North of Anderson and Coffey counties, where the Westphalia limestone is absent, the top of the upper *Sibley coal* is regarded as marking the base of the Vinland shale. A disconformity beneath a sandstone at the base of the Vinland cuts out the Westphalia limestone locally in Coffey county. The thickness of the Vinland member ranges from about 9 to 50 feet; average 20 feet

Local unconformity

● **Westphalia limestone member.**—A limestone characterized by abundant fusulines; not definitely recognized north of T. 19 S., but fairly persistent in southern Kansas. A limestone slightly above the *Sibley coal* in southern Douglas county possibly represents this limestone. The thickness of the member ranges from a featheredge to about 5 feet; average 2 feet

● **Tonganoxie sandstone member.**—Massive cross-bedded sandstone and sandy shale, containing several discontinuous coal beds, form the lower part of the Stranger formation in most places. The upper and lower *Sibley coals*, in the northern part of the outcrop area, are the most important coal beds of the unit. The top of the Tonganoxie member is rather indefinite, but in general the base of the Westphalia limestone or the top of the upper *Sibley coal* bed is defined as the boundary. Locally limestone conglomerate occurs at the base. Like the Ireland sandstone, the Tonganoxie is more massive and is thicker

PALEOZOIC
PENNSYLVANIAN
Virgilian
Douglas

Virgilian
Douglas

in southern Kansas, where the two sandstones form the Chautauqua Hills escarpment. The thickness ranges from a featheredge to 90 feet; average about 60 feet

An unconformity below the Tonganoxie sandstone brings deposits classified as lowermost Virgilian into contact with rocks ranging downward from the Iatan limestone, or perhaps a little higher, at least to the lower part of the Stanton limestone. Paleontological evidence indicating a significant hiatus in sedimentation and evidence of widespread, though not great, erosion support placement of the Missourian-Virgilian boundary in this position. The unconformity is obscure in the northern midcontinent area but it corresponds to the prominent unconformity in southern Oklahoma that records erosion of mountains formed by the Arbuckle orogeny.

~~~~~Unconformity~~~~~


**MISSOURIAN SERIES.**—Middle Pennsylvanian rocks of Kansas, classed as belonging to the Missourian series, lie between two regional disconformities that form boundaries between this division and adjoining strata. The Missourian deposits are essentially distinguished by paleontologic characters and to some extent by peculiarities of cyclic sedimentation. The outcrop area of this series in Kansas is a 20- to 40-mile wide belt, marked by fairly strong east-facing escarpments, extending from Doniphan county in the northeast to Montgomery county in the south. The thickness is about 700 feet.

**Pedee group.**—This topmost division of the Missourian rocks includes the Weston shale below, the Iatan limestone above, and possibly a few feet of sandy and clayey shale above the Iatan limestone. These beds, occurring between the Stanton limestone and the disconformity that defines the top of the Missourian series, are classified as a separate group, because the conformably underlying Lansing group is a compact unit consisting mainly of limestone. Average thickness is 90 feet.

■ **Iatan limestone.**—Light bluish-gray or nearly white limestone. The texture is very fine and dense, but there are numerous thin, irregular plates of clear calcite. Fossil remains are not abundant, but fusulines, brachiopods, bryozoans, crinoid fragments, and small corals are somewhat common locally; algal remains are abundant in the upper part at many places. The formation is extensive in northeastern Kansas but is cut out by the post-Missourian disconformity near Leavenworth and southward at least to the vicinity of southern Douglas county. The thickness ranges from about 4 to 22 feet; average ..... 10 feet

■ **Weston shale.**—Dark bluish to bluish-gray clay shale, generally unfossiliferous but containing thin, fossiliferous limestone beds locally. There are some beds of shaly and even-bedded sandstone toward the south. In many places the formation is characterized by flattish, elliptical concretions of ironstone. At many places the post-Missourian disconformity has cut out most or all of the Weston shale. The thickness ranges from a featheredge to 200 feet; average about ..... 80 feet

PALEOZOIC  
PENNSYLVANIANMissourian  
Pedee



(For explanation see figure 8)

FIG. 7. Generalized section of upper Pennsylvanian rocks in Kansas.

**Lansing group.**—This part of the Missourian includes a rather compact assemblage of two limestones and a thin shale formation. It makes an escarpment that is traced across Kansas readily. The group is recognizable in the subsurface of the eastern and central parts of the state, but in western Kansas it is not clearly separable from underlying Kansas City beds. The thickness averages about 85 feet.

■ **Stanton limestone.**—Three limestone and two shale members. In north-eastern Kansas the formation is rather uniform in character and thickness, but in east-central and southern Kansas there are great local variations. Near the Oklahoma-Kansas boundary the Stanton is difficult to trace. The thickness of the formation ranges from about 10 to 90 feet. The characteristic thickness is about ..... 42 feet

● **South Bend limestone member.**—The topmost division of the Stanton, called "Little Kaw" in some Kansas reports, is composed of dark-gray fine-grained limestone. The lower part is sandy and locally contains some sandy shale. *Fusulines* and the brachiopod *Meekella* are the most common fossils. The thickness is rather constantly about ..... 5 feet

● **Rock Lake shale member.**—Chiefly sandy shale and soft buff sandstone, and commonly a thin layer of greenish or gray clay shale in the lower part. In Anderson county the Rock Lake member locally contains an assemblage of well-preserved land plants, including *Walchia*, mixed with remains of amphibians, fishes, a scorpion, and marine invertebrates. The sandstone is generally even-bedded or massive, and contains marine fossils, mostly mollusks. This member has been designated as "Victory Junction shale" in some reports. The thickness ranges from about 1 to 15 feet; average ..... 6 feet

~~~~~ **Local disconformity?** ~~~~~

● **Stoner limestone member.**—Light bluish-gray to nearly white, mostly thin wavy-bedded limestone having thin shale partings between the layers. The rock is fine-grained, but locally there is much crystalline calcite. It is generally sparingly fossiliferous, but to the north it contains abundant *Triticites*. This limestone, called "Olathe" in some reports, makes a rather prominent escarpment in Kansas. It ranges in thickness from 11 to 40 feet; average 20 feet

● **Eudora shale member.**—Dark-gray and black fissile shale in the lower part and greenish-gray to bluish-gray clay shale in the upper part. Abundantly fossiliferous in northern Kansas but barren of megascopic fossils in northeastern Kansas. Thickness is locally 1 or 2 feet in some northern exposures and as much as 50 feet in part of Montgomery county; average 5 feet

● **Captain Creek limestone member.**—Dark to bluish-gray, granular or dense, brittle limestone. It is massive or even-bedded and in most places shows vertical joints. The upper few inches in some exposures is a brecciated silicified mottled pink and gray bed. The brachiopod *Enteleles pugnoides* is abundant and the fusuline *Triticites neglectus* occurs commonly on bedding planes. Characteristic thicknesses along Kansas river range from 4.5 to 5.5 feet. Locally in Miami county it is a little more than 10 feet thick, and near Independence, in Montgom-

PALEOZOIC
—
PENNSYLVANIAN
—
Missourian
—
Lansing

ery county, limestone 55 feet thick has been assigned to this member; average 6 feet

■ **Vilas shale.**—Gray to buff clayey and sandy shale, sparsely fossiliferous. Locally along Kansas river, hard gray ripple-marked sandstone occurs in the upper part, and farther south a considerable thickness of reddish-brown soft sandstone is found in some exposures. The thickness in the Kansas river valley and northward is about 15 feet, but to the south the thickness locally increases to 90 feet; average 20 feet

■ **Plattsburg limestone.**—Two limestone members separated by a shale member. The upper limestone is the most prominent, except locally in eastern Franklin county. The thickness of the formation ranges from a feather-edge to more than 100 feet at a few places in southern Kansas. The formation disappears near the Oklahoma line, but seemingly reappears farther south and is represented in the upper part of the *Ochelata group*. Average thickness of the Plattsburg limestone in Kansas is about 23 feet

● **Spring Hill limestone member.**—Bluish-gray, fine-grained, brittle, thin-bedded limestone in the lower part and light-gray algal (oölitic) or granular limestone in the upper part. Characteristic fossils include algal remains, several species of sponges, echinoid spines, and the brachiopods *Marginifera*, *Enteletes*, a robust *Composita*, *Neospirifer*, and *Isogramma*. The thickness ranges from 8 to 50 feet or more but is commonly about 15 feet

● **Hickory Creek shale member.**—Gray, green, and black shale. Where thin (0.5 to 2 feet), this member contains a black carbonaceous zone, and where thick (to 15 feet), it is mostly gray or yellowish in color and is clayey. Fossils are rare or lacking in this shale. The thickness in many places is less than 1 foot but averages 5 feet

● **Merriam limestone member.**—Commonly the lower part of this member is drab or light-gray limestone and the upper part is bluish-gray fine-grained limestone. Locally, because of accumulation of algal and granular limestone in the uppermost part, the Merriam is thicker than the overlying Spring Hill limestone. Numerous productids and a few other brachiopods are characteristic. Several species of nautiloid cephalopods are found locally where the member contains much algal (oölitic) limestone. The thickness ranges from less than 1 foot to about 10 feet (eastern Franklin county); average 3 feet

Kansas City group.—As defined in Kansas, this division of the Missourian series includes strata from the top of the Winterset limestone to the base of the Plattsburg limestone. These beds are classified as a separate group, chiefly because both the underlying Bronson deposits and the overlying Lansing strata are each lithologically distinct elements that are well differentiated along the outcrop. Average thickness of the Kansas City group is 275 feet. The Missouri Geological Survey classifies strata from the top of the Argentine limestone to the base of the Hertha limestone in the Kansas City formation.

■ **Bonner Springs shale.**—This formation consists of gray to buff shale, sandy shale, and sandstone. The upper part of the shale consists of olive-green clay shale containing a maroon band near the top in the northern part of the outcrop area. Soft, nodular, cavernous limestone commonly

Lansing

Missourian

PALEOZOIC

PENNSYLVANIAN

PALEOZOIC
PENNSYLVANIAN
Missourian
Kansas City

overlies the maroon shale. Sandstone is rather prevalent in the lower part of the Bonner Springs shale, and in some exposures it occupies definite channels cut in clay shale. South of the point in southeastern Franklin county where the underlying Wyandotte limestone disappears, the Bonner Springs shale rests directly on the Lane shale, and the part of the section between the Iola and Plattsburg limestones is designated as Lane-Bonner Springs shale. The thickness of the Lane-Bonner Springs ranges from 55 to 170 feet, averaging about 90 feet. Locally along Kansas river the Bonner Springs shale is absent; the maximum observed thickness is 60 feet in T. 17 S., R. 20 E.; average about 20 feet

■ **Wyandotte limestone.**—Three limestone and two shale members. The lower two limestones and their separating shale are more constant in lithologic character and thickness, but the entire formation, which is prominent in northeastern Kansas and especially along Kansas river, disappears at a point a short distance southwest of Lane in southeastern Franklin county. It reappears as thin limestone and sandstone beds a few miles north of Garnett in Anderson county, but can be identified for only a few miles. The thickness ranges from a featheredge to 75 feet; omitting consideration of attenuated southern outcrops, the average thickness is about 50 feet

● **Farley limestone member.**—This member comprises an extremely variable assemblage of limestone and shale beds that is recognized only north of T. 14 S. Many types of limestone are represented, but oölitic-appearing algal limestone, limestone breccia or conglomerate, and dense mottled pinkish-gray limestone are characteristic. A thin bed of dark greenish-gray sandy shale containing a molluscan fauna occurs rather persistently in the middle part. Cross bedding is common, both in algal limestone and in breccia. The algal and oölitic facies commonly occur in massive beds; the breccias and conglomerates are slabby. Wavy and thin-bedded, mottled, dense limestone, chiefly in the lower part, is somewhat similar to the main body of the underlying Argentine limestone. The Farley limestone is abundantly fossiliferous, mollusks predominating. The relative quantitative importance of limestone and shale in this member varies from place to place. Thickness ranges from a featheredge to about 35 feet; average 16 feet

● **Island Creek shale member.**—A gray, yellow, and bluish clay shale of highly variable thickness. Locally in Wyandotte county, several feet of sandstone, seemingly a channel filling, occupies this position. The thickness ranges from a featheredge to about 40 feet; average 10 feet

● **Argentine limestone member.**—This is a prominent escarpment-making light bluish-gray limestone, much of which weathers creamy-white. Wavy bedding is common; clay partings are numerous in some outcrops. Marine fossils are plentiful. The thickness ranges from a featheredge to 35 feet; average 20 feet

● **Quindaro shale member.**—Black fissile yellowish-gray calcareous shale, or soft shaly limestone, 1 to 4 feet thick; average 2 feet

● **Frisbie limestone member.**—Bluish-gray to dark-blue massive limestone, locally bearing numerous marine fossils. The thickness commonly ranges from 1 to 3 feet; average 2 feet

■ **Lane shale.**—Dark bluish-gray clay shale, gray and yellowish-brown unfossiliferous sandy shale, and thin-bedded sandstone. The thickness ranges from about 15 to 105 feet. Southward from the point where the Wyandotte limestone pinches out, the Lane and Bonner Springs shales merge. The thickness of the combined units (Lane-Bonner Springs shale) ranges from about 55 feet to 170 feet locally in Allen county. Average thickness of the Lane shale is about 50 feet

■ **Iola limestone.**—Two limestone members and a shale member. The upper limestone is the most conspicuous part of the formation. The thickness ranges from a featheredge along the Kansas-Oklahoma line and locally elsewhere in southern Kansas to about 30 feet in Allen county. In northeastern Kansas the thickness commonly is about 7.5 feet. Average for the formation in Kansas is about 12 feet

● **Raytown limestone member.**—Light-gray wavy-bedded limestone throughout much of the outcrop area; dark-gray to buff more massively bedded limestone in the Kansas river valley, where abundant light-colored large fossil shells give the rock a mottled appearance. The thickness in the Kansas City area is about 5 feet, in Miami county 6 feet, and in the vicinity of Iola 28 feet. It thins southward from Allen county and is absent locally in southern Kansas, but reappears in northern Oklahoma where it is named the *Avant limestone*. Average in Kansas, about 9 feet

● **Muncie Creek shale member.**—Black fissile shale overlain by gray or buff shale containing abundant phosphatic nodules. The thickness commonly ranges from about 1 to 3 feet; average 2 feet

● **Paola limestone member.**—Dark bluish-gray, massive-bedded, brittle, vertical-jointed limestone that weathers bluish-gray. "Worm borings," which are irregular subcylindrical bodies that differ slightly in color and texture from surrounding rock matrix, are more or less characteristic. The thickness ranges from a featheredge to about 2 feet; average 1 foot

■ **Chanute shale.**—Yellowish-brown sandy shale and dark-gray to greenish shale, sandstone, and one coal bed. The sandy shale and sandstone occur chiefly in the upper part, above the coal bed. Members have been differentiated in southern Kansas but are not distinguished north of Allen county. The thickness of the Chanute shale ranges from 12 feet near Kansas City to 165 feet in southern Kansas; average about 75 feet

● **Cottage Grove sandstone member.**—Yellowish-brown or tan thin-bedded to massive sandstone makes up the upper one third to one half of the Chanute shale in southeastern Kansas, but this sandstone occurs only locally in northeastern Kansas. Thickness ranges from a featheredge to 60 feet; average 25 feet

○ **Unnamed clay shale,** locally calcareous and nodular or sandy, occurs in the lower middle part of the Chanute shale. The persistent *Thayer coal*, or a thin bed of underclay, is commonly found at the top of this shale, next below the Cottage Grove sandstone. The Thayer coal ranges in thickness from less than an inch to 2.5 feet. The unnamed shale is 10 to 75 feet thick, the maximum being observed in places where the Noxie sandstone is absent; average thickness 10 feet

PALEOZOIC
PENNSYLVANIAN
Missourian
Kansas City

- **Noxie sandstone member.**—Sandstone at the base of the Chanute shale, commonly filling channels that extend as low as the Stark shale or perhaps lower, is known as the Noxie sandstone member. Locally a limestone conglomerate occurs at the base. This member of the Chanute is recognized only in south-central and southern Kansas where the thickness ranges locally to 100 feet; average 40 feet

The disconformity at the base of the Noxie sandstone is recognized at many places in southern Kansas, especially from T. 28 S. southward.

~~~~~ **Local disconformity** ~~~~~

- **Drum limestone.**—Two limestone members locally separated by a thin shale. The thickness ranges from a featheredge to about 60 feet. The average thickness is about ..... 9 feet

- **Corbin City limestone member.**—Chiefly light-gray oölitic cross-bedded limestone that is highly fossiliferous. This member is best developed in the vicinity of Independence where it is 50 feet or more thick. It is represented by a few feet of limestone conglomerate in southern Kansas near Coffeyville and by 1 foot or less of chiefly algal limestone in the Kansas City area. Fossils are abundant in the thick oölitic facies. The Corbin City limestone is identified at many outcrops in east-central Kansas where the Cement City member of the Drum is not present. A few inches of shale that lies between these two limestones at some exposures is included with the Corbin City member. Average thickness ..... 5 feet

*In southern Kansas the Corbin City limestone rests disconformably on the Cement City limestone. In the vicinity of Cherryvale in Montgomery county the oölitic limestone of the upper member fills hollows nearly 5 feet deep in the lower limestone.*

~~~~~ **Local disconformity** ~~~~~

- **Cement City limestone member.**—Bluish-gray limestone, mottled with brown on weathering, commonly fine-grained or dense, and more or less massive, comprises the Cement City limestone. Marine fossils are generally abundant. A horn coral identified as *Caninia torquia* is characteristic of the upper part of this limestone in northeastern Kansas. The thickness ranges from about 2 to 10 feet; average 4 feet

- **Quivira shale.**—This unit consists chiefly of olive-green clay shale having generally about 1 foot of black or maroon shale in the basal or middle part. Small brachiopods occur in the black facies. The thickness ranges from 3 to 11 feet. The Quivira shale is differentiated only in northeastern Kansas, from Miami county northward. The average thickness is about 7 feet

- **Westerville limestone.**—A persistent unit northward from Miami county, but variable in lithology. Cross-bedded oölite is characteristic in the Kansas City area and accordingly this limestone has been designated frequently as the "Kansas City oölite." Flint or chert, which is light pink and hence differs from other Pennsylvanian cherts, comprises 50 percent or more of the rock in some outcrops. The thickness ranges from less than a foot to 16 feet; average about 8 feet

PALEOZOIC

PENNSYLVANIAN

Missourian

Kansas City

Kansas City

■ **Wea shale.**—Chiefly olive-green clay shale containing a persistent thin zone of maroon silty shale in the upper part. For about 50 miles south of the place in Miami county where the Westerville limestone pinches out, it is feasible to designate the interval between the Block limestone and the Drum limestone as the Wea-Quivira shale. This combined unit is 30 to 85 feet thick. The thickness of the Wea, where it can be differentiated, ranges from about 15 to 35 feet; average 25 feet

■ **Block limestone.**—Bluish-gray, fine-grained, hard limestone. The rock is commonly massive and displays vertical jointing, but locally it is thin-bedded. In the Kansas City area the unit is represented by thin blue limestone beds separated by shale. Fusulines are generally common, and locally *Marginifera wabashensis* is abundant. The Block limestone has not been identified south of Linn county. Observed thicknesses range from about 3 to 8 feet; the common thickness is about 4 feet

■ **Fontana shale.**—Greenish-gray to buff shale bearing scattered calcareous nodules is defined under the name Fontana northward from Linn county. South of Linn county, where the Westerville and Block limestones have not been identified, the interval between the Drum and Winterset limestones is known as the Fontana-Quivira or *Cherryvale shale*. Near Cherryvale this unit comprises about 60 feet of bluish-gray silty shale containing layers of blue dense limestone near the top. In the vicinity of Coffeyville, similar limestone occurs only a few feet above the Winterset limestone. The Fontana shale ranges in thickness from about 5 to 25 feet; average about 15 feet

■ **Bronson group.**—Predominantly an assemblage of limestones, this group comprises beds from the top of the Dennis limestone to the base of the Hertha limestone. These limestones, together with the Swope limestone and separating shale formations, form a compact natural grouping of lower Missourian strata throughout most of the Kansas outcrop area. The Missouri Geological Survey, however, includes the Bronson rocks as a part of the Kansas City formation. The Bronson group ranges in thickness from about 85 feet in the Kansas City area to 175 feet in southern Kansas; shale predominates in the latter area. Average thickness is about 135 feet.

■ **Dennis limestone.**—The uppermost formation of the Bronson group contains two limestone members and a shale member. The capping limestone forms a long dip slope that terminates eastward at the prominent Bronson escarpment. In Oklahoma this formation is known as the *Hogshooter limestone*. The thickness of the Dennis ranges from 2 feet (in parts of Neosho county) to 60 feet (in northeastern Labette county); average about 40 feet

● **Winterset limestone member.**—Light bluish-gray and light-gray limestone characterized by abundant flint. In the Kansas river area the flint is almost black but elsewhere it is light gray. Along much of the outcrop belt cross-bedded oölite occurs in the upper part of the Winterset and there is much light-gray thin-bedded limestone having a brecciated appearance. Black platy shale containing land plant remains occurs in the middle part of the member in northern outcrops and calcareous gray shale occupies the same position in southern Kansas. Marine invertebrates are more or less abundant in various

PALEOZOIC
PENNSYLVANIAN

Missourian

Bronson

PALEOZOIC

PENNSYLVANIAN

Missourian

Bronson

zones; in places there are well-preserved molluscan faunas. Locally in Neosho county the Winterset limestone and possibly lower beds are cut out by the disconformity at the base of the Chanute shale. The thickness ranges from a featheredge to about 50 feet; average .. 36 feet

● *Stark shale member*.—Chiefly black shale but containing some gray and yellow shale in the upper part. The member is recognized only locally in Montgomery county and neighboring parts of Oklahoma but becomes persistent northward from northwestern Labette county. The common thickness is about 3 feet

● *Canville limestone member*.—Medium dark-gray dense to granular limestone, commonly massive but locally platy or slabby. This limestone is persistent from northwestern Labette county to Linn county. It is represented in parts of Montgomery county by about 1 foot of dark bluish-gray vertical-jointed limestone, and it reappears locally in Oklahoma. Thickness ranges from a featheredge to about 2 feet; average 1 foot

■ *Galesburg shale*.—Gray and yellow marine shale, sandy nonmarine shale, sandstone, and a little coal. Probably some of the sandy shale and sandstone is marine. In the Kansas river valley, the unit is about 3 feet thick and comprises calcareous nodular shale that is clearly distinguished by its light color and other lithologic features from the overlying black Stark shale. The thickness increases rather abruptly southward from Bourbon county and much sandstone is present in Neosho, Labette, and Montgomery counties. Plant fossils are fairly common in the sandy facies. A coal bed in the upper part of the Galesburg in southern Kansas is called the *Cedar Bluff coal*. Southward from the point where the Swope limestone disappears, a few miles north of the Oklahoma line, the Galesburg is not separable as a unit and it forms the upper part of the *Coffeyville formation*. The thickness of the Galesburg shale ranges from 3 to 75 feet; average about 35 feet

● *Dodds Creek sandstone member*.—Massive to thin-bedded sandstone, seemingly of deltaic origin, which occurs in the Galesburg shale in southern Kansas, is named the Dodds Creek sandstone member. The thickness of this subdivision ranges upward to about 40 feet

■ *Swope limestone*.—Two limestone members and a shale member. The upper limestone is the most persistent, but it is not recognized south of a point in eastern Montgomery county (northern part of T. 34 S.). The thickness ranges from a featheredge to about 35 feet; average 23 feet

● *Bethany Falls limestone member*.—Light-gray, dense, thin-bedded limestone overlain by mottled gray, massive, algal limestone or by nearly white oölitic limestone that is commonly cross-bedded. Fossils are more or less common in the lower thin-bedded part. The thickness of the thin-bedded division ranges from 1 to 20 feet and that of the massive division from 7 to 15 feet. The total thickness ranges from about 12 to 30 feet; average about .. 18 feet

● *Hushpuckney shale member*.—Bluish-gray clay shale in the upper part, black fissile shale in the lower part. This unit and the underlying Middle Creek limestone are persistent northward from Neosho county. The thickness ranges from a featheredge to about 6 feet; average 3 feet

Bourbon

PALEOZOIC
PENNSYLVANIAN
Missourian

● *Middle Creek limestone member.*—Dark bluish-gray limestone, commonly dense and brittle, vertical-jointed. The maximum observed thickness is in southern Linn county where it is about 8 feet; the common thickness is about 2 feet

■ *Ladore shale.*—Gray and brownish-yellow shale, sandstone, and some coal. Like the Galesburg shale, the formation thickens southward. The Ladore is a thin marine shale in the Kansas City area and east-central Kansas, but it expands into a sandy deltaic deposit in southern Kansas. In Montgomery and southern Labette counties it is not separated from the overlying Galesburg shale and the two form the upper more sandy part of the *Coffeyville formation*. Thin coal beds are present in Labette county, and plant fossils are plentiful there. The thickness ranges from 2 to 50 feet; average about 20 feet

■ *Hertha limestone.*—Two limestone members and a shale member. The two limestones are persistent and distinct throughout much of the Kansas outcrop area, especially in Miami, Linn, and northern Bourbon counties. In southern outcrops the shale member is thin or absent. The formation is recognized northward from a point in T. 33 S., Labette county. The thickness ranges from a featheredge to about 30 feet; average 16 feet

● *Sniabar limestone member.*—Gray and brown limestone, varying from more or less massive to thin-bedded. Marine fossils are moderately common and locally the coral "*Aulopora*" is abundant. The thickness ranges from a featheredge to about 10 feet; average 5 feet

● *Mound City shale member.*—Gray and yellow clayey to calcareous shale, containing a persistent 2-inch bed of crinoidal limestone in Linn and Bourbon counties. Marine fossils are locally plentiful. The thickness ranges from a featheredge to about 14 feet; average about.....6 feet

● *Critzer limestone member.*—Massive, brownish-gray, granular, partly algal limestone, and thin wavy-bedded gray limestone. In a part of the outcrop area this member is the most conspicuous part of the Hertha limestone, commonly forming a rim rock cliff along the Hertha escarpment. Much of the rock weathers deep brown. Bellerophontid gastropods are characteristic of the massive facies; brachiopods and corals are plentiful locally in the thin-bedded facies. The thickness ranges from a featheredge to about 11 feet; average about..... 5 feet

(?) Local disconformity

Bourbon shale.—Rocks lying between the base of the Hertha limestone and the disconformity that separates Missourian from Desmoinesian beds are mainly clastic sediments that for the most part represent mechanically weathered detritus derived from land and deposited in shallow seas which advanced over Kansas after a time of more or less prolonged emergence. Gray, yellow, and dark-gray to black clay shale predominates, but there is much sandstone and some limestone and coal. The thickness ranges from about 70 feet to 130 feet; average..... 100 feet

Bourbon

□ **Sandstone and interbedded black shale and flaggy limestone.**—In Linn and Miami counties the upper part of this division is partly occupied by massive sandstone (*Knobtown*) which is locally as much as 25 feet thick. In southern Linn and northern Bourbon counties, thin beds of dense blue limestone alter-

nating with thin beds of nearly black shale occupy the same position and attain a thickness of 45 feet. These deposits are absent for a short distance farther south, but black shale occurs in this position in Neosho and Labette counties. Near the point in Labette county where the Hertha limestone disappears, about 40 feet of black shale forms the top of the Bourbon section. Average thickness of this division is about 30 feet

□ **Light-colored shale.**—Gray to yellowish shale that locally contains a few thin limestone beds comprises most of the middle and lower parts of the Bourbon. This facies commonly extends downward to the Hepler sandstone. Fine-grained sandstone or siltstone forms the middle part of the Bourbon in some places. In southern Bourbon and northern Labette counties, light-colored shale extends to the base of the Hertha limestone. The thickness of this shale unit ranges from about 20 to 100 feet; average about 60 feet

■ **Checkerboard limestone.**—Gray limestone, locally a coquina of gastropods or crinoid fragments; where more fully developed this formation comprises two thin limestones separated by dark shale. The Checkerboard limestone is an important marker in the Pennsylvanian strata of eastern Oklahoma and is recognized in southern Kansas as far north as T. 31 S. The thickness ranges from a featheredge to about 6 feet; average 2 feet

■ **Hepler sandstone.**—Brown and gray sandstone named Hepler is a remarkably persistent sheetlike deposit overlying the regional disconformity at the base of the Missourian series. A calcareous facies of the Hepler sandstone is seen locally in southern Kansas. The thickness ranges from about 2 to 20 feet; average 10 feet

The unconformity below the Hepler sandstone brings deposits classified as lowermost Missourian into contact with rocks ranging from the Memorial shale downward to the upper part of the Bandera shale. Paleontological evidence and indication of widespread interruption in sedimentation, accompanied by some erosion, support placement of the Desmoinesian-Missourian boundary at this position.

Unconformity

Desmoinesian series.—The lowermost major division of the Pennsylvanian rocks in Kansas, now recognized, is named the Desmoinesian series. It is based on outcrops in Iowa. The series occupies the stratigraphic interval between important regional disconformities. The Desmoinesian deposits are set off from the overlying Missourian rocks by pronounced paleontologic and lithologic differences.

Marmaton group.—The upper 250 feet, approximately, of the Desmoinesian beds in Kansas is assigned to the Marmaton group. These strata are more calcareous and more dominantly marine than those of the underlying Cherokee shale. The Missouri Geological Survey now employs the term "Henrietta group" in a sense that makes it synonymous with Marmaton, which is the older, more widely used name.

■ **Memorial shale.**—Gray and yellowish clay shale, generally unfossiliferous. In Labette county it comprises dark-gray shale that bears marine fossils and contains two or more coal beds. The unit locally is absent owing to pre-Missourian erosion. The maximum known thickness is 30 feet, but the average is about 10 feet

PALEOZOIC

PENNSYLVANIAN

Missourian

Bourbon

Desmoinesian

Marmaton

PALEOZOIC
PENNSYLVANIAN
Desmoinesian
Marmaton

■ **Lenapah limestone.**—Two limestone members separated by a shale member. The formation is locally reduced in thickness or absent as a result of pre-Missourian erosion. The lower two members are absent in some places owing to nondeposition. The Lenapah is well marked in southeastern Kansas but inconspicuous in east-central Kansas. Thickness ranges from 1 to 18 feet; average 12 feet

● **Idenbro limestone member.**—Chiefly light-gray crystalline and pseudobrecciated limestone; in Linn and Bourbon counties represented by brownish-gray to dark-gray limestone containing limonitic inclusions, mostly crystalline, but dense and earthy locally. This member forms low escarpments and broad dip slopes in the southern part of the outcrop area. Characteristic thickness in the northern part is 1 to 2 feet; in the southern part it is 6 to 9 feet; average 4 feet

● **Perry Farm shale member.**—Light- to dark-gray clay shale containing nodules of limestone; grades laterally into nodular limestone south of the Kansas-Oklahoma line. The shale is abundantly fossiliferous locally, brachiopods occurring in the upper part and a molluscan fauna in the middle and lower parts. The member is locally absent in the northern part of the outcrop area. The thickness ranges from a featheredge to about 20 feet; average 7 feet

● **Norfleet limestone member.**—Dark brownish-gray massive to bluish-gray slabby limestone. The Norfleet resembles local facies of the Idenbro limestone in some northern outcrops. South of the Kansas-Oklahoma line, this member grades into massive dense limestone that is continuous with overlying nodular limestone facies of the Perry Farm shale, but distinction of the two members is clear. The thickness ranges from a few inches to about 3 feet; average 1.5 feet

■ **Nowata shale.**—Gray to yellow clay shale, sandy shale, and sandstone; marine and nonmarine. The formation is locally absent in Linn county and in the subsurface of Miami county, owing to pre-Missourian erosion. Measured thicknesses in Kansas, 3 to 30 feet; average 18 feet

● **Walter Johnson sandstone member.**—Thin-bedded to massive sandstone in the lower and middle Nowata shale, observed in parts of southern Kansas, has been named the Walter Johnson sandstone. It is equivalent to the "Wayside sand" of the subsurface. Outcrop thicknesses are commonly 4 to 10 feet

■ **Altamont limestone.**—Two limestone members separated by a shale member. The lower limestone is most prominent in the southern part of the outcrop area, and the upper limestone is most prominent in northern outcrops. The formation is locally absent in the subsurface of Miami county owing to pre-Missourian erosion. The thickness is about 6 to 25 feet; average..... 19 feet

● **Worland limestone member.**—Light-gray limestone, dense, semi-lithographic in upper part, remainder mostly crystalline, pseudo-brecciated and wavy-bedded. Marine fossils characteristically are few but there are prominent brachiopods and fusulines, the latter locally abundant. Thickness, commonly about 8 feet

● **Lake Neosho shale member.**—Light- and dark-gray and black shale; generally black in middle part, locally all black; containing

PALEOZOIC
PENNSYLVANIAN
Desmoinesian
Marnaton

abundant phosphatic concretions. The thickness ranges from 2 to 8 feet; average 5 feet

● *Tina limestone member*.—Light-gray, massive, coralline (*Chaetetes*) and dense limestone, associated with two shale beds in the southern part of the outcrop area. The limestone is darker, more granular and locally cross-bedded in the northern part. The thickness is 1 to 2 feet in northern part; about 10 feet in southern part; average..... 6 feet

■ *Bandera shale*.—Mainly nonmarine gray, yellow, and maroon clay shale, mostly blocky, well-bedded; massive to thin-bedded sandstone; and the *Mulberry coal* bed near the base. Maroon bands occur in the upper part. The formation contains limonitic concretions and veins and locally septarian limestone concretions. Plant fossils are found in the sandy facies, generally in the lower part. The Mulberry coal is a persistent bed in the lower part of the Bandera north of Crawford county. Shale below the coal is light to dark gray and carbonaceous. The thickness ranges from about 20 to 50 feet; average 35 feet

● *Bandera Quarry sandstone member*.—Gray and brown sandstone, generally thin-bedded, locally "flagstones," lenticular. The thickness ranges from a featheredge to 30 feet; average about 15 feet

■ *Pawnee limestone*.—The upper main part of the Pawnee formation comprises two persistent limestone members separated by a shale member; the lower part consists of shale and a thin nonpersistent basal limestone. Thickness ranges from about 15 to 60 feet; average about 30 feet

● *Laberdie limestone member*.—Light-gray, crystalline, and coralline (*Chaetetes*) limestone in thin wavy to massive beds that form extensive dip slopes and cap prominent escarpments. Fusulines and other marine fossils are locally plentiful. The thickness is about 10 to 20 feet; average 15 feet

● *Mine Creek shale member*.—Light- and dark-gray to black shale, flaky, platy and fissile, containing one or more thin limestone beds in the middle part locally. Generally brachiopods are abundant, corals fewer. This member is reduced to very slight thickness or is absent in the southern part of the outcrop area. The thickness ranges from a featheredge to 15 feet; average 8 feet

● *Myrick Station limestone member*.—Dark-gray, brown, and light-gray limestone. The lower part, comprising about 2 feet of massive dark-gray brown-weathering limestone, is most persistent, and this is overlain locally by lighter colored coralline (*Chaetetes*) limestone. The thickness ranges from 2 to about 8.5 feet; average 5 feet

● *Anna shale member*.—Black, platy, locally fissile shale, containing a few inches of gray shale near the top in some places; small smooth phosphatic concretions are common. A thin bed or lenses of dark, slabby, crystalline, locally crinoidal limestone occur in the basal part. Thickness commonly ranges from 2 to 5 feet; average 3 feet

■ *Labette shale*.—Gray and yellow clay shale, sandy shale, sandstone, and thin coal and limestone beds; a rather persistent black shale occurs in the basal part. Black shale occurs locally in the upper part. Some sections show several thin limestone and coal beds. A persistent limestone, 1 to 2 feet thick,

occurs in the middle lower part. Thickness ranges from about 30 to about 100 feet; average 50 feet

● *Englevale sandstone member*.—This is one of several bodies of sandstone, collectively referred to as *Warrensburg sandstone*, lying in the middle and lower parts of the Labette shale. This is the "Peru sand" of the subsurface. Average thickness, about 30 feet

■ **Fort Scott limestone**.—Comprises two limestones and an intervening shale member, the latter containing a persistent coal and a limestone bed in the northern part of the outcrop area. Characteristic thickness 33 feet

● *Higginsville limestone member*.—Chiefly light-gray thin-bedded to massive limestone characterized by the scattered occurrence of unusually large crinoid stems, irregular distribution of minute, medium, and large fusulines, and colonies of *Chaetetes* that generally are isolated but locally form the entire upper part. Characteristic thickness 15 feet

● *Little Osage shale member*.—Black platy shale containing small smooth phosphatic concretions, and gray clayey calcareous shale, coal, and limestone. Upper part generally light gray and calcareous, lower part commonly black and platy. The *Houx limestone* is a persistent thin bed of granular limestone that occurs a few inches to 2 or more feet below the Higginsville limestone. The *Summit coal*, which is persistent from northern Crawford county northeastward into Missouri, occurs in the upper middle part of the Little Osage shale. Thickness of the member ranges from about 4 to 12 feet; average 8 feet

● *Blackjack Creek limestone member*.—Brownish-gray, granular, and massive to light-gray, crystalline, pseudobrecciated, thinner bedded, and locally *Chaetetes*-bearing. The lower part, the only persistent part of the member, is approximately 2 feet thick and consists of brownish-gray, brown-weathering "cement rock." Thickness ranges from about 4 to 17.5 feet; average 10 feet

Cherokee shale.—Pennsylvanian rocks below the Fort Scott limestone in Kansas, which consist mainly of clastic rocks with light- and dark-colored shale predominating, have been classed as a formation in some reports and as a group in others. There is much sandstone and sandy shale, and the most important coal beds of the state belong to the Cherokee. The amount of limestone is small. The unit is not readily divisible into formations unless cyclic units (cyclothems) are so defined. Following are the more generally recognized stratigraphic units. The outcrop area of Cherokee beds in Kansas is confined to counties in the southeastern corner of the state. The thickness at the outcrop ranges from about 400 to 600 feet; average 450 feet

Mulky coal.—The upper few feet of the Cherokee shale consists of black platy shale underlain by a persistent coal bed, the Mulky. The black shale is characterized by small phosphatic concretions, and locally it contains large black limestone concretions. The coal is underlain by gray shale. It ranges in thickness from a featheredge to about 2 feet; average 1 foot

Breezy Hill limestone.—Lenticular sandy limestone occurring a few feet below the Mulky coal. Thickness ranges from a featheredge to about.... 2.5 feet

Marmaton

Desmoinesian

Cherokee

PALEOZOIC
PENNSYLVANIAN




FIG. 8. Generalized section of middle Pennsylvanian rocks in Kansas.

PALEOZOIC
PENNSYLVANIAN
Desmoinesian
Cherokee

Unnamed sandstone.—Lenticular sandstone bodies in the upper part of the Cherokee shale are designated as the "Squirrel sand."

Bevier coal.—A coal bed a short distance above the Ardmore limestone. Thickness averages about 2 feet

Ardmore limestone.—Persistent bed of dark-gray granular limestone 65 to 135 feet below the Fort Scott limestone. This unit is called *Verdigris limestone* in Oklahoma. The common thickness is about 2 feet

Croweburg coal.—Coal bed a few feet below the Ardmore limestone; characterized by overlying and underlying "fire clay." Average thickness about 1 foot

Fleming coal.—Second coal bed below the Ardmore limestone; average thickness about 1 foot

Mineral coal.—This coal, which is extensively mined by stripping, is overlain by a thin limestone bed. It occurs about 30 feet below the Ardmore limestone. The average thickness of the coal is about 1.8 feet

"Pilot" coal.—Thin coal bed about 27 feet above the Weir-Pittsburg bed. Thickness about 0.75 feet

Weir-Pittsburg coal.—The most important commercial coal bed in Kansas, which occurs about 115 feet below the Ardmore limestone, ranges in thickness from a featheredge to about 3.5 feet; average about 3 feet

Bluejacket sandstone.—Sandstone lenses in the lower middle part of the Cherokee section in Kansas and northern Oklahoma are named the Bluejacket sandstone at the outcrop and are known as "Bartlesville sand" in the subsurface. Thickness ranges to at least 50 feet

Columbus coal.—Coal bed a few feet below the Bluejacket sandstone; common thickness about 1 foot

Little Cabin sandstone.—Sandstone lenses at or near the base of the Kansas Cherokee section, which are correlated with the *Warner sandstone* in the McAlester formation of eastern Oklahoma. The sandstone commonly occurs in two or more beds separated by shale. Thickness is commonly 25 to 30 feet.

Riverton coal.—Coal bed at the base of the Little Cabin sandstone, in part of southeastern Kansas. It occurs a few feet above the base of the Cherokee section. Average thickness, about 2 feet

A regional unconformity of much importance defines the base of Pennsylvanian deposits in the northern midcontinent. At the outcrop, lowermost Pennsylvanian deposits of this region (but belonging stratigraphically far above the lowermost Pennsylvanian of some other regions) commonly lie parallel on older rocks, but the unconformable contact locally has topographic irregularity amounting to 100 feet. In Kansas, outcropping rocks next below the Pennsylvanian belong to the lower part of the Mississippian system, but in Missouri and other near-by states, the Pennsylvanian is found resting on Devonian, Silurian, Ordovician, Cambrian, or pre-Cambrian rocks. Beneath the sur-

face in Kansas, the basal Pennsylvanian unconformity is locally angular and pre-Cambrian or lower Paleozoic rocks occur in places next below the Pennsylvanian.

~~~~~ Regional unconformity ~~~~~

**MISSISSIPPIAN SYSTEM.**—Rocks of Mississippian age consisting of light-gray cherty limestone, medium to coarse crystalline in texture, crop out in the extreme southeastern corner of Kansas. The area of exposure is about 100 square miles, but in adjoining parts of Missouri and Oklahoma the outcrop area is very broad. These rocks contain the rich lead and zinc ores of the Tri-State district, and they contain commercial oil and gas accumulations in several Kansas fields. Late Mississippian rocks belonging to the Chesterian series occur a short distance south of the Kansas-Oklahoma line and probably extend a short distance into Kansas at the outcrop. They are important beneath the surface in much of Kansas. Total thickness of outcropping formations of the Mississippian in Kansas is about 50 feet.

**MERAMECIAN SERIES.**—The Mississippian limestones exposed in southeastern Kansas include some beds of Warsaw age that are classed as belonging to the Meramecian series.

■ **Warsaw limestone.**—These limestones consist of bluish-gray cherty limestone containing glauconite near the base. They have been included in the so-called "Boone formation." The thickness of these rocks is about ..... 15 feet

*A break in sedimentation that has considerable stratigraphic importance separates rocks of Meramecian age from older deposits.*

~~~~~ Disconformity ~~~~~

OSAGIAN SERIES.—This division of the Mississippian system is mainly characterized by abundance of crinoidal remains and common occurrence of chert in the limestone beds. These limestones have an average thickness of 200 feet in Missouri.

■ **Keokuk limestone (?)**.—Light bluish-gray, fine- to medium-grained, cherty limestone, commonly in beds 0.5 to 2 feet thick, characterized by the common occurrence of fossil bryozoans along with other marine organisms. It has been included in the "Boone formation." Thickness about 35 feet

No rocks older than Keokuk are known to be exposed at the surface in Kansas.

IGNEOUS ROCKS.—Outcrops of igneous rocks of unknown age occur in two Kansas counties. In Riley county three small pluglike masses of dark basic rock are exposed, and in Woodson county a granite dike is exposed. The Riley county igneous rocks are intruded in sedimentary rocks of Wolfcampian age, and the Woodson county granite cuts rocks of Missourian age.

□ **Granite.**—A dike of coarse-grained granite occurs in Woodson county. The outcrop extends from near the center of sec. 13, T. 26 S., R. 16 E. eastward to a point slightly beyond the township line.

□ **Basic volcanic rock.**—Three outcrops of basic volcanic rock occur in Riley county. The locations of the three small outcrops are: (1) sec. 22, T. 8 S., R. 5 E.; (2) sec. 23, T. 8 S., R. 6 E.; (3) sec. 6, T. 9 S., R. 5 E.

PALEOZOIC

MISSISSIPPIAN




FIG. 9. Generalized section of lower Pennsylvanian rocks in Kansas.

BIBLIOGRAPHY OF OUTCROPPING ROCKS¹

CENOZOIC

Eastern Kansas

- FRYE, J. C., 1941, Reconnaissance of ground-water resources in Atchison county, Kansas: Kansas Geol. Survey, Bull. 38, pt. 9, pp. 237-260, figs. 1-6, pls. 1-3.
- KAY, G. F., and APFEL, E. T., 1928, The pre-Illinoian Pleistocene geology of Iowa: Iowa Geol. Survey Ann. Rept., vol. 34, pp. 1-304, figs. 1-63, pls. 1-3.
- SCHOEWE, W. H., 1923, Glacial geology of Kansas: Pan-Am. Geologist, vol. 40, no. 2, pp. 102-110, pl. 1.
- , 1930, Evidences for a relocation of the drift border in eastern Kansas: Jour. Geology, vol. 38, no. 1, pp. 67-74, figs. 1, 2.
- , 1938, The west Atchison glacial section (abstract): Kansas Acad. Sci. Trans., vol. 41, p. 227.
- , 1939, Evidences for the relocation of west drift border in eastern Kansas: Kansas Acad. Sci. Trans., vol. 42, p. 367, fig. 1.
- TODD, J. E., 1918, Kansas during the ice age: Kansas Acad. Sci. Trans., vol. 28, pp. 33-47.
- , 1918, History of Kaw lake: Kansas Acad. Sci. Trans., vol. 28, pp. 187-199, figs. 1-3.

Central Kansas

- FRYE, J. C., and HIBBARD, C. W., 1941, Stratigraphy and paleontology of a new middle and upper Pliocene formation of south-central Kansas: Jour. Geology, vol. 49, no. 3, pp. 261-278, figs. 1-6.
- FRYE, J. C., LEONARD, A. B., and HIBBARD, C. W., 1943, Westward extension of the Kansas "*Equus* beds": Jour. Geology, vol. 51, no. 1, pp. 33-47, figs. 1-3.
- HAWORTH, ERASMUS, and BEEDE, J. W., 1897, the McPherson *Equus* beds: Kansas Univ. Geol. Survey, vol. 2, pp. 285-296.
- KNIGHT, G. L., 1934, Gerlane formation (abstract): Geol. Soc. America, Proc. 1933, p. 91.
- LOHMAN, S. W., and FRYE, J. C., 1940, Geology and ground-water resources of the "*Equus* beds" area in south central Kansas: Econ. Geology, vol. 35, no. 7, pp. 839-866, figs. 1-5.
- WING, M. E., 1930, The geology of Cloud and Republic counties, Kansas: Kansas Geol. Survey, Bull. 15, pp. 1-51, figs. 1, 2, pls. 1-18.

Southwestern Kansas

- FRYE, J. C., 1942, Geology and ground-water resources of Meade county, Kansas: Kansas Geol. Survey, Bull. 45, pp. 1-152, figs. 1-13, pls. 1-12.
- FRYE, J. C., and HIBBARD, C. W., 1941, Pliocene and Pleistocene stratigraphy and paleontology of the Meade basin, southwestern Kansas: Kansas Geol. Survey, Bull. 38, pt. 13, pp. 389-424, figs. 1-3, pls. 1-4.

¹ References that give the most adequate description of the formations included in this report have been selected. The paper in which a formation was originally named is not necessarily included if it has been superseded by a more recent or more comprehensive discussion. The references are grouped by stratigraphic divisions.

- HIBBARD, C. W., 1941, New mammals from the Rexroad fauna, Upper Pliocene of Kansas: *Am. Midland Naturalist*, vol. 26, no. 2, pp. 337-368, figs. 1-12, pls. 1-3.
- , 1941, The Borchers fauna, a new Pleistocene interglacial fauna from Meade county, Kansas: *Kansas Geol. Survey, Bull.* 38, pt. 7, pp. 197-220, pls. 1, 2.
- , 1944, Stratigraphy and vertebrate paleontology of Pleistocene deposits of southwestern Kansas: *Geol. Soc. America Bull.*, vol. 55, no. 6, pp. 707-754, figs. 1-20, pls. 1-3.
- JOHNSON, W. D., 1901, The High Plains and their utilization: *U.S. Geol. Survey, 21st Ann. Rept.*, pt. 4, pp. 601-741, pls. 113-156; *22d Ann. Rept.*, pt. 4, pp. 631-669, pls. 51-65.
- LATTA, B. F., 1941, Geology and ground-water resources of Stanton county, Kansas: *Kansas Geol. Survey, Bull.* 37, pp. 1-119, figs. 1-6, pls. 1-9.
- , 1944, Geology and ground-water resources of Finney and Gray counties, Kansas: *Kansas Geol. Survey, Bull.* 55, pp. 1-272, figs. 1-21, pls. 1-12.
- McGREW, P. O., 1944, An early Pleistocene (Blancan) fauna from Nebraska: *Field Mus. Nat. History, Geol. Ser.*, vol. 9, no. 2, pp. 33-66, figs. 14-22.
- McLAUGHLIN, T. G., 1942, Geology and ground-water resources of Morton county, Kansas: *Kansas Geol. Survey, Bull.* 40, pp. 1-126, figs. 1-6, pls. 1-9.
- , 1943, Geology and ground-water resources of Hamilton and Kearny counties, Kansas: *Kansas Geol. Survey, Bull.* 49, pp. 1-220, figs. 1-18, pls. 1-17.
- SMITH, H. T. U., 1940, Geologic studies in southwestern Kansas: *Kansas Geol. Survey, Bull.* 34, pp. 1-212, figs. 1-22, pls. 1-34.
- WAITE, H. A., 1942, Geology and ground-water resources of Ford county, Kansas: *Kansas Geol. Survey, Bull.* 43, pp. 1-250, figs. 1-22, pls. 1-16.

Northwestern Kansas

- DARTON, N. H., 1899, Preliminary report on the geology and water resources of Nebraska west of the one hundred and third meridian: *U.S. Geol. Survey, 19th Ann. Rept.*, pt. 4, pp. 719-785, pls. 74-118. Reprinted as *U.S. Geol. Survey, Prof. Paper* 17 (1903).
- ELIAS, M. K., 1931, The geology of Wallace county, Kansas: *Kansas Geol. Survey, Bull.* 18, pp. 1-254, figs. 1-7, pls. 1-42.
- , 1937, Geology of Rawlins and Decatur counties with special reference to water resources: *Kansas Geol. Survey, Mineral Resources Circ.* 7, pp. 1-25, figs. 1-4.
- HAWORTH, ERASMUS, 1897, Physical properties of the Tertiary: *Kansas Univ. Geol. Survey*, vol. 2, pp. 247-284, pls. 36-44.
- HIBBARD, C. W., 1939, Notes on additional fauna of Edson Quarry of the Middle Pliocene of Kansas: *Kansas Acad. Sci. Trans.*, vol. 42, pp. 457-462, figs. 1-6.
- HIBBARD, C. W., FRYE, J. C., and LEONARD, A. B., 1944, Reconnaissance of Pleistocene deposits in north-central Kansas: *Kansas Geol. Survey, Bull.* 52, pt. 1, pp. 1-28, figs. 1, 2, pls. 1, 2.

LEONARD, A. B., and FRYE, J. C., 1943, Additional studies of the Sanborn formation, Pleistocene, in northwestern Kansas: *Am. Jour. Sci.*, vol. 241, pp. 453-462, figs. 1-3, pl. 1.

MESOZOIC

CRETACEOUS

BASS, N. W., 1926, The geology of Ellis county, Kansas: *Kansas Geol. Survey, Bull. 11*, pt. I, pp. 11-52, figs. 1-12, pls. 1-4.

———, 1926, The geology of Hamilton county, Kansas: *Kansas Geol. Survey, Bull. 11*, pt. II, pp. 53-83, figs. 13-26, pls. 5, 6.

DOTT, R. H., 1941, Regional stratigraphy of Mid-Continent: *Am. Assoc. Petroleum Geologists Bull.*, vol. 25, no. 9, pp. 1619-1705, figs. 1-8.

ELIAS, M. K., 1931, The geology of Wallace county, Kansas: *Kansas Geol. Survey, Bull. 18*, pp. 1-254, figs. 1-7, pls. 1-42.

FRYE, J. C., and BRAZIL, J. J., 1943, Ground water in the oil-field areas of Ellis and Russell counties, Kansas: *Kansas Geol. Survey, Bull. 50*, pp. 1-104, figs. 1-9, pls. 1, 2.

LANDES, K. K., 1930, The geology of Mitchell and Osborne counties, Kansas: *Kansas Geol. Survey, Bull. 16*, pp. 1-55, fig. 1, pls. 1-15.

———, and KEROHER, R. P., 1942, Mineral resources of Phillips county: *Kansas Geol. Survey, Bull. 41*, pt. 8, pp. 277-312, figs. 1-5, pls. 1-4.

LATTA, B. F., 1941, Geology and ground-water resources of Stanton county, Kansas: *Kansas Geol. Survey, Bull. 37*, pp. 1-119, figs. 1-6, pls. 1-9.

LOGAN, W. N., 1897, The Upper Cretaceous of Kansas: *Kansas Univ. Geol. Survey, vol. 2*, pp. 199-234, figs. 10, 11, pls. 25-34.

McLAUGHLIN, T. G., 1943, Geology and ground-water resources of Hamilton and Kearny counties, Kansas: *Kansas Geol. Survey, Bull. 49*, pp. 1-220, figs. 1-18, pls. 1-17.

MOORE, R. C., and LANDES, K. K., 1937, Geologic map of Kansas, scale 1:500,000. *Kansas Geol. Survey.*

MOSS, R. G., 1932, The geology of Ness and Hodgeman counties, Kansas: *Kansas Geol. Survey, Bull. 19*, pp. 1-48, fig. 1, pls. 1-7.

PLUMMER, NORMAN, and ROMARY, J. F., 1942, Stratigraphy of the pre-Greenhorn Cretaceous beds of Kansas: *Kansas Geol. Survey, Bull. 41*, pt. 9, pp. 313-348, figs. 1-4, pls. 1, 2.

PROSSER, C. S., 1897, The Upper Permian and the Lower Cretaceous: *Kansas Univ. Geol. Survey, vol. 2*, pp. 55-194, figs. 2-9, pls. 9-24.

RUBEY, W. W., and BASS, N. W., 1925, The geology of Russell county, Kansas: *Kansas Geol. Survey, Bull. 10*, pp. 1-104, figs. 1-11, pls. 1-7.

TESTER, A. C., 1931, The Dakota stage of the type locality: *Iowa Geol. Survey, vol. 35*, pp. 195-332, figs. 25-44, pls. 3, 4.

TWENHOFEL, W. H., 1924, The geology and invertebrate paleontology of the Comanchean and "Dakota" formations of Kansas: *Kansas Geol. Survey, Bull. 9*, pp. 1-135, pls. 1-23.

WAITE, H. A., 1942, Geology and ground-water resources of Ford county, Kansas: Kansas Geol. Survey, Bull. 43, pp. 1-250, figs. 1-22, pls. 1-16.

WING, M. E., 1930, The geology of Cloud and Republic counties, Kansas: Kansas Geol. Survey, Bull. 15, pp. 1-51, figs. 1, 2, pls. 1-18.

TRIASSIC

McLAUGHLIN, T. G., 1942, Geology and ground-water resources of Morton county, Kansas: Kansas Geol. Survey, Bull. 40, pp. 1-126, figs. 1-6, pls. 1-9.

PALEOZOIC

PERMIAN

ADAMS, G. I., GIRTY, G. H., and WHITE, DAVID, 1903, Stratigraphy and paleontology of the Upper Carboniferous rocks of the Kansas section: U.S. Geol. Survey, Bull. 211, pp. 1-123, pls. 1-4.

ADAMS, J. E., and others, 1939, Standard Permian section of North America: Am. Assoc. Petroleum Geologists Bull., vol. 23, no. 11, pp. 1673-1681.

BASS, N. W., 1929, The geology of Cowley county, Kansas: Kansas Geol. Survey, Bull. 12, pp. 1-203, figs. 1-23, pls. 1-12.

BEEDE, J. W., 1898, The stratigraphy of Shawnee county: Kansas Acad. Sci. Trans., vol. 15, pp. 27-34.

———, 1909, Formations of the Marion stage of the Kansas Permian: Kansas Acad. Sci. Trans., vol. 22, pp. 248-256.

BOOS, M. F., 1929, Stratigraphy and fauna of the Luta limestone (Permian) of Oklahoma and Kansas: Jour. Paleontology, vol. 3, no. 3, pp. 241-253, figs. 1-3, pl. 1.

BROADHEAD, G. C., 1884, Carboniferous rocks of eastern Kansas: St. Louis Acad. Sci. Trans., vol. 4, pt. 3, pp. 481-493.

CONDRA, G. E., 1927, The stratigraphy of the Pennsylvanian system in Nebraska: Nebraska Geol. Survey, 2d ser., Bull. 1, pp. 1-291, figs. 1-38, pls. 1-7.

CONDRA, G. E., and BENGSTON, N. A., 1915, The Pennsylvanian formations of southeastern Nebraska: Nebraska Acad. Sci., vol. 9, no. 2, pp. 1-60.

CONDRA, G. E., and BUSBY, C. E., 1933, The Grenola formation: Nebraska Geol. Survey, Paper 1, pp. 1-31, figs. 1, 2.

CONDRA, G. E., and UPP, J. E., 1931, Correlation of the Big Blue series in Nebraska: Nebraska Geol. Survey, 2d ser., Bull. 6, pp. 1-74, figs. 1-15.

CRAGIN, F. W., 1896, The Permian system in Kansas: Colorado College Studies, vol. 6, pp. 1-48.

DOTT, R. H., 1941, Regional stratigraphy of Mid-Continent: Am. Assoc. Petroleum Geologists Bull., vol. 25, no. 9, pp. 1619-1705, figs. 1-8.

ELIAS, M. K., 1934, Cycles of sedimentation in the Big Blue series of Kansas (abstract): Geol. Soc. America, Proc. 1933, p. 366.

———, 1937, Depth of deposition of the Big Blue (Late Paleozoic) sediments in Kansas: Geol. Soc. America Bull., vol. 48, no. 3, pp. 403-432, figs. 1-4, pl. 1.

FATH, A. E., 1921, Geology of the ElDorado oil and gas field: Kansas Geol. Survey, Bull. 7, pp. 1-187, figs. 1-9, pls. 1-19.

- HAY, ROBERT, 1891, *Kansas State Board Agr., 7th Biennial Rept.*, pp. 83-96.
- , 1893, *Kansas State Board Agr., 8th Biennial Rept.*, pp. 99-162, map.
- , 1896, *The geology of the Fort Riley Military Reservation and vicinity, Kansas: U.S. Geol. Survey, Bull. 137*, pp. 1-35, pls. 1-8.
- JEWETT, J. M., 1933, *Evidence of cyclic sedimentation in Kansas during the Permian period: Kansas Acad. Sci. Trans.*, vol. 36, pp. 137-140, figs. 1, 2.
- , 1941, *The geology of Riley and Geary counties, Kansas: Kansas Geol. Survey, Bull. 39*, pp. 1-164, figs. 1, 2, pls. 1-17.
- KIRK, M. Z., 1896, *A geologic section along the Neosho and Cottonwood rivers: Kansas Univ. Geol. Survey, vol. 1*, pp. 72-85.
- MEEK, F. B., and HAYDEN, F. V., 1860, *Geological explorations in Kansas Territory: Acad. Nat. Sci. Philadelphia Proc.*, vol. 11, pp. 8-75.
- MOORE, R. C., 1918, *The environment of Camp Funston: Kansas Geol. Survey, Bull. 4*, pp. 1-81, figs. 1-37, pls. 1-11.
- , 1940, *Carboniferous-Permian boundary: Am. Assoc. Petroleum Geologists Bull.*, vol. 24, no. 2, pp. 282-336, figs. 1-5.
- MOORE, R. C., ELIAS, M. K., and NEWELL, N. D., 1934, *Pennsylvanian and Permian rocks in Kansas: Kansas Geol. Survey, chart*.
- MOORE, R. C., and LANDES, K. K., 1937, *Geologic map of Kansas, scale 1:500,000: Kansas Geol. Survey*.
- MOORE, R. C., and MOSS, R. G., 1934, *Permian-Pennsylvanian boundary in the northern Midcontinent area (abstract): Geol. Soc. America, Proc. 1933*, p. 100.
- MOORE, R. C., and others, 1932, *Kansas Geol. Soc. Guidebook 6th Ann. Field Conference*, pp. 1-124, figs. and maps.
- , 1936, *Kansas Geol. Soc. Guidebook 10th Ann. Field Conference*, pp. 1-73, figs. 1-47.
- NORTON, G. H., 1939, *Permian redbeds of Kansas: Am. Assoc. Petroleum Geologists Bull.*, vol. 23, no. 12, pp. 1751-1819, figs. 1-24.
- PROSSER, C. S., 1894, *Kansas river section of the Permo-Carboniferous and Permian rocks of Kansas: Geol. Soc. America Bull.*, vol. 6, pp. 29-54.
- , 1895, *The classification of the upper Paleozoic rocks of central Kansas: Jour. Geology*, vol. 3, pp. 682-705, 764-800.
- , 1897, *The upper Permian and the lower Cretaceous: Kansas Univ. Geol. Survey, vol. 2*, pp. 54-194, figs. 2-9, pls. 9-24.
- , 1902, *Revised classification of the upper Paleozoic formations of Kansas: Jour. Geology*, vol. 10, pp. 700-737.
- , and BEEDE, J. W., 1904, *Description of the Cottonwood Falls quadrangle (Kansas): U.S. Geol. Survey, Geol. Atlas Cottonwood Falls folio (no. 109)*, pp. 1-6, 2 maps.
- SWALLOW, G. C., 1866, *Preliminary report: Kansas Geol. Survey*, pp. 1-198.

PENNSYLVANIAN

- ABERNATHY, G. E., 1937, *The Cherokee group of southeastern Kansas: Kansas Geol. Soc. Guidebook 11th Ann. Field Conference*, pp. 18-23, figs. 5, 6.

- ADAMS, G. I., 1896, A geologic section from Galena to Wellington, Kansas: Kansas Univ. Geol. Survey, vol. 1, pp. 16-30.
- , Girty, G. H., and White, David, 1903, Stratigraphy and paleontology of the Upper Carboniferous rocks of the Kansas section: U.S. Geol. Survey, Bull. 211, pp. 1-123, pls. 1-4.
- BEEDE, J. W., and ROGERS, A. F., 1908, Coal Measures faunal studies; faunal divisions of the Kansas Coal Measures: Kansas Univ. Geol. Survey, vol. 9, pp. 318-385, figs. 1-3, pl. 42.
- BENNETT, JOHN, 1896, Geologic section along the Missouri Pacific railway from the State line, Bourbon county, to Yates Center: Kansas Univ. Geol. Survey, vol. 1, pp. 86-98, fig. 4.
- BOWSHER, A. L., and JEWETT, J. M., 1943, Coal resources of the Douglas group in east-central Kansas: Kansas Geol. Survey, Bull. 46, pp. 1-94, figs. 1-12, pls. 1-6.
- CONDRA, G. E., 1927, The stratigraphy of the Pennsylvanian system in Nebraska: Nebraska Geol. Survey, 2d ser., Bull. 1, pp. 1-291, figs. 1-38, pls. 1-7.
- , and REED, E. C., 1943, The geological section of Nebraska: Nebraska Geol. Survey, Bull. 14, pp. 1-82, figs. 1-25.
- DOTT, R. H., 1941, Regional stratigraphy of Mid-Continent: Am. Assoc. Petroleum Geologists Bull., vol. 25, no. 9, pp. 1619-1705, figs. 1-8.
- HAWORTH, ERASMUS, 1896, Special report on coal: Kansas Univ. Geol. Survey, vol. 3, pp. 1-347, figs. 1-54, pls. 1-70.
- , and BENNETT, JOHN, 1908, Special report on oil and gas: Kansas Univ. Geol. Survey, vol. 9, pp. 42-121, pls. 4-40.
- JEWETT, J. M., 1932, Brief discussion of the Bronson group in Kansas: Kansas Geol. Soc. Guidebook 6th Ann. Field Conference, pp. 99-103.
- , 1937, Lateral changes in the lower Missouri beds of southeastern Kansas: Kansas Geol. Soc. Guidebook 11th Ann. Field Conference, pp. 35-37, figs. 11, 12.
- , 1940, Oil and gas in Linn county, Kansas: Kansas Geol. Survey, Bull., 30, pp. 1-29, figs. 1-7, pls. 1-3.
- , 1941, Classification of the Marmaton group, Pennsylvanian, in Kansas: Kansas Geol. Survey, Bull. 38, pt. 11, pp. 285-344, pls. 1-9.
- , and NEWELL, N. D., 1935, The geology of Wyandotte county, Kansas: Kansas Geol. Survey, Bull. 21, pt. 2, pp. 151-205, fig. 2, pls. 13-23.
- MOORE, R. C., 1931, Pennsylvanian cycles in the northern Mid-Continent region: Illinois Geol. Survey, Bull. 60, pp. 247-257, figs. 1-3.
- , 1932, A reclassification of the Pennsylvanian system in the northern Mid-Continent region: Kansas Geol. Soc. Guidebook 6th Ann. Field Conference, pp. 79-98, figs. 1-4.
- , 1935, Late Paleozoic crustal movements of Europe and North America: Am. Assoc. Petroleum Geologists Bull., vol. 19, no. 9, pp. 1253-1307, figs. 1-14.
- , 1936, Stratigraphic classification of the Pennsylvanian rocks of Kansas: Kansas Geol. Survey, Bull. 22, pp. 1-256, figs. 1-12.
- , 1936, "Carboniferous" rocks of North America: 16th Internat. Geol. Cong. Rept., vol. 1, pp. 593-617, figs. 1-8, pls. 1-3.

208 *Geological Survey of Kansas—1944 Reports of Studies*

- MOORE, R. C., and LANDES, K. K., 1937, Geologic map of Kansas, scale 1: 500,000: Kansas Geol. Survey.
- MOORE, R. C., and Others, 1932, Kansas Geol. Soc. Guidebook 6th Ann. Field Conference, pp. 1-124, figs. and maps.
- , 1936, Kansas Geol. Soc. Guidebook 10th Ann. Field Conference, pp. 1-73, figs. 1-47.
- , 1937, Kansas Geol. Soc. Guidebook 11th Ann. Field Conference, pp. 1-108, figs. 1-34, map.
- , 1944, Correlation of the Pennsylvanian formations of North America: Geol. Soc. America Bull., vol. 55, no. 6, pp. 657-706, pl. 1.
- NEWELL, N. D., 1935, Geology of Johnson and Miami counties, Kansas: Kansas Geol. Survey, Bull. 21, pt. 1, pp. 1-120, fig. 1, pls. 1-12.
- OAKES, M. C., and JEWETT, J. M., 1943, Upper Desmoinesian and lower Missourian rocks in northeastern Oklahoma and southeastern Kansas: Am. Assoc. Petroleum Geologists Bull., vol. 27, no. 5, pp. 632-640, fig. 1.
- PIERCE, W. G., and COURTIER, W. H., 1935, Englevalle channel sandstone of Pennsylvanian age, southeastern Kansas: Am. Assoc. Petroleum Geologists Bull., vol. 19, no. 7, pp. 1061-1068, figs. 1-3.
- , 1938, Geology and coal resources of the southeastern Kansas coal field: Kansas Geol. Survey, Bull. 24, pp. 1-122, figs. 1-13, pls. 1-12.

MISSISSIPPIAN

- FOWLER, G. M., 1935, Geology of the Tri-State district (abstract with discussion): Tulsa Geol. Soc. Digest, pp. 43-47.
- FOWLER, G. M., LYDEN, J. P., GREGORY, F. E., and AGAR, W. M., 1935, Chertification in the Tri-State (Oklahoma-Kansas-Missouri) mining district (with discussion): Am. Inst. Min. Met. Eng. Trans., vol. 115, Mining geology, pp. 106-163, figs. 1-32.
- LANDES, K. K., 1937, The Tri-State zinc-lead district: Kansas Geol. Soc. Guidebook 11th Ann. Field Conference, pp. 96-98.
- MOORE, R. C., FOWLER, G. M., and LYDEN, J. P., 1939, in Contributions to a knowledge of the lead and zinc deposits of the Mississippi valley region, edited by E. S. Bastin: Geol. Soc. America, Special Paper 24, pp. 1-12, pl. 1.
- MOORE, R. C., and LANDES, K. K., 1937, Geologic map of Kansas, scale 1: 500,000: Kansas Geol. Survey.

IGNEOUS ROCKS

- JEWETT, J. M., 1941, The geology of Riley and Geary counties, Kansas: Kansas Geol. Survey, Bull. 39, pp. 1-164 (note pp. 96-99), figs. 1-6, pls. 1-17.
- KNIGHT, G. L., and LANDES, K. K., 1932, Kansas laccoliths: Jour. Geology, vol. 40, no. 1, pp. 1-15, figs. 1-3.
- MOORE, R. C., and HAYNES, W. P., 1920, An outcrop of basic igneous rocks in Kansas: Am. Assoc. Petroleum Geologists Bull., vol. 4, no. 2, pp. 183-187.
- TWENHOFEL, W. H., 1926, Intrusive granite of the Rose dome, Woodson county, Kansas: Geol. Soc. America Bull., vol. 37, no. 2, pp. 403-412, figs. 1, 2.

INDEX OF STRATIGRAPHIC NAMES

(*Abandoned or unrecognized terms in italics.*)

Aarde shale, 176
Abilene conglomerate, 146
Admire group, 168
Aftonian interglacial deposits, 146
Alluvium, 145
Altamont limestone, 195
Amazonia limestone, 182
Americus limestone, 165, 168
Anna shale, 196
Ardmore limestone, 199
Argentine limestone, 187, 188
Aspinwall limestone, 169
Auburn shale, 175
Avant limestone, 189
Avoca limestone, 179

Bachelor Creek limestone, 176
Bader limestone, 166
Bandera Quarry sandstone, 196
Bandera shale, 194, 196
Barneston limestone, 164
"Bartlesville sand," 199
Beattie limestone, 166
Beecner Island shale, 151
Beil limestone, 180
Belleville formation, 146
Belvidere formation, 154
Bennett shale, 168
Bethany Falls limestone, 192
Bevier coal, 199
Big Blue series, 160
Bigelow formation, 165
Big Springs limestone, 180
Blackjack Creek limestone, 197
Blaine formation, 158
Block limestone, 191
Blue Hill shale, 152
Bluejacket sandstone, 199
Blue Rapids shale, 165
Blue Springs shale, 164
Bonner Springs shale, 187
Boone formation, 200
Bourbon shale, 193
Breezy Hill limestone, 197
Bridge Creek limestone, 153
Bronson group, 191
Brownville limestone, 170, 172
Burlingame limestone, 175
Burr limestone, 167

Calhoun shale, 178
Caneyville limestone, 172
Canville limestone, 192
Captain Creek limestone, 186
Carboniferous system, 170

Carlile shale, 152
Carlton limestone, 160
Cedar Bluff coal, 192
Cedar Hills sandstone, 158
Cedar Vale shale, 176
Cement City limestone, 190
Cenozoic rocks, 143, 145, 150, 159
Champion shell bed, 154
Chanute shale, 189
Chase group, 160
Checkerboard limestone, 194
Cherokee shale, 197
Cherryvale shale, 191
Chert gravels, 148
Chesterian series, 200
Cheyenne sandstone, 153, 154, 158
Chicago Mound formation, 169
Chikaskia member, 159
Church limestone, 176
Clay Creek limestone, 180
Coal Creek limestone, 177
Cockrum sandstone, 153
Codell sandstone, 152
Coffeyville formation, 192, 193
Colorado group, 152
Columbus coal, 199
Comanchean series, 154
Corbin City limestone, 190
Cottage Grove sandstone, 189
Cottonwood limestone, 165, 167
Council Grove group, 165
Cresswell limestone, 163
Cretaceous system, 144, 151, 155
Crisfield sandstone, 159
Critzler limestone, 193
Crouse limestone, 165
Croweburg coal, 199
"Cryptozoan limestone," 175
Curzon limestone, 178

Dakota formation, 153, 154
Dakota group, 153, 154
Day Creek dolomite, 157
Deer Creek limestone, 178
Dennis limestone, 191
Desmoinesian series, 194
?Dockum group, 156
Dodds Creek sandstone, 192
Dog Creek shale, 157, 158
Doniphan shale, 180
Douglas group, 182
Dover limestone, 170, 173
Doyle shale, 163
Drum limestone, 190

- Dry shale, 173
 Du Bois limestone, 177
 Dune sand, 145
- Easily Creek shale, 166
 Eiss limestone, 166
 Elgin sandstone, 180
 Elk River beds, 154
 Ellsworth formation, 153
 Elmdale shale, 165
 Elmo coal, 176
 Elmont limestone, 174
 Emma Creek formation, 148
 Emporia limestone, 174
 Englevale sandstone, 197
 Ervine Creek limestone, 179
 Eskridge shale, 165, 167
 Eudora shale, 186
- Fairport chalky shale, 152
 Falls City limestone, 169
 Farley limestone, 188
 "Fencepost limestone bed," 153
 Five Point limestone, 169
 Fleming coal, 199
 Florena shale, 166
 Florence limestone, 164
 Flowerpot shale, 158
 Fontana shale, 191
 Foraker limestone, 168
 Fort Hays limestone, 152
 Fort Riley limestone, 164
 Fort Scott limestone, 197
 French Creek shale, 173
 Friedrich shale, 173
 Frisbie limestone, 188
 Funston limestone, 165
- Gage shale, 163
 Galesburg shale, 192
 Garrison shale, 165, 166
 Gerlane formation, 146
 Glenrock limestone, 168
 Grandhaven limestone, 173
 Graneros shale, 153
 Grant shale, 163
 Grayhorse limestone, 172
 Greenhorn limestone, 152
 Greenleaf sandstone, 154
 Grenola limestone, 167
 Guadalupian series, 156
 Gulfian series, 151
 Hamlin shale, 169
 Happy Hollow limestone, 176
 Harper sandstone, 159
 Hartford limestone, 178
 Hartland shale, 153
 Harveyville shale, 174
 Haskell limestone, 183
- Haskew gypsum, 158
 Havensville shale, 165
 Hawxby shale, 169
 Heebner shale, 181
 Henrietta group, 194
 Hepler sandstone, 194
 Herington limestone, 154, 160
 Hertha limestone, 187, 191, 193, 194
 Heumader shale, 181
 Hickory Creek shale, 187
 Higginsville limestone, 197
 Hogshooter limestone, 191
 Hollenberg limestone, 160
 Holmesville shale, 164
 Holt shale, 177
 Hooser shale, 166
 Houchen Creek limestone, 169
 Houx limestone, 197
 Howard limestone, 176
 Howe limestone, 167
 Hughes Creek shale, 168
 Hushpuckney shale, 192
 Hutchinson salt, 160
- Iatan limestone, 184
 Idenbro limestone, 195
 Igneous rocks, 144, 200
 Indian Cave sandstone, 170
 Iola limestone, 189
 Iowa Point shale, 178
 Ireland sandstone, 182
 Island Creek shale, 188
- Jackson Park shale, 181
 Janssen clay, 153
 Jetmore chalk, 153
 Jim Creek limestone, 173
 Johnson shale, 168
 Jones Point shale, 177
 Jurassic, 151, 154
- Kansan till, 146, 148
 Kansas City formation, 187, 191
 Kansas City group, 187
 Kansas City oolite, 190
 Kanwaka shale, 180
 Keokuk limestone, 200
 Kereford limestone, 181
 King Hill shale, 180
 Kingman sandstone, 159
 Kingsdown silt, 147
 Kinney limestone, 164
 Kiowa shale, 153, 154
 Knobtown sandstone, 193
 Krider limestone, 163
- Laberdie limestone, 196
 Labette shale, 196
 Ladore shale, 193
 Lake Creek shale, 151

- Lake Neosho shale, 195
 Lane-Bonner Springs shale, 188
 Lane shale, 189
 Langdon shale, 173
Lanphier beds, 154
 Lansing group, 186
 Larsh-Burroak shale, 179
 Laverne formation, 149
 Lawrence shale, 182
 Leavenworth limestone, 181
 Lecompton limestone, 179
 Legion shale, 167
 Lenapah limestone, 195
 Leonardian series, 157
 Lincoln limestone, 153
 Little Cabin sandstone, 199
Little Kaw limestone, 186
 Little Osage shale, 197
 Loess, 145, 146
 Long Creek limestone, 168
Luta limestone, 163

 McAlester formation, 199
 McPherson formation, 146
 Maple Hill limestone, 173
 Marlow sandstone, 157
 Marmaton group, 194
Marquette sandstone, 153
 Matfield shale, 164
 Meade formation, 147
Medicine bed, 154
 Medicine Lodge gypsum, 158
 Memorial shale, 194
Mentor beds, 154
 Meramecian series, 200
 Merriam limestone, 187
 Mesozoic rocks, 143, 151, 156
 Middleburg limestone, 166
 Middle Creek limestone, 193
 Milan limestone, 160
 Mine Creek shale, 196
 Mineral coal, 199
 Mississippian system, 200
 Missourian series, 184, 200
 Montana group, 151
 Morrill limestone, 166
 Mound City shale, 193
 Mulberry coal, 196
 Mulky coal, 197
 Muncie Creek shale, 189
 Myrick Station limestone, 196

Natural Corral sandstone, 154
 Nebraska City limestone, 172
 Nebraskan till, 146, 148
 Nescatunga gypsum, 158
 Neva limestone, 165, 167
 Ninnescah shale, 159
 Niobrara chalk, 152
 Nippewalla group, 157

 Nodaway coal, 176
 Nolans limestone, 160
 Norfleet limestone, 195
 Nowata shale, 195
 Noxie sandstone, 190
 Nyman coal, 173

 Oaks shale, 169
 Ochelata group, 187
 Odell shale, 163
 Ogallala formation, 149
 Oketo shale, 164
Olathe limestone, 186
 Oread limestone, 181
 Osagian series, 200
 Oskaloosa shale, 179
 Ost limestone, 179
 Ozawkie limestone, 179

 Paddock shale, 163
 Paleozoic rocks, 143, 156
 Paola limestone, 189
 Pawhuska formation, 177
 Pawnee limestone, 196
 Pedee group, 184
 Pennsylvanian system, 170, 185, 198, 201
 Permian system, 143, 156, 161, 162, 171
 Perry Farm shale, 195
 "Peru sand," 197
 Pfeifer shale, 153
 Pierre shale, 151
Pierson Point shale, 174
 "Pilot" coal, 199
 Plattsburg limestone, 187
 Plattsmouth limestone, 181
 Pleistocene series, 145, 146, 148
 Pliocene series, 148
 Pony Creek shale, 172

 Quartermaster formation, 156, 157
 Quartermaster group, 156
 Quaternary system, 145
 Queen Hill shale, 180
 Quindaro shale, 188
 Quivira shale, 190

 Raytown limestone, 189
 Reading limestone, 174
 Recent series, 145
 Red Eagle limestone, 167
Reeder sandstone, 153
 Relay Creek? dolomite and sandstone, 157
 Rexroad member, 149
 Riverton coal, 199
 Robbins shale, 183
 Roca shale, 167
 Rock Bluff limestone, 179
 Rock Lake shale, 186
Rocktown channel sandstone, 153
 Rulo limestone, 175
 Runnymede sandstone, 159

Salem Point shale, 167
 Sallyards limestone, 167
 Salt Creek gravel beds, 146
 Salt Grass shale, 151
 Salt Plain formation, 159
 Sanborn formation, 147
 Schroyer limestone, 165
Scranton shale, 175
 Severy shale, 176
 Sharon Springs shale, 152
 Shawnee group, 177
 Sheldon limestone, 177
 Shimer gypsum, 158
 Sibley coal, 183
 Silver Lake shale, 175
 Smoky Hill chalk, 152
 Sniabar limestone, 193
 Snyderville shale, 182
 Soldier Creek shale, 175
Solomon formation, 153
 South Bend limestone, 186
 Speiser shale, 165
 Spring Branch limestone, 180
Spring Creek clay, 153
 Spring Hill limestone, 187
 "Squirrel sand," 199
 Stanton limestone, 184, 186
 Stark shale, 192
 Stearns shale, 166
 Stine shale, 169
Stokes sandstone, 154
 Stonebreaker limestone, 174
 Stone Corral dolomite, 159
 Stoner limestone, 186
 Stovall limestone, 163
 Stranger formation, 183
 Stull shale, 180
 Summit coal, 197
 Sumner group, 159
 Swope limestone, 192

Table Creek shale, 173
 Taloga formation, 157
 Tarkio limestone, 174
 Tecumseh shale, 179
 Terrace deposits, 146, 147
 Terra Cotta clay, 153
 Tertiary system, 143, 148
 Thayer coal, 189

Threemile limestone, 165
 Tina limestone, 196
 Tonganoxie sandstone, 183
 Topeka limestone, 177
 Toronto limestone, 182
 Towanda limestone, 163
 Towle shale, 170
 Triassic system, 151, 154
 Turner Creek shale, 177

 Utopia limestone, 176

 Verdigris limestone, 199
Victory Junction shale, 186
 Vilas shale, 187
 Vinland shale, 183
 Virgilian series, 170

 Wabaunsee group, 170
 Wakarusa limestone, 175
 Walter Johnson sandstone, 195
 Wamego shale, 174
 Warner sandstone, 199
 Warrensburg sandstone, 197
 Warsaw limestone, 200
 "Wayside sand," 195
 Wea shale, 191
 Wea-Quivira shale, 191
 Weir-Pittsburg coal, 199
 Wellington formation, 160
 Weskran shale, 152
 West Branch shale, 169
 Westerville limestone, 190
 Weston shale, 184
 Westphalia limestone, 183
 White Cloud shale, 176
 Whitehorse sandstone, 157
 Willard shale, 174
 Williamsburg coal, 182
 Winfield limestone, 163
 Winterset limestone, 187, 191
 Winzeler shale, 176
 Wolfcampian series, 160, 200
 Wolf River limestone, 178
 Woodhouse clay, 149
 Woods Siding formation, 172
 Worland limestone, 195
 Wreford limestone, 165
 Wyandotte limestone, 188
 Wymore shale, 164

STATE GEOLOGICAL SURVEY OF KANSAS

Recent Publications

- BULLETIN 32.** Coal Resources of Kansas; Post-Cherokee Deposits, by R. E. Whitla, 64 pages, 1940. Mailing charge, 20 cents.
- BULLETIN 33.** Subsurface Mississippian Rocks of Kansas, by Wallace Lee, 114 pages, 1940. Mailing charges, 25 cents.
- BULLETIN 34.** Geologic Studies in Southwestern Kansas, by H. T. U. Smith, 244 pages, 1940. Mailing charge, 25 cents.
- BULLETIN 36.** Exploration for Oil and Gas in Western Kansas during 1940, by Walter A. Ver Wiebe, 109 pages, 1941. Mailing charge, 25 cents.
- BULLETIN 37.** Geology and Ground-Water Resources of Stanton County, Kansas, by Bruce F. Latta, 119 pages, 1941. Mailing charge, 25 cents.
- BULLETIN 38.** 1941 Reports of Studies (Parts issued separately). Mailing charge, 10 cents each part.
- BULLETIN 39.** The Geology of Riley and Geary Counties, Kansas, by J. M. Jewett, 164 pages, 1941. Mailing charge, 25 cents.
- BULLETIN 40.** Geology and Ground-Water Resources of Morton County, Kansas, by Thad G. McLaughlin, 126 pages, 1942. Mailing charge, 25 cents.
- BULLETIN 41.** 1942 Reports of Studies (Ten parts issued separately). Mailing charge, 10 cents each part.
- BULLETIN 42.** Exploration for Oil and Gas in Western Kansas During 1941, by Walter A. Ver Wiebe, 123 pages, 1942. Mailing charge, 25 cents.
- BULLETIN 43.** Geology and Ground-Water Resources of Ford County, Kansas, by Herbert A. Waite, 250 pages, 1942. Mailing charge, 25 cents.
- BULLETIN 44.** Geophysical Investigations in the Tri-State Zinc and Lead Mining District, by J. J. Jakosky, R. M. Droyer, and C. H. Wilson, 151 pages, 1942. Mailing charge, 25 cents.
- BULLETIN 45.** Geology and Ground-Water Resources of Meade County, Kansas, by John C. Frye, 152 pages, 1942. Mailing charge, 25 cents.
- BULLETIN 46.** Coal Resources of the Douglas Group in East-Central Kansas, by Arthur L. Bowsher and John M. Jewett, 94 pages, 1943. Mailing charge, 25 cents.
- BULLETIN 48.** Exploration for Oil and Gas in Western Kansas During 1942, by Walter A. Ver Wiebe, 88 pages, 1943. Mailing charge, 25 cents.
- BULLETIN 49.** Geology and Ground-Water Resources of Hamilton and Kearny Counties, Kansas, by Thad G. McLaughlin, 220 pages, 1943. Mailing charge, 25 cents.
- BULLETIN 50.** Ground Water in the Oil-Field Areas of Ellis and Russell Counties, Kansas, by John C. Frye and James J. Brazil, 104 pages, 1943. Mailing charge, 25 cents.
- BULLETIN 51.** The Stratigraphy and Structural Development of the Forest City Basin in Kansas, by Wallace Lee, 142 pages, 1943. Mailing charge, 25 cents.
- BULLETIN 53.** McLouth Gas and Oil Field, Jefferson and Leavenworth Counties, Kansas, by Wallace Lee and Thomas G. Payne, 193 pages, 1944. Mailing charge, 25 cents.
- BULLETIN 54.** Exploration for Oil and Gas in Western Kansas During 1943, by Walter A. Ver Wiebe, 104 pages, 1944. Mailing charge 25 cents.
- BULLETIN 55.** Geology and Ground-Water Resources of Finney and Gray Counties, Kansas, by Bruce F. Latta, 272 pages, 1944. Mailing charge, 25 cents.
- RESOURCE-FULL KANSAS,** Kenneth K. Landes and Oren R. Bingham, 65 pages. Mailing charge, 10 cents.

STATE GEOLOGICAL SURVEY OF KANSAS

BULLETIN 47

1943 REPORTS OF STUDIES

- PART 1. NEW CARBONIFEROUS AND PERMIAN SPONGES**, by Ralph H. King, pp. 1-36, pls. 1-3, figs. 1-2, February 27, 1943.
- PART 2. KANSAS OIL FIELD BRINES AND THEIR MAGNESIUM CONTENT**, by Walter H. Schoewe, pp. 37-76, figs. 1-3, June 30, 1943.
- PART 3. DEEP WATER WELL AT THE JAYHAWK ORDNANCE WORKS IN CHEROKEE COUNTY KANSAS**, by G. E. Abernathy, pp. 77-112, figs. 1-4, September 10, 1943.
- PART 4. A PROCESS FOR EXTRACTING ALUMINA FROM KANSAS CLAY**, by E. D. Kinney, pp. 113-136, December 10, 1943.

BULLETIN 52

1944 REPORTS OF STUDIES

- PART 1. RECONNAISSANCE OF PLEISTOCENE DEPOSITS IN NORTH-CENTRAL KANSAS**, by Claude W. Hibbard, John C. Frye, and A. Byron Leonard, pp. 1-28, pls. 1, 2, figs. 1, 2, February 20, 1944.
- PART 2. GROUND-WATER CONDITIONS IN THE NEOSHO RIVER VALLEY IN THE VICINITY OF PARSONS, KANSAS**, by Charles C. Williams, pp. 29-80, pls. 1-3, figs. 1-9, March 15, 1944.
- PART 3. COAL RESOURCES OF THE KANSAS CITY GROUP, THAYER BED, IN EASTERN KANSAS**, by Walter H. Schoewe, pp. 81-136, pls. 1-5, figs. 1-8, July 25, 1944.
- PART 4. TABULAR DESCRIPTION OF OUTCROPPING ROCKS IN KANSAS**, by Raymond C. Moore, John C. Frye, and John Mark Jewett, pp. 137-212, figs. 1-9, October 25, 1944.
- PART 5. MINED AREAS OF THE WEIR-PITTSBURG COAL BED**, by G. E. Abernathy, pp. 213-228, pl. 1, fig. 1, October 25, 1944.