



# Cannibalism in Juvenile Tropical House Geckos, *Hemidactylus mabouia* (Squamata: Gekkonidae)

Daylon Fundora Caballero<sup>1</sup> and Llilian Martínez Pérez<sup>2</sup>

<sup>1</sup>Delegación Territorial del CITMA, Subdelegación de Medio Ambiente, Ciego de Ávila 65200, Cuba (daylon8910@nauta.cu)

<sup>2</sup>Centro de Estudios del Jardín Botánico de Villa Clara, Universidad Central “Marta Abreu” de Las Villas, Villa Clara 54830, Cuba (llilian@uclv.cu)

**C**annibalism among Tropical House Geckos (*Hemidactylus mabouia*) has been documented (Zamprogno and Teixeira 1998; Pombal and Pombal 2010; Rufino de Albuquerque et al. 2013) but only Lyakurwa (2017) described such behavior and cannibalism in juveniles has not been reported. Herein we present an observation of cannibalism in two juvenile Tropical House Geckos and the first documented report of cannibalism in the species in Cuba.



**Fig.1.** A juvenile Tropical House Gecko (*Hemidactylus mabouia*) preying on a smaller conspecific in Santa Clara, Villa Clara, Cuba. Photograph by Llilian Martínez Pérez.

At 2000 h on 30 June 2019 on a wall near a lamp in a house in urban Santa Clara, Villa Clara, central Cuba (22.3927°S, 79.9484°W; WGS 84), we observed a juvenile *H. mabouia* (SVL approx. 4 cm) preying on a smaller conspecific (SVL approx. 2 cm). The larger gecko had a firm grasp on the head and forelimbs of its prey (Fig. 1). We were unable to verify if ingestion was successful because the predatory gecko escaped into a crack in the wall.

## Acknowledgement

Tomás M. Rodríguez-Cabrera (Sociedad Cubana de Zoología) provided helpful assistance with this manuscript.

## Literature Cited

- Lyakurwa, J.V. 2017. *Hemidactylus mabouia* (Tropical House Gecko). Cannibalism. *Herpetological Review* 48: 646.
- Pombal, R. and J.P. Pombal. 2010. *Hemidactylus mabouia* (Tropical House Gecko). Cannibalism. *Herpetological Review* 41: 223–224.
- Rufino de Albuquerque, N., A. da Silva Costa-Urquiza, M. Pereira Soares, L. Saucedo Alves, and M.V. Santiago Urquiza. 2013. Diet of two sit-and-wait lizards, *Phyllopezus pollicaris* (Spix, 1825) (Phyllodactylidae) and *Hemidactylus mabouia* (Moreau de Jonnès, 1818) (Gekkonidae) in a perianthropic area of Mato Grosso do Sul, western Brasil. *Biotá Neotropical* 13: 376–381. <http://dx.doi.org/10.1590/S1676-06032013000400032>.
- Zamprogno, C. and R.L. Teixeira. 1998. Hábitos alimentares da lagartixa de parede *Hemidactylus mabouia* (Reptilia, Gekkonidae) da planície litorânea do norte do Espírito Santo, Brasil. *Revista Brasileira de Biologia* 58: 143–150.